



## **Morphological and Biochemical Bases of Resistance of Some Groundnut Germplasms against Tobacco Caterpillar, *Spodoptera litura* (Fabricius) and Leaf Miner, *Approaerema modicella* (Deventer)**

**Shaik Moizur Rahman <sup>a\*</sup>, K Vijayalakshmi <sup>b</sup>, Ch. V. Durga Rani <sup>c</sup>,  
S. Ameer Basha <sup>d</sup> and C. Srinivas <sup>a</sup>**

<sup>a</sup> Department of Entomology, College of Agriculture, Rajendranagar, PJTSAU, Hyderabad-30, India.

<sup>b</sup> Department of Entomology, Agriculture College, Palem, PJTSAU-509215, India.

<sup>c</sup> Department of Genetics and Plant Breeding, Institute of Biotechnology, PJTSAU, Hyderabad-30, India.

<sup>d</sup> Department of Plant Pathology, College of Agriculture, Rajendranagar, PJTSAU, Hyderabad-30, India.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/IJECC/2021/v11i1230571

Editor(s):

(1) Dr. Daniele De Wrachien, State University of Milan, Italy.

Reviewers:

(1) V. N. Kapadia, AAU, India.

(2) Esameldin Bashir Mohamed Kabbashi, The National Food Research Centre, Sudan.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:

<https://www.sdiarticle5.com/review-history/80664>

**Original Research Article**

**Received 07 October 2021**

**Accepted 15 December 2021**

**Published 16 December 2021**

### **ABSTRACT**

A screening experiment of forty germplasm lines was conducted at College Farm, College of Agriculture, Rajendranagar, Hyderabad in *rabi*, 2019-20 and was laid in Randomized block design (RBD) having two replicates. Three germplasm lines *viz.*, ICGV 16679 (13.71% leaf damage), ICGV 07222 (14.00% leaf damage) and ICGV 93468 (14.25% leaf damage) showed greater resistance than the resistant check, ICGV 86031 (15.04% leaf damage) against tobacco caterpillar and with respect to resistance against leaf miner, no germplasm line was found to be superior than the resistant check, ICGV 86031 (7.82% leaf damage). However, the germplasm line, ICGV 02266 (8.35% leaf damage) was next best to the resistant check. The various morphological and

biochemical characters were recorded and correlated with insect pest damage and incidence to know their role in imparting resistance/susceptibility. Resistance to various insect pests in germplasm lines was due to significantly higher trichome density on leaf lamina, higher phenol and tannin content.

**Keywords:** Screening; tobacco caterpillar; leaf miner; groundnut; biochemical; morphological & resistance.

## 1. INTRODUCTION

Groundnut (*Arachis hypogaea* L.) is an important leguminous food crop in India and is also known as peanut, earthnut, monkey nut, and goobers [1]. It has originated in South America, where the genus *Arachis* is widely distributed. It is cultivated mostly in the semi-arid tropical and sub-tropical regions [2]. Low productivity in groundnut is attributed to several constraints and the attack of insect-pests is one among these. The avoidable yield loss due to major insect pests of groundnut was recorded to be 48.57 percent in pod and 42.11 percent in fodder [3].

Host plant resistance is an effective biological approach for plant protection [4] and using insect resistant varieties is an important strategy of integrated pest management [5]. The biophysical traits can be used as phenotypic markers to identify groundnut germplasm lines with resistance to tobacco caterpillar and leaf miner. The main reasons of variability in resistance shown by different genotypes were explained by Painter [6]. He pointed out three mechanisms of resistance, viz., non-preference (antixenosis), antibiosis and tolerance.

The morphological features of plants are associated with attraction, feeding and egg laying of the insect pests [7]. The identification of important morphological and bio-chemical characteristics of germplasm lines will help to understand the resistance mechanisms of plant against insect pests which in turn can be used in the breeding programmes for development of resistant varieties to these insect pests.

## 2. MATERIALS AND METHODS

The present investigations were conducted at College Farm, College of Agriculture, Rajendranagar, Hyderabad (Telangana) under field conditions during *rabi*, 2019-20. Geographic location of Hyderabad pertains to 17.3850° North latitude, 78.4867° East longitude and elevation of 536 metres above mean sea level (MSL).

## 2.1 Experimental Layout

The experiment was laid out in a randomized block design (RBD) with 40 treatments (Plate 1), each replicated twice. The plot size was 225 m<sup>2</sup>. Each treatment was sown in two rows of 3 m each with row to row distance of 30 cm and plant to plant spacing of 10 cm along with susceptible (ICGV 91114) and resistant (ICGV 86031) check. The crop was raised as per the package of practices recommended by PJTSAU (Vyavasaya Panchangam) [8] except for the plant protection measures.

## 2.2 Observations

### 2.2.1 Tobacco caterpillar

The larval counts were recorded on five randomly selected plants in each line under each replication and also percent leaf area damaged was worked out by assessing leaf damage at top, middle and bottom leaves from ten randomly selected plants showing maximum damage due to tobacco caterpillar at 10 days interval. By considering the mean percent leaf area damage, the germplasm was given damage rating by following the scale adopted by Sharma et al. [2].

### 2.2.2 Leaf miner

Observations on leaf miner were made on top five leaves from ten randomly selected plants in each replication for the number of leaflets damaged by leaf miner and percent leaf damage was worked out. (AICRP on groundnut [9]; Chakravarthy and Selvanarayanan [10]).

The morphological that include plant height, no. of branches, main stem thickness, no. of trichomes on leaf lamina, midrib and petiole and biochemical parameters were estimated by following standard procedures as prescribed by Hedge and Hofreiter [11] (Total sugars), AOAC [12] and Mariotti *et al.* [13] (proteins); Sadasivam and Manickam [14] (phenols) and Schanderl [15] (tannins). These parameters were correlated with




Plate 1. List of germplasm lines

tobacco caterpillar and leaf miner incidence and infestation to study their relationship.

### 3. RESULTS AND DISCUSSION

#### 3.1 Incidence and Infestation of Tobacco Caterpillar

The mean no. of larvae per plant on all the germplasm lines were on par with the resistant check, ICGV 86031 (1.31) except five lines viz., ICGV 00351 (1.45), K 6 (1.47), ICGV 13189 (1.53), ICGV 10021 (1.55) and JCG 5834 (1.62). Similarly, considering the mean per cent leaf damage per plant, the germplasm lines ICGV 07222, ICGV 93468 and ICGV 02266 recorded 14.00, 14.25 and 16.96 percent leaf damage per plant, respectively and were on par with the resistant check, ICGV 86031 (15.04%) and are significantly different from the susceptible check, ICGV 91114 (34.50%) (Table 1).

#### 3.2 Infestation of Groundnut Leaf Miner

The overall mean leaf damage due to the leaf miner in test lines ranged from 7.82 to 18.15 percent. The resistant check, ICGV 86031 recorded the lowest per cent damage of 7.82 percent and it was on par with the germplasm lines viz., ICGV 15083, ICGV 181052, ICGV 181011, ICGV 171015, ICGV 13189, ICGV 15426, ICGV 99195, ICGV 93468, ICGV 00298, ICGV 00350, ICGV 00351, ICGV 06040, ICGV 02266, ICGV 10021, ICGV SM 90704 and KDG 128 (8.35-11.85%). The highest leaf damage was recorded in JCG 3341 (18.40%) (Table 1).

#### 3.3 Morphological Parameters of Groundnut Germplasm Lines

Various morphological parameters viz., plant height, number of branches per plant, trichomes on leaf lamina, leaf midrib and petiole, main stem thickness were recorded and correlated with insect pests to know the morphological bases of resistance in the test germplasm lines.

The plant height, no. of branches and main stem thickness ranged between 12.68 cm (ICGV 15083) to 27.90 cm (ICGV 91114), 4.20 (K 6) to 8.00 (ICGV 06424) and 2.07 cm (ICGV 06040) to 3.33 (ICGV 86031), respectively with a mean of 19.82 cm, 5.35 and 2.49 cm, respectively. The no. of trichome on leaf lamina, midrib and petiole per 0.25 cm<sup>2</sup> ranged between 25.68 (K 6) to 54.56 (ICGV 171015), 38.25 (ICGV 06424) to

96.01 (ICGV 86031) and 52.10 (ICGV 06424) to 102.87 (ICGV 86031), respectively with a mean of 36.03, 60.65 and 72.47, respectively (Table 1).

#### 3.3.1 Tobacco caterpillar incidence and infestation

The correlation studies indicated significant positive correlation between tobacco caterpillar larval population (no./plant) and percent leaf damage with plant height whereas significant negative relationship with no. of branches per plant, main stem thickness, trichome density on lamina, trichome density on midrib and trichome density on petiole (Table 2).

Krishnaiah [16] reported that plant height had positive correlation with *S. litura* damage in groundnut which corroborates with our studies. Similar to our studies, Sharma et al. [2] reported significant negative relationship between *S. litura* damage and hairiness and main stem thickness of the groundnut varieties screened. The present findings are in line with Mohammad Saleem et al. [17] who reported negative significant correlation between damage due to *S. litura* and trichomes on midrib and leaf lamina indicating the role of trichomes in imparting resistance against *S. litura*. Near to midrib of the leaf, *S. litura* prefers to lay eggs. So, midrib trichome density hinders in the oviposition of the insect, thus imparting resistance to *S. litura*.

#### 3.3.2 Leaf miner infestation

Positive non-significant correlation was observed between plant height and damage due to mining by leaf miner. The correlation studies indicated negative relationship between no. of branches per plant (non-significant), main stem thickness (non-significant), trichome density on lamina (significant), trichome density on midrib (non-significant) and trichome density on petiole (non-significant) and damage due to mining (Table 2).

The present findings are in conformity with the results of Ranga Rao [18] and Vishalakshi [19] who recorded significantly greater number of trichomes on midrib as well as on leaf lamina in resistant genotypes to leaf miner. Peeru Saheb et al. [20] reported that the trichomes had shown a negative correlation with fifth instar duration of groundnut leaf bud borer. On contrary to our findings, Sharma et al. [2] reported positive correlation between main stem thickness and trichome density on leaf lamina to *Spodoptera*.

**Table 1. Incidence and infestation of tobacco caterpillar, leaf miner and morphological characters of groundnut germplasm lines**

Genotype	Tobacco caterpillar		Leaf miner	Plant height (cm)	No. of branches	Main stem thickness (cm)	Trichome density (No. of trichomes / 0.25 cm <sup>2</sup> )		
	Mean no. of larvae/plant <sup>a</sup>	Mean leaf damage/plant (%) <sup>b</sup>	Mean leaf damage/plant (%) <sup>b</sup>				Leaf lamina	Midrib	Petiole
ICGV 15083	0.88 (1.37)	19.92 (26.42)	10.30 (18.42)	12.68	7.50	2.55	34.25	54.15	63.55
ICGV 181052	1.33 (1.53)	26.38 (30.79)	9.70 (17.60)	15.83	7.40	2.25	30.96	55.13	65.78
ICGV 181011	1.17 (1.47)	24.96 (29.93)	10.05 (18.15)	16.73	5.80	2.43	35.25	66.33	76.00
ICGV 171015	1.03 (1.42)	22.33 (28.15)	9.90 (18.09)	26.60	5.80	2.51	54.56	60.01	59.37
ICGV 16679	0.65 (1.28)	13.71 (21.67)	17.55 (24.58)	18.00	5.60	2.88	34.85	68.59	76.60
ICGV 03043	1.07 (1.44)	22.71 (28.37)	16.20 (23.58)	21.70	5.60	2.45	40.85	63.66	78.86
ICGV 07222	0.83 (1.35)	14.00 (21.92)	14.20 (22.05)	16.82	7.80	2.58	44.65	73.22	80.55
ICGV 06424	0.97 (1.40)	19.50 (26.04)	15.20 (22.74)	21.75	8.00	2.54	31.95	38.25	52.10
ICGV 13189	1.53 (1.59)	32.83 (34.90)	11.40 (19.52)	24.10	4.80	2.22	30.89	62.55	68.55
ICGV 13200	1.35 (1.53)	29.50 (32.85)	14.40 (22.04)	18.30	4.30	2.31	27.12	56.96	62.58
ICGV 14421	1.13 (1.46)	24.13 (29.32)	13.45 (21.23)	16.31	6.00	2.28	33.88	59.65	73.99
ICGV 15423	1.57 (1.60)	31.50 (34.08)	13.05 (21.03)	22.88	5.80	2.19	27.67	48.54	64.20
ICGV 15426	1.18 (1.48)	24.58 (29.66)	9.60 (17.83)	18.15	4.60	2.42	31.39	48.56	65.23
ICGV 93468	0.75 (1.32)	14.25 (22.12)	11.85 (20.03)	13.15	4.50	2.94	47.56	68.26	84.75

Genotype	Tobacco caterpillar		Leaf miner	Plant height (cm)	No. of branches	Main stem thickness (cm)	Trichome density (No. of trichomes / 0.25 cm <sup>2</sup> )		
	Mean no. of larvae/ plant <sup>a</sup>	Mean leaf damage/ plant (%) <sup>b</sup>	Mean leaf damage/ plant (%) <sup>b</sup>				Leaf lamina	Midrib	Petiole
ICGV 99195	1.00 (1.41)	21.25 (27.41)	10.95 (19.15)	22.70	5.60	2.54	47.35	66.89	71.64
ICGV 00298	1.18 (1.48)	25.46 (30.27)	10.90 (19.10)	21.89	4.80	2.43	30.93	61.50	70.25
ICGV 00350	1.23 (1.49)	26.66 (31.04)	11.45 (19.73)	16.43	5.00	2.38	33.96	60.36	68.95
ICGV 00351	1.45 (1.57)	29.08 (32.60)	9.80 (18.29)	16.57	5.20	2.31	38.23	44.15	61.96
ICGV 06040	1.30 (1.52)	28.58 (32.26)	9.20 (17.46)	18.71	5.00	2.07	34.37	48.70	62.58
ICGV 02266	0.77 (1.33)	16.96 (24.26)	8.35 (16.45)	14.97	5.00	3.12	45.66	78.55	83.55
ICGV 86015	1.27 (1.51)	25.50 (30.17)	14.10 (21.81)	18.18	4.80	2.45	27.57	62.55	94.84
ICGV 93437	1.27 (1.51)	26.92 (31.15)	17.00 (24.04)	21.25	7.50	2.47	34.25	40.66	57.97
ICGV 93382	1.13 (1.46)	23.83 (29.15)	12.70 (20.81)	20.85	7.00	2.55	31.17	66.95	73.84
ICGV 10001	1.22 (1.49)	28.63 (32.23)	14.35 (21.87)	26.16	5.50	2.36	29.54	53.98	69.31
ICGV 10021	1.55 (1.60)	34.92 (36.19)	11.50 (19.72)	22.90	4.20	2.35	31.56	40.56	61.11
ICGV 15264	1.42 (1.56)	33.33 (35.22)	18.15 (25.13)	19.65	4.50	2.28	32.25	63.56	77.17
ICGV 15307	1.15 (1.47)	26.96 (31.24)	15.30 (22.65)	18.40	4.40	2.51	29.58	55.89	66.31
ICGV 87141	1.37 (1.54)	32.21 (34.54)	15.15 (22.80)	21.90	4.60	2.29	41.56	51.65	71.38

Genotype	Tobacco caterpillar		Leaf miner	Plant height (cm)	No. of branches	Main stem thickness (cm)	Trichome density (No. of trichomes / 0.25 cm <sup>2</sup> )		
	Mean no. of larvae/ plant <sup>a</sup>	Mean leaf damage/ plant (%) <sup>b</sup>	Mean leaf damage/ plant (%) <sup>b</sup>				Leaf lamina	Midrib	Petiole
ICGV SM 90704	1.15 (1.47)	25.71 (30.41)	10.35 (18.53)	20.00	5.40	2.18	36.25	70.65	86.17
ICGV 90320	0.90 (1.38)	21.75 (27.77)	12.35 (20.44)	21.40	4.50	2.56	38.65	72.68	87.31
JCG 4798	1.43 (1.56)	31.67 (34.22)	12.80 (20.57)	19.45	4.30	2.34	42.56	53.65	61.91
JCG 5834	1.62 (1.62)	37.46 (37.69)	14.35 (22.05)	25.60	4.30	2.43	36.90	66.95	80.33
JCG 2141	1.02 (1.42)	21.83 (27.76)	12.45 (20.58)	20.45	5.20	2.49	43.65	59.69	66.85
JCG 3341	1.23 (1.49)	28.08 (31.96)	18.40 (25.22)	20.65	5.20	2.33	30.65	66.94	87.97
K 6	1.47 (1.57)	35.33 (36.44)	15.95 (23.20)	23.65	4.20	2.37	25.68	50.65	56.87
K 9	1.13 (1.45)	22.75 (28.38)	14.45 (22.17)	18.10	5.60	2.45	44.25	71.25	81.56
KDG 128	0.98 (1.40)	21.54 (27.54)	9.75 (17.97)	13.37	5.20	2.51	34.68	76.43	87.25
Dharani	1.43 (1.56)	32.08 (34.42)	14.45 (22.26)	16.35	4.50	2.92	32.56	63.89	73.55
ICGV 86031 (RC)	0.72 (1.31)	15.04 (22.72)	7.82 (15.94)	22.45	4.60	3.33	48.65	96.01	102.87
ICGV 91114 (SC)	1.42 (1.56)	34.50 (35.95)	17.79 (24.88)	27.90	4.40	2.88	33.26	57.56	63.37
<b>Mean</b>	1.18 (1.48)	25.71 (30.42)	12.93 (20.99)	19.82	5.35	2.49	36.03	60.65	72.47
<b>S.Em±</b>	0.09	0.96	1.47						
<b>CD (P=0.05%)</b>	0.25	2.68	4.10						

RC- Resistant check, SC-Susceptible check; a - Figures in parentheses indicate square root transformed  $\sqrt{(x + 0.5)}$  values; b- Figures in parentheses indicate angular transformed values

**Table 2. Relationship between morphological characters of germplasm lines and tobacco caterpillar incidence and infestation**

S.No	Parameters (X)	Correlation coefficient		
		Tobacco caterpillar larva (Y)	Tobacco caterpillar leaf damage (Y)	Leaf miner leaf damage (Y)
1.	Plant height	0.3801*	0.4511**	0.2968
2.	No. of branches	-0.3150*	-0.4259**	-0.0431
3.	Main stem thickness	-0.6122**	-0.5434**	-0.0767
4.	Trichome density on lamina	-0.5262**	-0.5237**	-0.3325*
5.	Trichome density on midrib	-0.5294**	-0.4732**	-0.2213
6.	Trichome density on petiole	-0.4249*	-0.4005*	-0.1001

\*Significant at 0.05 level; \*\* Significant at 0.01 level

### 3.4 Biochemical Attributes of Resistance/Susceptibility in Selected Groundnut Germplasm Lines

About 13 germplasm lines were selected under different levels of resistance and their biochemical attributes like total sugars, proteins, phenols and tannins were analyzed (Table 3) and correlated with percent damage and population counts of insects to know their role in imparting resistance/susceptibility to germplasm lines.

#### 3.4.1 Tobacco caterpillar incidence and infestation

There was significant positive correlation between incidence and infestation of *S. litura* with total sugars and proteins and significant negative correlation with tannins. With respect to phenols *S. litura* population showed significant negative correlation whereas with leaf damage it

showed non-significant negative correlation (Table 4).

Our findings are in line with Mohammad Saleem et al. [17] who reported significant positive correlation with total sugars against *S. litura* in groundnut. Phenols are plant secondary metabolites that give resistance in plants towards herbivores including insects. Phenols affect the biology of insects in so many different ways. It can act as feeding deterrents, antifeedants and also can cause less damage by larva. Rao [21] showed that higher phenol content in the leaf reduced the incidence of *S. litura* in groundnut suggesting that biochemical components can be effectively utilized in breeding for resistance.

#### 3.4.2 Leaf miner infestation

There was non-significant positive relationship between percent mining and total sugars whereas significant positive relationship was seen with proteins. Phenols were found to have

**Table 3. Biochemical characters of selected germplasm lines**

S.No	Germplasm lines	Total sugars (mg/g)	Proteins (mg/g)	Phenols (mg/g)	Tannins (mg/g)
1	ICGV 15083	2.67	2.57	0.94	0.0026
2	ICGV 181011	4.90	2.39	0.82	0.0030
3	ICGV 13189	5.91	2.58	0.76	0.0028
4	ICGV 93382	2.50	2.89	0.68	0.0023
5	ICGV 10001	3.54	2.92	0.78	0.0032
6	ICGV 10021	4.81	2.98	0.64	0.0022
7	ICGV 02266	4.10	2.36	0.73	0.0040
8	ICGV 00298	3.15	2.35	0.90	0.0047
9	ICGV 93468	2.24	2.14	0.92	0.0050
10	K 6	3.85	2.52	0.75	0.0032
11	JCG 4798	6.82	2.82	0.81	0.0035
12	ICGV 86031 (R)	2.53	2.23	0.97	0.0045
13	ICGV 91114 (S)	4.86	2.94	0.61	0.0024



**Table 4. Relationship between biochemical characters of selected germplasm lines and incidence and infestation of tobacco caterpillar and leaf miner**

Biochemical characters	<i>S. litura</i> (No./plant)	<i>S. litura</i> leaf damage (%)	Leaf miner damage (%)
Total sugars	0.707**	0.672**	0.218
Proteins	0.677**	0.710**	0.594*
Phenols	-0.640*	-0.669**	-0.561*
Tannins	-0.602*	-0.626*	-0.411

\*Significant at 0.05 level; \*\* Significant at 0.01 level

significant negative correlation with percent mining while non-significant negative correlation occurred with tannins (Table 4).

Ravi Chandra Reddy [22] reported that phenols were significantly negatively correlated and total sugars were positively correlated with leaf miner damage in groundnut. Strong negative correlation between percent foliage damage by leaf miner with total phenol content in the leaf was reported in groundnut by Chandramani [23]. Praveen [24] reported significant positive correlation between total sugars and leaf miner damage in groundnut variety KRG 1. Similar results were reported by Ambenagare et al. [25] and Halder et al. [26] in soybean.

#### 4. CONCLUSION

Among the forty germplasm lines screened against tobacco caterpillar and leaf miner, three germplasm lines viz., ICGV 16679 (13.71% leaf damage), ICGV 07222 (14.00% leaf damage) and ICGV 93468 (14.25% leaf damage) showed greater resistance than the resistant check, ICGV 86031 (15.04% leaf damage) against tobacco caterpillar infestation and with respect to resistance against leaf miner, no germplasm line was found to be superior than the resistant check, ICGV 86031 (7.82% leaf damage). However, the germplasm line, ICGV 02266 (8.35% leaf damage) was next best to the resistant check. The characters like trichome density, phenols and tannins showed significant negative correlation with insect damage in the test lines which is of significant important in breeding for resistance.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Dwivedi SL, Crouch JH, Nigam SN, Ferguson ME, Paterson AH. Molecular

- breeding of groundnut for enhanced productivity and food security in the semi-arid tropics: Opportunities and challenges. *Advances in Agronomy*. 2003;80:153-221.
- Sharma HC, Pampapathy G, Dwivedi SL, Reddy LJ. Mechanisms and diversity of resistance to insect pests in wild relatives of groundnut. *Journal of Economic Entomology*. 2003;96(6):1886-1897.
- Dabhade PL, Bapodra JG, Jethva DM, Rathod RT, Dabhi MV. Estimation of yield losses due to major insect pests of groundnut in Gujarat. *Legume Research-An International Journal*. 2012;35(4):354-356.
- Iqbal J, ul Hasan M, Ashfaq M, Sahi ST, Ali A. Studies on Correlation of *Amrasca biguttula biguttula* (Ishida) Population with Physio-morphic Characters of Okra, *Abelmoschus esculentus* (L.) Monech. *Pakistan Journal of Zoology*. 2011;43(1).
- Rama Prasad G. Integrated management of *Spodoptera litura* (F.) in India. In Proc. Nation. Scientists Forum *Spodoptera litura*. 1997; 96-103.
- Painter RH. Insect resistance in crop plants. The Macmillan Publishers Limited; 1951.
- Bhatti MA, Saeed M, Chatta N, Iqbal S. Host plant resistance and importance to insect population suppression in cotton crop. In Proc. Cott. Prod. Seminar, ESSO Pak. Fertilizer Co. Ltd. 1976;132-142.
- Vyavasaya Panchangam. Available: <https://pjtsau.edu.in/files/publications/2018/VPanchangam2018-19.pdf>
- AICRP on groundnut. Proceedings of VII annual rabi-groundnut workshop. Rahuri Mahatma Phule Agriculture University. 1986;4-5.
- Chakravarthy AK, Selvanarayanan S. Experimental techniques in host plant resistance. Springer Nature Singapore Pte Ltd. 2019;215.
- Hedge JE, Hofreiter BT. In: *Methods in Carbohydrate Chemistry* (Eds. Whistler,

- R.L and Be Miller, J.N), Academic press, New York. 1962;17:420.
12. A.O.A.C. Official Methods of Analysis, 16th edition. Association of Official Analytical Chemists, Washington, D.C; 1995.
  13. Mariotti F, Tome D, Mirand PP. Converting nitrogen into protein- Beyond 6.25 and Jones factors. *Critical Reviews in Food Science and Nutrition*. 2008;48:177-184.
  14. Sadasivan S, Manickam A Phenolics. (in) *Biochemical methods, New age international publishers, New Delhi, India*. 2008;3:203-204.
  15. Schanderl SH. *Methods in food analysis*, Academic press, New York. 1970;709.
  16. Krishnaiah M. Screening of promising groundnut genotypes to insect pest complex and their management with certain insecticides. M.Sc (Agri) Thesis ANGRAU; 2011.
  17. Mohammad Saleem A, Gopalakrishna Naidu K, Tippannavar PS, Nadaf HL. Biophysical and biochemical mechanism of resistance to *Spodoptera litura* in groundnut (*Arachis hypogaea* L.). *Journal of Entomology and Zoology studies*. 2019; (4):86-96.
  18. Ranga Rao GV. Studies on seasonal occurrence and host preference of groundnut leaf miner, *Aproaerema modicella* in some selected groundnut lines. Ph. D. Thesis ANGRAU Hyderabad; 1991.
  19. Visalakshi M. Biochemical basis of resistance in groundnut cultures to groundnut leaf miner *Aproaerema modicella* Deventer. *Journal of Research, ANGRAU*. 1997;23(3):15-17.
  20. Peeru Saheb Y, Hari Prasad KV, Swarajya Lakshmi K, Sailaja Rani J. Mechanisms of resistance in groundnut genotypes against leaf bud borer, *Anarsia ephippias* (Meyrick). *Journal of Entomology and Zoology Studies*. 2018;6(2): 694-705.
  21. Rao KR. Influence of host plant nutrition on the incidence of *Spodoptera litura* and *Helicoverpa armigera* on groundnut. *Indian Journal of Entomology*. 2003;65: 386-392.
  22. Ravi Chandra Reddy P. Field screening of groundnut genotypes against leaf miner (*Aproaerema modicella* Deventer) and its management with insecticides. M.Sc. (Agri.) Thesis ANGRAU Guntur; 2019.
  23. Chandramani P, Rajendran R, Sivasubramanian P and Muthiah C. Impact of biophysical factors as influenced by organic sources of nutrients on major pests of rice. *Journal of Biopesticides*. 2009;2:1-5.
  24. Praveen YV. Studies on groundnut leaf miner *Aproaerema modicella* Deventer in Northern dry zone of Karnataka. M.Sc. (Agri.) Thesis Univ Agric Sci Dharwad, Karnataka (India); 2010.
  25. Ambenagare RM, Shetgar SS and Takankhar VG. Biochemical basis of resistance in soybean cultivars to leaf miner *Aproaerema Modicella* Deventer. *Indian Journal of Entomology*. 2011;73(1): 93-94.
  26. Halder J, Srinivasan S and Muralikrishna T. Biochemical basis of resistance to spotted pod borer, *Maruca vitrata* (Geyer) in Mungbean. *Journal of Entomological Research*. 2006;30(4):313-316.

© 2021 Rahman et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/80664>