



Effectiveness of Health Interventions towards HIV/AIDS in Developing Countries: A Systematic Review

**Sarmad Jamal Siddiqui^{1,2}, Rosnah Sutan^{1*}, Zaleeha Md Isa¹,
Arshad Hussain Laghari³ and Vija Kumar Gemnani²**

¹*Department of Community Health, Faculty of Medicine, University Kebangsaan Malaysia, Kuala Lumpur, Malaysia.*

²*Faculty of Community Medicine & Public Health Sciences, Shaheed Mohtarma Benazir Bhutto Medical University, Larkana, Sindh, Pakistan.*

³*Department of Biochemistry, Ghulam Muhammad Mahar Medical College Sukkur @ Shaheed Mohtarma Benazir Bhutto Medical University Larkana, Malaysia.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i47A33044

Editor(s):

(1) Dr. Rafik Karaman, Al-Quds University, Palestine.

Reviewers:

(1) Harrison Daka, University of Zambia, Zambia.

(2) Taiwo Modupe Balogun, Igbinedion University, Nigeria.

(3) Bahizire Riziki Richard, Université de Développement Durable en Afrique Centrale, Congo.

Complete Peer review History: <https://www.sdiarticle4.com/review-history/75801>

Systematic Review

Received 13 August 2021

Accepted 27 October 2021

Published 28 October 2021

ABSTRACT

Background: Plentiful development has been achieved in interventions for the prevention of HIV. Although, progression of prevention programs based on evidence – informed methods that interpret the effectiveness of these approaches in population is still a challenge. In developing countries, not many interventions are implemented for reduction of HIV burden. The single most important identified problem is lack of demand, supply, and adherence approaches. In current systemic review, recent evidence for the prevention of HIV in a cascade manner is described to see status of current interventions and further needs for improvements.

Methodology: Systemic reviews regarding effectiveness on interventions of HIV prevention were searched. Primary studies were identified from eligible review that evaluated one of following

factors: prevalence of HIV, incidence of HIV, testing uptake of HIV and use of condom. Interventions were categorized that pursued demand for prevention of HIV, improvement in supply for preventive approaches, support related to preventive behaviors or prevent HIV directly. A rating was assigned for each intervention based on evidence strength or randomized controlled trials.

Results: Out of 91 eligible reviews, 264 primary studies were included in this review. Primary studies related to direct mechanisms of prevention that showed strong data for circumcision and effectiveness of pre – exposure prophylaxis. Evidence implies that interventions related to increased supply of preventive methods including clean needles or condoms can be operative. Interventions related to demand – side and adherence approaches were less clear with some studies showing effectiveness. Quality evidence was assessed among various categories. Various interventions showed supportive outcomes and results. In our findings, it was observed that difference between behavioral and structural has not evidently distinguished the interventions.

Conclusion: Growing data is present for the support of effectiveness of products, behaviors, and procedures for prevention of HIV. In developing countries, negligible data is present for implementation of such approaches on community level. Interventions will be required for transforming this evidence to produce impact on population. It will empower the demand for prevention of HIV, supply of preventive technologies and utilization of preventive approaches against HIV. The findings can be eye opener to see actual burden of HIV and their implanted interventions and can be useful to design further intervention programs in future.

Keywords: HIV; prevention; reviews; effectiveness; Health education.

1. INTRODUCTION

There has been advancement in delivery and development of interventions regarding prevention against human immunodeficiency virus (HIV), but every year more than one million individuals are emerging with new HIV cases [1]. HIV prevention reinvigoration have been called by United Nations Program for HIV/AIDS (UNAIDS) concerning methods and demonstrated that globally 25% expenditure of HIV should be reserved for activities of prevention [2]. Cascade of HIV prevention have been growing and used for supporting the implementation and development of interventions and for the facilitation of resource distribution. For HIV prevention, specific cascades elaborates that the target and outcome can be achieved by implanting key components including demand – side interventions for awareness and perception acceptability; supply – side interventions production of easy available and accessible products; and adherence interventions for behavior acceptance. Reframing of interventions regarding HIV prevention that are systematized nearby a cascade of HIV prevention has been suggested by Hargreaves et al. [3]. This reformed cascade can assimilate evidence from various disciplines and can be very useful for programmers. Effective treatment and preventive measures are necessary in those at risk. Such behaviors or products are direct mechanisms for the exertion of effectiveness of program for HIV prevention. The concept of cascade is used to

see effectiveness of programs for the prevention of mother to child transmission. Such approaches have demonstrated the needs for suppression of HIV in communities and is being used widely for assessing the programs. As by the treatment cascade, the prevention cascade can also be powerful intervention by provision of population proportion estimation which were lost at certain steps of implementation, explaining the steps in productive interventions for prevention, which represent components to identify the inadequacies and produce further action plan. Characteristics of interventions for treatment of HIV relies on presenting data i.e., from health facilities that track the patients or studies that are population based. There can be a variety in the approach as per its provision i.e., it follows longitudinal cohorts' pattern that describe the evolution over time or population cross – sectional view at time. Such interventions perhaps include prevention of HIV but highlight the diagnosis steps, treatment linkage, initiation of treatment and suppression of viral load. Testing of HIV is important pathway for both prevention and treatment of HIV. To prevent HIV, achievement of all interventions in single approach is difficult due to variability in different approaches they include. Each intervention can be described and separated out in terms of how it can fail or succeed in epidemiological context to elaborate the required steps for specific approach. An observational data was used by Garnet et al. for implementation of an idea for

cascade of HIV prevention to use as monitoring tool [4].

In current review, the available HIV prevention evidence have been reviewed as was reflected in previous reviews of interventions of HIV prevention. The evidence in relation with cascade of HIV prevention is mapped, that explains intervention characteristics in relevance to each cascade area, assessment of available evidence types on these interventions and identification of area and gaps for further research.

2. METHODOLOGY

2.1 Search Methods and Reviews Selection

Three different systemic searches were performed for the identification of HIV prevention and intervention systemic reviews. Terms included HIV/AIDS MeSH terms, behavior, prevention, and intervention. We searched MEDLINE, Cochrane Library, ClinicalTrials.gov. and ISI Web of Knowledge.

Extraction of data was done from review by data extraction tool, which included their reviews, study design, location, population, intervention, and outcome. Reviews with effectiveness of prevention and intervention on HIV were included. Reviews from both observational and experimental studies were included. Literature review, scoping views and broad overviews were excluded. Reviews having structural and behavioral interventions done in developed countries only were excluded as we focused on areas with highest HIV burden and so the interventions effectiveness could be specific to the context and data synthesis would present heterogeneity of studies.

2.2 Identification of Primary Study and Extraction of Data

Primary studies were extracted from reviews if they meet one of following criteria: Prevalence of HIV, incidence of HIV, condom usage reporting and HIV testing uptake. In studies of only direct mechanisms, incidence of HIV was the primary outcome to match inclusion. HIV testing uptake and condom use were included as adjacent intervention effectiveness outcome. Though some reviews identified intervention and prevention of vertical transmission (mother-to-

child), with special consideration to transmission by needle-sharing and intercourse.

Extraction of minimal data at primary study level was developed including target population, country of focus, design of study and reported outcomes for each study. Primary studies and reviews with prevention cascade of HIV as per Hargreaves et al. [3] Many primary studies qualify for multiple categories, but in current study, based on most projecting component we assigned each study into single category (Table 1).

Studies with demand – side domain in which interventions were influencing the behavior by strengthening awareness of, targeting risk perception and positive attitude, prevention of HIV by technologies and behavior were judged. Such interventions were those providing education, information, and communication as well as those proposed to impact apparent patterns via peer-based methods. Interventions were distributed to various target populations and in wide range of settings.

Studies with supply-side domain in which interventions were influencing the messages and products for prevention of HIV, such as treatment planning for sexually transmitted diseases (STDs), initiatives for needle exchange mainstreaming prevention of HIV and mass condom distribution were judged. Some of these interventions are identified as structural interventions in published literature.

Adherence domain included studies in which interventions supported maintenance of behaviors for prevention or adoption were judged. Such interventions influenced behavioral skills or self – efficacy such as longitudinal risk counselling. In this category, the interventions with targeted determinants of social behaviors theorized for acting as barriers to capability of individuals to adhere or access to prevention were also included such as interventions of employment. Some of these interventions are also characterized as structural interventions in published literature.

To determine the efficacy of procedures, the studies conduct randomized control trials as direct mechanism.

Specific kinds of interventions were identified among each domain. As per target populations, each intervention was classified (Table 1). Based

on reported findings and study design, the direction and type of evidence were assessed for each outcome. Framework produced by Mavedzenge et al. was used for assessing adolescents and young individuals [5]. In each category, based on number of randomized controlled trials for certain outcome, the study design was described with ratings A/B/C (Table 2-3). Effectiveness of interventions was assessed by allocating a score from 1-4 (Table 2). The evidence for behavioral and structural interventions were assessed by two reviewers. After consultation, discrepancies were resolved.

3. RESULTS

A total of 91 reviews were eligible for inclusion [6-97] (Fig. 1), out of which 264 primary studies were extracted (Fig. 2). 178 primary studies were related to demand-, supply-side or adherence interventions. Of them, 118 (66.29%) had observational type of study design. Out of 79 randomized controlled trials, 27 (34.17%) used direct prevention mechanisms against HIV. Out of those studies, 18 (10.11%) included prevalence or incidence as primary outcome; through rest demonstrated use of condoms.

Various approaches were used for the influence on risk perception, attitude, and awareness regarding preventive measures such as text

messages, multimedia, and posters etc. For instance, in Zimbabwe, the Helping Each Other Act Responsibly Together (HEART) campaigns used programs through multimedia, advertisements etc. to convey message regarding risk reduction of sexually transmitted infections and HIV [94]. In KwaZulu – Natal, a school-based program was organized for provision of education of HIV prevention and sexual health through performances in drama and various booklets. [95]. More than half of communication, education and information studies were emphasized on young individuals. A randomized controlled trial by Kwa Vijana provided the data of education regarding sexual health in children of school by a participatory program along with health worker trainings to provide sexual health services [96]. In majority of studies, the primary outcome was the evaluation of condom.

Peer – led interventions were used for female sex workers for their support and to produce a sagacity of community. In Philippines, a peer-delivered program regarding education with use of condoms and trainings on information of HIV [97-98]. 10 studies showed peer – based interventions in young individuals. In Kenya, a project involved peer teaching of students by educators regarding HIV information through quizzes, songs, and competitive methods [99-100].




Fig. 1. Systemic reviews identification

Table 1. Categories of prevention intervention against HIV in terms of prevention cascade of HIV

Review & References	Type of Intervention	Subcategory	Outcomes
Demand-side interventions [6-8, 10-12, 14, 17, 19, 27-28, 30, 32, 43, 56-57, 60, 71, 95-96]	Information, education, and communication Peer	Young males, females, individuals on drugs and mass media Young homosexual males, female sex workers, individuals on alcohol or drugs	Awareness Target risk perception Positive attitude
Supply-side interventions [9, 13, 18, 21-22, 24, 26, 31, 33-36, 40, 42, 49-51, 58, 63-64, 79-81, 94-95]	Syringe or needle programs HIV services integration Sexually transmitted infections interventions on community-level Distribution of condoms	---	Messages Products for HIV prevention
Adherence interventions [15-16, 37-39, 41, 44-45, 48, 53-55, 57-58, 60-62, 64-65]	Socioeconomics Counselling	Interventions related to microfinance and transfer of cash Counselling individually, counselling of couples, testing of HIV	Behavior for prevention and adoption (behavioral skills, self-efficacy)
Direct HIV prevention mechanisms [20, 23, 25, 29, 46-47, 52, 54-55, 58-59, 66-70, 72-80, 82-93, 97]	Voluntary circumcision medically Pre-exposure prophylaxis Condoms Treatment of sexually transmitted infections Microbicides Vaccines	Male-to-female and female-to-male transmission, homosexual males	Incidence of HIV HIV testing uptake Condom use

Table 2. Strength assessment of prevention interventions against HIV along intervention types

	No Randomized Controlled Trials (Only observational studies)	1-2 Randomized Controlled Trials	≥3 Randomized Controlled Trials
Consistent effectiveness	C1	B1	A1
Effectiveness on larger scale	C2	B2	A2
Mixed results (beneficial + harmful)	C3	B3	A3
No effectiveness/ harmful effects	C4	B4	A4

	Prevalence		Incidence		Testing of HIV		Use of Condoms	
	Number	QAR	Number	QAR	Number	QAR	Number	QAR
Direct Mechanisms								
Circumcision for heterosexual risk (male-to-female)	-	-	6	B3	-	-	-	-
Circumcision for heterosexual risk (female-to-male)	-	-	33	A1	-	-	-	-
Circumcision for homosexual risk (male-to-male)	-	-	15	C3	-	-	-	-
Individuals-level studies on condoms (heterosexual)	-	-	5	C1	-	-	-	-
Individuals-level studies on PrEP	-	-	5	A2	-	-	-	-
Individual-level studies on microbicide prophylaxis	-	-	9	A3	-	-	-	-
Individual-level studies on STI	-	-	6	A4	-	-	-	-
Individual-level studies on HIV vaccine	-	-	4	A3	-	-	-	-
Interventions based on Demand-Side								
Efficacy of IEC on young individuals	1	B4	2	B4	-	-	18	A3
Efficacy of IEC on males	-	-	-	-	1	C1	7	A2
Efficacy of IEC on females	-	-	-	-	-	-	2	B3
Efficacy of IEC using mass media	-	-	1	B3	-	-	7	B4
Efficacy of IEC on individuals on drugs	-	-	-	-	-	-	3	A1
Efficacy of peer-based interventions on young individuals	-	-	1	B4	2	C1	10	C2
Efficacy of peer-based interventions on male homosexuals	-	-	-	-	1	C1	2	B1
Efficacy of peer-based interventions on sex workers (females)	3	C4	3	C4	3	C1	19	B2
Efficacy of peer-based interventions on individuals on alcohol or drugs	1	B4	2	B4	-	-	3	B3
Interventions based on Supply-Side								
Efficacy of interventions using integration of HIV services routinely	-	-	-	-	3	C1	1	C1

Efficacy of programs concerning clean syringe or needle	4	C1	2	C3	-	-	-	-
Efficacy of interventions based on distribution of condoms	2	C1	-	-	-	-	16	A1
Efficacy of interventions based on STI on community-level	-	-	3	A4	-	-	1	B4
Interventions based on Adherence								
Efficacy of counselling (couples-based)	-	-	1	C1	3	A3	5	A1
Efficacy of counselling (individual-level)	-	-	1	B3	2	B3	10	A1
Efficacy of counselling and testing of HIV	-	-	1	B4	2	B3	11	A1
Efficacy of prevention counselling related to HIV-positive individuals	-	-	-	-	-	-	6	A3
Efficacy of interventions based on microfinance	-	-	1	B4	1	B1	6	A3
Efficacy of interventions based on transfer of cash	2	B1	2	B4	-	-	1	B4

Number indicates the number of studies.
 QAR indicates the Quality Assessment Rating
 IEC = Information, Education and Communication
 PrEP = Pre-exposure Prophylaxis
 STI = Sexually Transmitted Infection

Table 3. PICO Template for questions

Population	Adult males and females
Intervention	Direct mechanisms Demand-side intervention Supply-side intervention Adherence intervention
Comparison	Not applicable
Outcome	Prevalence, incidence, HIV testing, condom use
HIV OR AIDS	
AND	
Behavior OR prevention	
AND	
Intervention	




Fig. 2. Identification of primary studies

Policy changes facilitated the approaches such as promoting the use of condoms to population at risk or accessibility to free and clean needles. In 1989, policy for 100% use of condoms in Thailand was launched to promote safety among sex workers with collaboration between various factors [101]. In Cameroon and Dominican Republic, same approaches have been executed for other groups such as young adults [102-103]. An intensive sexually – transmitted infections control program was carried out in Rakai, Uganda through home – based mass treatment with antibiotic as a randomized controlled trial [104].

Ten studies described interventions on individual level. In South Africa, a program was focused on HIV – free individuals and conveyed a 1-hour counselling session regarding risk reduction from health educators [105]. Six posttest counseling studies described prevention counselling regarding patients positive for HIV. In South

Africa, a randomized controlled trial was carried out, between HIV patients and counsellors during their routine clinical visits focusing on risk reduction [106].

Aim of intervention was to enhance the awareness in schools and reduction in rates of HIV in young individuals [107]. An exigence model of management was used by some interventions including programs regarding cessation of smoking in which behavioral monitoring on regular basis was assessed along with incentives for demonstration of required behavior [108]. In few studies, carried out in Zimbabwe and Kenya, the interventions were carried out along with interventions of life – skills and distribution of condoms [109-110].

4. DISCUSSION

Evident data from various randomized controlled trials were found to support the direct mechanism

efficacy for HIV prevention. From evidence, it is also suggested that interventions based on supply-side that enhance approach to such useful technologies can be beneficial and effective, and there is requirement for continued interventional research for increasing the demand and direct mechanisms adherence for HIV prevention. Interventions and their combinations needed for translation of efficient direct mechanism on the level of population impact will necessitate such domain monitoring for the understanding of breaches and for supporting developmental interventions.

For the attainment of impact on population – level much is to be discovered about augmentation to support the adherence and coverage. A range of possible interventions related to such components have been identified in this review. Interventions based on supply-side including initiative of syringe and needle sharing and distribution of mass condom also have influence of such used methods. Though, on outcome of HIV, the efficacy of such interventions has been explored by only few studies.

Interventions based on demand – side described findings including communication, education and information and peer – based interventions on outcome of HIV are unsatisfactory as they rarely reduce prevalence or incidence of HIV. Few studies and trials showed in reviews assessed interventions for increasing circumcision and pre – exposure prophylaxis demand, though data from these components are still in emerging stage [110-116]. Additional research is needed for understating the reason of decreased uptake for these approaches and assessments needed for improved interventions to increase the adherence and uptake. Data is growing for increase in circumcision demand, so systemic reviews of such approaches are defensible. By emergence of direct mechanisms such as vaccines and microbicides, current interventions can be helpful for improvement in accessibility [116-120].

Effectiveness and evidence of interventions based on supply-side is judicious sign of improvements that can be made in prevention of HIV by accessibility of prevention products to population and community. In situations when there is threat to efforts for reduction in incidence of HIV, such interventions can be used for the prevention of HIV and ultimately help in incidence reduction. Our review showed similar

findings and conclusion as described by Mavadzenge et al. [5].

Our review suggests that difference between behavioral and structural has not evidently distinguished the interventions, so the classification in this manner may produce some misunderstanding. In one review there was identification of Steppingstone which targeted young males and females and for the exploration of intervention effect on outcome of individual biological factors through changes in gender customs to structural level [100]. An intervention for youth through social marketing of condom use was added in reviews and was assessed through structural and behavioral search [103]. Such examples show that categorizing the level of intervention operation might not be as useful as compared to its categorization by intervention objective.

This review also showed some gaps that are present in literature data on intervention effectiveness for the prevention of HIV; especially in terms of demand-, supply-side and adherence interventions. Though, large number of studies were identified across these approaches. Majority of such studies were having observation type of design and were dependent on self – reported outcomes in behaviors. This can be understanding in the way that for effectiveness, these studies are less helpful as compared to studies carried out by randomized controlled trials. Although, when randomized controlled trials are not practicable or unethical, then observational design are necessary. Analysis of approaches of HIV preventions can help in identification of barriers for effective population prevention, shown by application of various prevention formulations from data of high-risk areas. These are used to highlight the inadequacy of service delivery in prevention, low risk perception and poor HIV prevention uptake.

Our methodology has few limitations. As we performed review of reviews, the studies were assessed which themselves comprised of systemic reviews; there were no additional biomedical preventive studies were included. Many studies were identified including primary studies of interventions which were complex that contain factors intended to enhance the demand through peer interventions, communication, education, and information. This review aimed for intervention mapping for prevention of HIV framework by main component of intervention.

Although, there was subjective classification and were classified differently by reviewers or might have chosen the intervention cataloguing into many categories. In this regard, some data might be under – reported categories. In this review, the classification was intended into single category so that exaggeration of the evidence can be avoided. Where possible, we elaborated the evidence for outcome of HIV. Although, outcomes from self – reported behavioral studies were only interventional approach shown in studies; especially studies concerning demand-, supply-side and adherence interventions. Due to inclusion of these studies, it was established that adjacent measures can be influenced by interventions. The objective of this systemic review did not include the assessment of methodological precision of primary studies showed by these reviews. We categorized the primary studies rather than reviews, into domains of definite cascade. This review delivered overview of evidence strength and exactness; though, quality of primary studies is not given in details. Due to these minimal extraction of data at the level of data from primary studies, remarks on population heterogeneity in studies. It is important to understand the study design e.g., observational etc., to demonstrate their usefulness and their similarity with other study designs for efficacy. In subsequent step, consideration of significant connection between studied populations and quality of evidence is necessary.

5. CONCLUSION

Recent evidence shows that we have approaches that slog on individual level and reach the level of effect on population by various interventions to prevent HIV. The cascade of HIV prevention that contains these factors gives an outline for the understanding of the reason for failure of direct mechanisms impact on population level and support the implementation and development of interventions for targeting of these domains. HIV prevention has been categorized according to prevention approach so that it can be helpful in highlighting and recognition of gaps in interventions. Development of understanding can be started for specified problems for better demand, supply, and adherence approaches.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. UNAIDS. The Gap Report. Geneva: UNAIDS; 2016.
2. UNAIDS. End Inequalities. End AIDS. Global AIDS Strategy. Geneva: UNAIDS; 2021.
3. Hargreaves JR, Delany-Moretlwe S, Hallett TB, Johnson S, Kapiga S, Bhattacharjee P et al. The HIV prevention cascade: integrating theories of epidemiological, behavioural, and social science into programme design and monitoring. *Lancet HIV*. 2016;3:e318–22.
4. Garnett GP, Hallett TB, Takaruzza A, Hargreaves J, Rhead R, Warren M et al. Providing a conceptual framework for HIV prevention cascades and assessing feasibility of empirical measurement with data from east Zimbabwe: a case study. *Lancet HIV*. 2016;3:e297–306.
5. Mavedzenge SN, Luecke E, Ross DA. Effective approaches for programming to reduce adolescent vulnerability to HIV infection, HIV risk, and HIV-related morbidity and mortality: a systematic review of systematic reviews. *J Acquir Immune Defic Syndr*. 2014;66:S154–69.
6. Sanii AS, Abraham C, Danford et al. School-based sexual health education interventions to prevent STI/HIV in sub-Saharan Africa: a systemic review and meta-analysis. *BMC Public Health*. 2016;16. Available:<http://doi.org/10.1186/s12889-016-3715-4>.
7. Bertrand JT, O'Reilly K, Denison J, Anhang R, Sweat M. Systematic review of the effectiveness of mass communication programs to change HIV/AIDS-related behaviors in developing countries. *Health Educ Res*. 2006;21:567–97.
8. Fonner VA, Armstrong KS, Kennedy CE, O'Reilly KR, Sweat MD. School based sex education and HIV prevention in low- and middle-income countries: a systematic

- review and meta-analysis. *PLoS One*. 2014;9:e89692.
9. Levy E, Kaufman MR, Gidron Y, Deschepper R, Olley BO. Interventions targeting social cognitive determinants of condom use in the general Sub-Saharan population: A Systemic Review. 2019;6:1.
 10. Muthoni CN, Kneipp SM, Gichane MW. A Systemic Review of HIV Interventions for Young Women in Sub-Saharan Africa. *AIDS Behav*. 2020;24:3395-3413.
 11. Lameiras-Fernandez MV, Rodriguez-Castro Y. Sex Education in the Spotlight: What is Working? Systemic Review. *Int J Environ Res Public Health*. 2021;18:2555.
 12. LaCroix JM, Snyder LB, Huedo-Medina TB, Johnson BT. Effectiveness of mass media interventions for HIV prevention, 1986–2013: a meta-analysis. *J Acquir Immune Defic Syndr*. 2014;66(suppl 3):S329–40.
 13. Mwale M, Muula AS. Systemic review: a review of adolescent behavior change interventions [BCI] and their effectiveness in HIV and AIDS prevention in sub-Saharan Africa. *BMC Public Health*. 2017;17. Available:<http://doi.org/10.1186/s12889-017-4729-2>.
 14. He J, Wang Y, Du Z et al. Peer education for HIV prevention among high-risk groups: a systemic review and meta-analysis, *BMC Infect Dis*. 2020;20:338.
 15. Iwelunmor J, Nwaozuru U, Obeizu-Umeh C, Uzoaru F, Ehiri J, Curley J et al. Is it time to RE-AIM? A systemic review of economic empowerment as HIV prevention intervention for adolescent girls and young women in sub-Saharan Africa using RE-AIM framework. *Implement Sci Commun*. 2020;1:53.
 16. Sileo KM, Miller AP, Huynh TA, Kiene SM. A systemic review of interventions for reducing heavy episodic drinking in sub-Saharan African settings. *PLOS ONE*. 2020;15(12):e0242678.
 17. Mason-Jones AJ, Sinclair D, Mathews C, Kagee A, Hillman A, Lombard C. School-based interventions for preventing HIV, sexually transmitted infections and pregnancy in adolescents. *Cochrane Database of Systemic Review*. 2016;11:CD006417.
 18. Malekinejad M, Parriott A, Blodgett JC, Horvath H, Shrestha RK, Hutchison AB et al. Effectiveness of community-based condom distribution interventions to prevent HIV in the United States: A systemic review and meta-analysis. *PLOS ONE*. 2017;12(8):e0180718.
 19. Questa K, Das M, King R et al. Community engagement interventions for communicable disease control in low- and lower-middle-income countries: evidence from a review of systemic reviews. *Int J Equity Health*. 2020;19:51.
 20. Yuan T, Fitzpatrick T, Ko N et al. Circumcision to prevent HIV and other sexually transmitted infections in men who have sex with men: a systemic review and meta-analysis of global data. *Lancet Glob Health*. 2019;7:e436-47.
 21. Faini D, Hanson C, Baisley K, Kapiga S, Hayes R. Sexual behavior, changes in sexual behavior and associated factors among women at high risk of HIV participating in feasibility studies for prevention trials in Tanzania. *PLOS ONE*. 2020;15(4):e231766.
 22. Wulandari LPL, Guy R, Kaldor J. The burden of HIV infection among men who purchase sex in low- and middle-income countries – a systemic review and meta-analysis. *PLOS ONE*. 2020;15(9):e238639.
 23. Shen Q, Wang Y, Wang et al. Understanding condom use among unmarried female migrants: a psychological and personality factors modified information – motivation – behavioral skills model. *BMC Public Health*. 2019;223. Available: <http://doi.org/10.1186/s12889-019-6541-7>.
 24. Henderson JT, Senger CA, Henninger M, Bean SI, Redmond N, O'Connor EA. Behavioral Counseling Interventions to Prevent Sexually Transmitted Infections: Updated Evidence Report and Systematic Review for the US Preventive Services Task Force. *JAMA*. 2020;324(7): 682–699.
 25. Jumbe S, Kamninga TM, Mwalwimba et al. Determinants of adolescent substance use in Africa: a systemic review and meta-analysis protocol. *Syst Rev*. 2021;10:125.
 26. Hansell AH, Giacobbi PR, Voelker DK. A Scoping Review of Sport-Based Health Promotion Interventions With Youth in Africa. *Health Promotion Practice*. 2021; 22(1):31-40.
 27. Siddiqui M, Kataria I, Watson K, Chandra-Mouli V. A systemic review of the evidence on peer education programmes for promoting the sexual and reproductive health of young people in India. *Sexual*

- and Reproductive Health Matters. 2020; 28:1.
28. Morar NS, Naidoo S, Goolam A, Ramjee G. Research participants' skills development as HIV prevention peer educators in their communities. *Journal of Health Psychology*. 2018;23(10):1343-1349.
 29. Gannon-Loew KE, Holland-Hall C. A review of current guidelines and research on the management of sexually transmitted infections in adolescents and young adults. *Therapeutic Advances in Infectious Disease*; 2020. DOI:10.1177/2049936120960664.
 30. Berg RC, Page S, Ogard-Repal A. The effectiveness of peer-support for people living with HIV: A systemic review and meta-analysis. *PLOS ONE*. 2021;16(6): e252623.
 31. Herbst JH, Sherba RT, Crepaz N, et al. For the HIV/AIDS Prevention Research Synthesis Team. A meta-analytic review of HIV behavioral interventions for reducing sexual risk behavior of men who have sex with men. *J Acquir Immune Defic Syndr*. 2005;39:228–41.
 32. Broccateli C, Wang P, McDaid L et al. Social Network Research contribution to evaluating process in feasibility study of peer-led and school-based sexual health. *Sci Rep*. 2021;11:12244.
 33. Xin M, Viswanath K, Li AY et al. The Effectiveness of Electronic Health Interventions for Promoting HIV-Preventive Behaviors Among Men Who Have Sex With Men: Meta-Analysis Based on an Integrative Framework of Design and Implementation Features. *J Med Internet Res*. 2020;22(5):e15977.
 34. Cheng W, Xu H, Tang W et al. Online HIV prevention intervention on condomless sex among men who have sex with men: A web-based randomized controlled trial. *BMC Infect Dis*. 2019;19:644.
 35. Herbst JH, Beeker C, Mathew A, et al, for the Task Force on Community Preventive Services. The effectiveness of individual-, group-, and community-level HIV behavioral risk-reduction interventions for adult men who have sex with men: a systematic review. *Am J Prev Med*. 2007; 32(suppl):S38–67.
 36. Herbst JH, Sherba RT, Crepaz N, et al, for the HIV/AIDS Prevention Research Synthesis Team. A meta-analytic review of HIV behavioral interventions for reducing sexual risk behavior of men who have sex with men. *J Acquir Immune Defic Syndr*. 2005;39:228–41.
 37. De Vasconcelos S, Toskin I, Cooper B et al. Behaviour change techniques in brief interventions to prevent HIV, STI and unintended pregnancies: A systematic review. *PLOS ONE*. 2018;13(9):e0204088.
 38. Sagherian MJ, Huedo-Medina TB, Pellowski JA et al. Single-Session Behavioral Interventions for Sexual Health Risk Reduction: A Meta-Analysis. *Ann Behav Med*. 2016;50:920-934.
 39. Chemaitelly H, Weiss HA, Calvert C et al. HIV epidemiology among female sex workers and their clients in the Middle East and North Africa: systemic review, meta-analysis, and meta-regression. *BMC Med*. 2019;17:119.
 40. Atuhaire L, Adetokunboh O, Shumba C et al. Effect of community-based interventions targeting female sex workers along the HIV care cascade in sub-Saharan Africa: a systemic review and meta-analysis. *Syst Rev*. 2021;10:137.
 41. McGranahan M, Nakyeyune J, Baguma C et al. Right based approaches to sexual and reproductive health in low and middle-income countries: A Systemic Review. *PLOS ONE*. 2021;16(4):e0250976.
 42. Atuhaire L, Adetokunboh O, Shumba C, et al. Effect of female sex work-targeted communitybased interventions along the HIV treatment cascade in sub-Saharan Africa: a systematic review protocol. *BMJ Open*. 2020;10:e039495.
 43. Tan JY, Huedo-Medina TB, Warren MR, Carey MP, Johnson BT. A meta-analysis of the efficacy of HIV/AIDS prevention interventions in Asia, 1995–2009. *Soc Sci Med*. 2012;75:676–687.
 44. Rinaldi G, Kiadaliri AA, Haghparast-Bidgoli H. Cost effectiveness of HIV and sexual reproductive health interventions targeting sex workers: a systemic review. *Cost Eff Resour Alloc*. 2018;63. Available:<http://doi.org/10.1186/s12962-018-0165-0>.
 45. Gilchrist G, Swan D, Widyaratna K et al. A Systemic Review and Meta-analysis of Psychosocial Interventions to Reduce Drug and Sexual Blood Borne Virus Risk Behaviors Among People Who Inject Drugs. *AIDS Behav*; 2017. Available:<http://doi.org/10.1007/s10461-017-1755-0>.

46. Johnson BT, Carey MP, Marsh KL, Levin KD, Scott-Sheldon LA. Interventions to reduce sexual risk for the human immunodeficiency virus in adolescents, 1985–2000: A Research Synthesis. *Arch Pediatr Adolesc. Med.* 2003;157:381–88.
47. Johnson BT, Scott-Sheldon LA, Huedo-Medina TB, Carey MP. Interventions to reduce sexual risk for human immunodeficiency virus in adolescents: a meta-analysis of trials, 1985–2008. *Arch Pediatr Adolesc Med.* 2011;165:77–84.
48. Narasimhan M, Yeh PT, Haberlen S et al. Integration of HIV testing services into family planning services: a systemic review. *Reprod Health.* 2019;16:61.
49. Fernandes RM, Cary M, Duarte G et al. Effectiveness of needle and syringe Programmes in people who inject drugs – An overview of systemic reviews. *BMC Public Health.* 2017;17:309.
50. Resiak D, Mpofu E, Rothwell R. Sustainable Harm Reduction Needle and Syringe Programs for People Who Inject Drugs: A Scoping Review of Their Implementation Qualities. *Sustainability.* 2021;13:2834.
51. Hong Y, Poon AN, Zhang C. HIV/STI prevention interventions targeting FSWs in China: a systematic literature review. *AIDS Care.* 2011;23 (suppl 1):54–65.
52. Adamson B, Garrison L, Barnabas RV, Carlson JJ, Kublin J, Dimitrov D. Competing biomedical HIV prevention strategies: potential cost-effectiveness of HIV vaccines and PrEP in Seattle, WA. *Journal of the International AIDS Society.* 2019;22:e25373.
53. Arden-Close E, McGrath N. Health behavior change interventions for couples: A systemic review. *British Journal of Health Psychology.* 2017;22(2):215-237.
54. Fonner Virginia A, Denison J, Kennedy Caitlin E, O'Reilly K, Sweat M. Voluntary counseling and testing (VCT) for changing HIV-related risk behavior in developing countries. *Cochrane Database Syst Rev.* 2012;9:CD001224.
55. El-Bassel N, Gilbert L, Goddard-Eckrich D, et al. Effectiveness of a Couple-Based HIV and Sexually Transmitted Infection Prevention Intervention for Men in Community Supervision Programs and Their Female Sexual Partners: A Randomized Clinical Trial. *JAMA Netw Open.* 2019;2(3):e191139.
56. Kennedy CE, Medley AM, Sweat MD, O'Reilly KR. Behavioural interventions for HIV positive prevention in developing countries: a systematic review and meta-analysis. *Bull World Health Organ.* 2010; 88:615–23.
57. Zhang H, Yu Q, Li Z et al. Efficacy of Psychological Interventions Towards the Reduction of High-Risk Sexual Behaviors Among People Living with HIV: A Systemic Review and Meta-Analysis, 2010-2020. *AIDS Behav;* 2021. Available: <http://doi.org/10.1007/s10461-021-03181-4>.
58. Jason G, Mitra S, David G et al. HIV/STI prevention interventions: A systemic review and meta-analysis. *Open Medicine.* 2017;12(1):450-467.
59. Gonçalves TR, Faria ER, Carvalho FT, Piccinini CA, Shoveller JA. Behavioral interventions to promote condom use among women living with HIV: a systematic review update. *Cad Saude Publica.* 2017;33(1):e00202515.
60. Nkenfou CN, Gouekem J, Socpa A et al. Differential Risky Behaviors among HIV Positive and HIV Negative People. *Int J Virol AIDS;* 2017. Available:<http://doi.org/10.23937/2469-567X/1510037>.
61. Tsai LC, Witte SS, Aira T, Riedel M, Offringa R, Chang M. Efficacy of a microsavings intervention in increasing income and reducing economic dependence upon sex work among women in Mongolia. *International Social Work.* 2018;61(1):6-22.
62. Nadkarni S, Genberg B, Galárraga O. Microfinance Interventions and HIV Treatment Outcomes: A Synthesizing Conceptual Framework and Systematic Review. *AIDS Behav.* 2019;23(9):2238-2252.
63. Gibbs A, Willan S, Misselhorn A, Mangoma J. Combined structural interventions for gender equality and livelihood security: a critical review of the evidence from southern and eastern Africa and the implications for young people. *J Int AIDS Soc.* 2012;15 (suppl 1): 1–10.
64. Kennedy CE, Fonner VA, O'Reilly KR, Sweat MD. A systematic review of income generation interventions, including microfinance and vocational skills training, for HIV prevention. *AIDS Care.* 2014;26: 659–73.

65. MacPhail C, Khoza N, Selin A et al. Cash transfers for HIV prevention: what do young women spend it on? Mixed methods findings from HPTN 068. *BMC Public Health*. 2018;18:10.
66. Olapade-Olaopa EO, Salami MA, Lawal TA. Male circumcision and global HIV/AIDS epidemic challenges. *Afr J Urol*. 2019;25:3
67. Farley TMM, Samuelson J, Grabowski MK, Ameyan W, Gray RH, Baggaley R. Impact of male circumcision on risk of HIV infection in men in a changing epidemic context – systemic review and meta-analysis. *Journal of the International AIDS Society*. 2020;23:e25490.
68. Sharma SC, Raison N, Khan S, Shabbir M, Dasgupta P, Ahmed K. Male circumcision for the prevention of human immunodeficiency virus (HIV) acquisition: a meta-analysis. *BJU Int*. 2018;121:515-526.
69. Darby R. Targeting Patients Who Cannot Object? Re-Examining the Case for Non-Therapeutic Infant Circumcision. *SAGE Open*;2016. DOI:10.1177/2158244016649219.
70. Gao Y, Yuan T, Zhan et al. Association between medical male circumcision and HIV risk compensation among heterosexual men: a systemic review and meta-analysis. *Lancet Glob Health*. 2021;9:e932-41.
71. US Preventive Services Task Force. Behavioral Counseling Interventions to Prevent Sexually Transmitted Infections: US Preventive Services Task Force Recommendation Statement. *JAMA*. 2020;324(7):674–681.
72. Lei JH, Liu LR, Wei Q, et al. Circumcision status and risk of HIV acquisition during heterosexual intercourse for both males and females: a meta-analysis. *PLOS ONE*. 2015;10:e0125436.
73. Gazimbi MM, Magadi MA, Krugar C. The Association between Male Circumcision and HIV Infection in Sub-Saharan Africa: A Systemic Review of the Literature. *Int Arch Pubic Health Community Med*. 2019; 3:022.
74. Awad SF, Sgaier SK, Lau FK et al. Could Circumcision of HIV-Positive Males Benefit Voluntary Medical Male Circumcision Programs in Africa? Mathematical Modeling Analysis. *PLOS ONE*. 2017;12 (1): e0170641.
75. Ahmed P, Farid GM, Afroz T. Male Circumcision and Risk of HIV Transmission from Male to Women. *Medicine Today*. 2016;28(1):22-26.
76. Zhang C, Qian H-Z, Liu Y, Vermund SH. Voluntary medical male circumcision and HIV infection among men who have sex with men: Implications from a systematic review. *SAGE Open Medicine*; 2019. DOI:10.1177/2050312119869110.
77. Millett GA, Flores SA, Marks G, Reed JB, Herbst JH. Circumcision status and risk of HIV and sexually transmitted infections among men who have sex with men: a meta-analysis. *JAMA*. 2008;300: 1674–84.
78. Fankem SL, Wiysonge CS, Hankins CA. Male circumcision and the risk of HIV infection in men who have sex with men. *Int J Epidemiol*. 2008;37:353–55.
79. Giannou FK, Tsiara CG, Nikolopoulos GK, Talias M, Benetou V, Kantzanou M et al. Condom effectiveness in reducing heterosexual HIV transmission: a systematic review and meta-analysis of studies on HIV serodiscordant couples. *Expert Rev Pharmacoecon Outcomes Res*. 2016;16(4):489-99.
80. Johnson WD, O’Leary Ann, Flores SA. Per-partner condom effectiveness against HIV for men who have sex with men. *AIDS*. 2018;32:1499-1505.
81. Hahn A, Hinz R, Meyer T et al. HIV prevention strategies and risk of infection: A model-based analysis. *Epidemiology and Infection*. 2018;146(8):1015-1025.
82. US Preventive Services Task Force. Preexposure Prophylaxis for the Prevention of HIV Infection. Recommendation Statement. *American Family Physician*. 2019;100(10):636F-636J.
83. Traeger MW, Schroeder SE, Wright EJ et al. Pre-exposure Prophylaxis for the Prevention of Human Immunodeficiency Virus Infection on Sexual Risk Behavior in Men Who Have Sex With Men: A Systemic Review and Meta-analysis. *Clinical Infectious Diseases*. 2018;67(5):676-686.
84. Sidebottom D, Ekstrom AM, Stromdahl S. A systemic review of adherence to oral pre-exposure prophylaxis of HIV – how can we improve uptake and adherence? *BMC Infect Dis*. 2018;18:581.
85. Corneli A, Meagher K, Henderson G et al. How Biomedical HIV Prevention Trials Incorporate Behavioral and Social Science Research: A Typology of Approaches. *AIDS Behav*. 2019;23:2146-2154.

86. Obiero J, Ogongo P, Mwethera PG, Wiysonge CS. Topical microbicides for preventing sexually transmitted infections. *Cochrane Database Syst Rev.* 2021;13; 3(3):CD007961.
87. Musekiwa A, Fernando NB, Abariga SA. Effectiveness of vaginal microbicides in preventing HIV transmission. *Tropical Medicine and International Health.* 2020;25 (7):790-802.
88. Ghimire S, Hallett J, Gray C, Lobo R, Crawford G. What Works? Prevention and Control of Sexually Transmitted Infections and Blood-Borne Viruses in Migrants from Sub-Saharan Africa, Northeast Asia and Southeast Asia Living in High-Income Countries: A Systematic Review. *International Journal of Environmental Research and Public Health.* 2019;16 (7):1287.
89. Wilkinson D, Ramjee G, Tholandi M, Rutherford G. Nonoxynol-9 for preventing vaginal acquisition of sexually transmitted infections by women from men. *Cochrane Database Syst Rev.* 2002;4:CD003939.
90. Wilkinson D, Ramjee G, Tholandi M, Rutherford G. Nonoxynol-9 for preventing vaginal acquisition of HIV infection by women from men. *Cochrane Database Syst Rev.* 2002;4:CD003936.
91. Balkus JE, Manhart LE, Lee J et al. Periodic Presumptive Treatment for Vaginal Infections May Reduce the Incidence of Sexually Transmitted Bacterial Infections. *The Journal of Infectious Diseases.* 2016;213(12):1932-1937.
92. Kularatne RS, Muller EE, Maseko DV, Kufa-Ch akezha T, Lewis DA. Trends in the relative prevalence of genital ulcer disease pathogens and association of HIV infection in Johannesburg, South Africa, 2007-2015. *PLOS ONE.* 2018;13(4): e0194125.
93. Cook RL, Rosenberg MJ. Do spermicides containing nonoxynol-9 prevent sexually transmitted infections? A meta-analysis. *Sex Transm Dis.* 1998;25:144-50.
94. Underwood C, Hachonda H, Serlemitos E, Bharath-Kumar U. Reducing the risk of HIV transmission among adolescents in Zambia: psychosocial and behavioral correlates of viewing a risk-reduction media campaign. *J Adolesc Health.* 2006; 38: 55.
95. Harvey B, Stuart J, Swan T. Evaluation of a drama-in-education programme to increase AIDS awareness in South African high schools: a randomized community intervention trial. *Int J STD AID.S.* 2000;11: 105-11.
96. Ross DACJ, Changalucha J, Obasi AI, et al. Biological and behavioural impact of an adolescent sexual health intervention in Tanzania: a community-randomized trial. *AIDS.* 2007;21:1943-55.
97. Chiao CMD, Ksobiech K, Malow R. Promoting HIV testing and condom use among Filipina commercial sex workers: findings from a quasi-experimental intervention study. *AIDS Behav.* 2009;13: 892-901.
98. Morisky DE, Chiao C, Stein JA, Malow R. Impact of social and structural influence interventions on condom use and sexually transmitted infections among establishment-based female bar workers in the Philippines. *J Psychol Hum Sex.* 2005; 17:45-63.
99. van der Maas F, Otte WM. Evaluation of HIV/AIDS secondary school peer education in rural Nigeria. *Health Educ Res.* 2009;24:547-57.
100. Jewkes R, Nduna M, Levin J, et al. Impact of Stepping Stones on incidence of HIV and HSV-2 and sexual behaviour in rural South Africa: cluster randomised controlled trial. *BMJ.* 2008;337:a506.
101. Jose JEd, Sakboonyarat B, Kana K, et al. Prevalence of HIV infection and related risk factors among young Thai men between 2010 and 2011. *PLOS ONE.* 2020;15(8):e0237649.
102. Carrasco M, Barrington C, Perez M, Donastorg Y, Kerrigan D. Social cohesion, condom use, and sexually transmitted infections among female sex workers living with HIV in the Dominican Republic. *International Journal of STD & AIDS.* 2019;30(1):64-71.
103. Meekers D, Agha S, Klein M. The impact on condom use of the "100% Jeune" social marketing program in Cameroon. *J Adolesc Health.* 2005;36:530.
104. Mafigiri R, Matovu JKB, Makumbi FE et al. HIV prevalence and uptake of HIV/AIDS services among youths (14-25 Years) in fishing and neighboring communities of Kasensero, Rakai District, South Western Uganda. *BMC Public Health.* 2017; 17:251.
105. Kalichman SC, Simbayi LC, Vermaak R, Cain D, Jooste S, Peltzer K. HIV/AIDS risk reduction counseling for alcohol using

- sexually transmitted infections clinic patients in Cape Town, South Africa. *J Acquir Immune Defic Syndr.* 2007;44:594–600.
106. Hlongwa M, Peltzer K, Hlongwana K. Risky sexual behaviors among women of reproductive age in a high HIV burdened township in KwaZulu-Natal, South Africa. *BMC Infect Dis.* 2020;20:563.
107. Smith LM, Hein NA, Bagenda D. Cash transfers and HIV/HSV-2 prevalence: A replication of a cluster randomized trial in Malawi. *PLOS ONE.* 2019;14(1): e0210405.
108. Ladapo JA, Tseng CH, Sherman SE. Financial Incentives for Smoking Cessation in Hospitalized Patients: A Randomized Clinical Trials. *The American Journal of Medicine.* 2020;133(6):741-749.
109. Dunbar MS, Maternowska MC, Kang MS, Laver SM, Mudekunye-Mahaka I, Padian NS. Findings from SHAZ!: a feasibility study of a microcredit and life-skills HIV prevention intervention to reduce risk among adolescent female orphans in Zimbabwe. *J Prev Interv Community.* 2010;38:147–61.
110. Erulkar AS, Onoka CJ, Phiri A. What is youth-friendly? Adolescents' preferences for reproductive health services in Kenya and Zimbabwe. *Afr J Reprod Health.* 2005; 9:51–58.
111. Diabate S, Kra O, Biekoua J, et al. Pre-exposure prophylaxis among men who have sex with men in Cote d'Ivoire: a quantitative study of acceptability. *AIDS Care.* 2020.
Available:<http://doi.org/10.1080/09540121.2020.1785997>.
112. Rerks-Ngarm S, Pitisuttithum P, Nitayaphan S, et al, for the MOPH-TAVEG Investigators. Vaccination with ALVAC and AIDSVAX to prevent HIV-1 infection in Thailand. *N Engl J Med.* 2009;361:2209–20.
113. Laher F, Bekker LG, Garrett N et al. Review of preventive HIV vaccine trials in South Africa. *Arch Virol.* 2020;165:2439-2452.
114. Ng'uni T, Chasara C, Ndhlovu ZM. Major Scientific Hurdles in HIV Vaccine Development: Historical Perspective and Future Directions. *Front Immunol.* 2020; 11:590780.
115. Musunguzi N, Kidoguchi L, Mugo NR et al. Adherence to recommendations for ART and targeted PrEP use among HIV serodiscordant couples in East Africa: the "PrEP as a bridge to ART" strategy. *BMC Public Health.* 2020;20:1621.
116. Weiss SM, Zulu R, Jones DL, Redding CA, Cook R, Chitalu N. The Spear and Shield intervention to increase the availability and acceptability of voluntary medical male circumcision in Zambia: a cluster randomised controlled trial. *Lancet HIV.* 2015; 2:e181–89.
117. Rees H, Delany-Moretlwe SA, Lombard C, Baron D, Panchia R, Myer L et al, eds. FACTS 001 phase III trial of pre-coital tenofovir 1% gel for HIV prevention in women. *Conferences on Retroviruses and Opportunistic Infections (CROI); Seattle, WA, USA; 2015.*
118. McCormack S, Dunn DT, Desai M, et al. Pre-exposure prophylaxis to prevent the acquisition of HIV-1 infection (PROUD): effectiveness results from the pilot phase of a pragmatic open-label randomised trial. *Lancet.* 2016;387:53–60.
119. Molina JM, Charreau I, Spire B et al. Efficacy, safety, and effect on sexual behavior of on-demand pre-exposure prophylaxis for HIV in men who have sex with men: an observational cohort study. *The Lancet HIV.* 2017;4(9): e402-e410.
120. Effects of cash transfer for the prevention of HIV in young South African women. 8th International AIDS Society (IAS) Conference on HIV Pathogenesis, Treatment and Prevention; Vancouver, BC, Canada; 2015.

© 2021 Siddiqui et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle4.com/review-history/75801>