



# Effect of Liquid Jeevamrutha on Growth Yield and Quality of China Aster (*Callistephus chinensis* [L.] Nees.)

Bhagyashree Narayana Hegde <sup>a++\*</sup>, Anand B. Mastiholi <sup>b#</sup>,  
B. C. Patil <sup>a†</sup>, N. Thammaiah <sup>c</sup>, T. Shantappa <sup>d†</sup>,  
Rudragouda Patil <sup>a‡</sup> and Suvarna Patil <sup>e^</sup>

<sup>a</sup> Department of Floriculture and Landscape Architecture, K. R. C. College of Horticulture, Arabhavi, India.

<sup>b</sup> ZBNF Project, RHREC, Dharwad, India.

<sup>c</sup> University Librarian, Sarvajna Library, UHS, Bagalkot, India.

<sup>d</sup> ADRE, RHREC, Dharwad, India.

<sup>e</sup> AICRP Vegetable Crops, RHREC, Dharwad, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2023/v35i32779

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/97136>

Original Research Article

Received: 22/12/2022

Accepted: 27/02/2023

Published: 01/03/2023

<sup>++</sup> Ph. D Scholar;

<sup>#</sup> Professor of Agronomy and Principal Investigator;

<sup>†</sup> Professor and Head;

<sup>‡</sup> Assistant Professor;

<sup>^</sup> Assistant Professor and Head;

\*Corresponding author: E-mail: shreehegde888@gmail.com;

## ABSTRACT

The experiment on “Effect of liquid jeevamrutha on growth yield and quality of China aster (*Callistephus chinensis* [L.] Nees.)” was carried at Department of Floriculture and Landscape Architecture, K. R. C. College of Horticulture, Arabhavi. Among all the treatments, application of liquid jeevamrutha at 1500 liter per hectare at an interval of 7 days ( $D_3F_1$ ) was recorded significantly higher plant height (47.98 cm and 58.78 cm at 60 and 90 DAT, respectively), plant spread (N-S) *i.e.*, 17.90 26.35 and 31.31 cm at 60 and 90 DAT, respectively, plant spread (E-W) *i.e.*, 27.10 and 30.35 cm at 60 and 90 DAT, respectively, number of primary branches (7.45 and 10.28 at 60 and 90 DAT, respectively), number of flowers per plant (38.42), flower yield (111.39 g/plant, 5.41 kg/plot and 8.06 t/ha, respectively), flower diameter (5.76 cm), individual flower weight (3.04 g), shelf life (3.43 days) and vase life (8.79 days) in pooled data.

**Keywords:** China aster; liquid jeevamrutha; dosage; frequency; growth; flower yield and quality.

## 1. INTRODUCTION

China aster is one of the most popular annuals in addition to chrysanthemum and marigold. The aster is cultivated commercially in India, France, Netherlands, United Kingdom, Siberia, Russia, North America, Japan, Europe and Switzerland. In India, China aster is grown extensively in Karnataka, Tamil Nadu, Telangana, Andhra Pradesh, Maharashtra and West Bengal. In Karnataka, it is cultivated in 316 hectares, producing around 2317 MT and productivity of 7.34 tonnes per hectare (Anon, 2022) [1]. China aster is cultivated throughout the year around Bengaluru area because of its suitability and market. It is adapted to a variety of agro-climatic situations and blooms round the year and is grown all over the world. It is effectively grown in open situations for the year-round production during *kharif*, *rabi* and summer, ensuring a steady supply of flowers for the market. Commercial cultivation of China aster demands excessive usage of synthetic inputs leading to soil and environmental degradation and also agro-ecological imbalances.

At the same time, the cost of synthetic or inorganic fertilizers is increasing enormously. These inputs are also becoming non-affordable to farmers. The excessive and indiscriminate use of chemical fertilizers and pesticides affected the beneficial soil micro flora and fauna and polluted soil and groundwater. Due to mining of nutrients by high value and commercial crops farmers are forced to add fertilizers and inputs externally. If this trend continues, we will be left with little or no chemical fertilizer reserves in future. Hence, we have to find an alternate ways to suffice the nutrient needs of crops. This is compelling the farming community to find out an alternate method which is eco-friendly, low cost, locally

adoptable and enhance or maintain productivity sustainably. Growing flower crops in natural ways by adopting natural farming principles is one of the recent approaches being adopted by some farmers. The use of beejamrutha, ghanajeevamrutha and liquid jeevamrutha, besides organic mulch application in crop production, are the core practices in natural farming which are to be standardized for sustainable flower production. In this regard present study was carried out with the objective to know the effect of liquid jeevamrutha on growth yield and quality of China aster (*Callistephus chinensis* [L.] Nees.).

## 2. MATERIALS AND METHODS

The present investigation was carried out at Kittur Rani Channamma College of Horticulture Arabhavi, Karnataka, under the University of Horticultural Sciences, Bagalkot during *Rabi* 2020-21 and *Kharif* 2021-22. The experiment was laid out in two Factorial Randomized Block Design with fourteen treatments ( $3 \times 4 + 2$ ) and three replications. Factor-A includes  $D_1$ : 500 litre/ha,  $D_2$ : 1000 litre/ha and  $D_3$ :1500 litre/ha, Factor-B includes  $F_1$ : Once in week (7 days),  $F_2$ : Once in two weeks (14 days),  $F_3$ : Once in three weeks (21 days) and  $F_4$ : Once in four weeks (30 days) and these treatment combinations were compared with control treatment *i.e.*, INM: Integrated nutrient management (50% RDN through FYM (9 t/ha) + 50% RDN through inorganic fertilizer (45 kg N/ha) + 120 kg P/ha + K @ 60 Kg/ha + 20 t FYM/ha + *Azospirillum* @ 2.5 kg/ha + PSB @ 2.5 kg/ha and RPP: Recommended package of practice (NPK @ 90:120:60 kg/ha + 20 t FYM/ha). The treatment details were given in Table 1. Vegetative parameter like plant height, plant spread (N-S and E-W) and number of primary branches were taken at 30, 60 and 90 days after planting.

**Table 1. Treatment combinations (D x F)**

T <sub>1</sub>	D <sub>1</sub> F <sub>1</sub>	Application of liquid jeevamrutha @ 500 l/ha once in a week
T <sub>2</sub>	D <sub>1</sub> F <sub>2</sub>	Application of liquid jeevamrutha @ 500 l/ha once in two week
T <sub>3</sub>	D <sub>1</sub> F <sub>3</sub>	Application of liquid jeevamrutha @ 500 l/ha once in three week
T <sub>4</sub>	D <sub>1</sub> F <sub>4</sub>	Application of liquid jeevamrutha @ 500 l/ha once in four week (one month)
T <sub>5</sub>	D <sub>2</sub> F <sub>1</sub>	Application of liquid jeevamrutha @ 1000 l/ha once in a week
T <sub>6</sub>	D <sub>2</sub> F <sub>2</sub>	Application of liquid jeevamrutha @ 1000 l/ha once in two week
T <sub>7</sub>	D <sub>2</sub> F <sub>3</sub>	Application of liquid jeevamrutha @ 1000 l/ha once in three week
T <sub>8</sub>	D <sub>2</sub> F <sub>4</sub>	Application of liquid jeevamrutha @ 1000 l/ha once in four week (one month)
T <sub>9</sub>	D <sub>3</sub> F <sub>1</sub>	Application of liquid jeevamrutha @ 1500 l/ha once in a week
T <sub>10</sub>	D <sub>3</sub> F <sub>2</sub>	Application of liquid jeevamrutha @ 1500 l/ha once in two week
T <sub>11</sub>	D <sub>3</sub> F <sub>3</sub>	Application of liquid jeevamrutha @ 1500 l/ha once in three week
T <sub>12</sub>	D <sub>3</sub> F <sub>4</sub>	Application of liquid jeevamrutha @ 1500 l/ha once in four week (one month)
T <sub>13</sub>	INM	Integrated nutrient management (INM) (50% RDN through FYM (9 t/ha) + 50% RDN through inorganic fertilizer (45kg N/ha) + P @ 120 kg/ha + K @ 60 kg/ha + 20 t FYM/ha + Azospirillum @ 2.5 kg/ha + PSB @ 2.5 kg/ha)
T <sub>14</sub>	RPP	Recommended package of practice (RPP) (NPK @ 90:120:60 kg/ha + 20t FYM/ha)

Note: Beejamrutha treatment, ghanajeevamrutha @ 1000kg per hectare at the time of planting and application of mulch was common to all the treatments except T<sub>13</sub> and T<sub>14</sub>, treatments were imposed after 15 DAT

### 3. RESULTS AND DISCUSSION

The dosage of liquid jeevamrutha @ 1500 liters per hectare (D<sub>3</sub>) and frequency of liquid jeevamrutha applied at 7 days interval (F<sub>1</sub>) was found better in influencing the growth parameters significantly than other dosage and frequency levels. Further, the interaction treatment D<sub>3</sub>F<sub>1</sub> (liquid jeevamrutha applied @ 1500 l/ha applied at 7 days interval) resulted in higher values for all the growth parameters. Based on the pooled analysis of the two years data it was observed that, significantly higher plant height (43.28 and 56.27 cm at 60 and 90 DAT, respectively), plant spread N-S (24.92 and 28.51 cm at 60 and 90 DAT, respectively), plant spread E-W (24.19 and 25.51 cm at 60 and 90 DAT, respectively) and number of primary branches per plant (6.23 and 9.34 at 60 and 90 DAT, respectively) was recorded in D<sub>3</sub> (liquid jeevamrutha applied @ 1500 l/ha). Similarly, frequency of liquid jeevamrutha applied at 7 days interval (F<sub>1</sub>) also recorded higher plant height (46.13 and 56.64 cm at 60 and 90 DAT, respectively), plant spread in both North-South (25.19 and 31.86 cm at 60 and 90 DAT, respectively) and East-West directions (24.81 and 26.68 cm at 60 and 90 DAT, respectively) and number of primary branches per plant (7.03 and 10.23 at 60 and 90 DAT, respectively). The interaction of dosage and frequency of liquid jeevamrutha applied @ 1500 liter per hectare at an interval of 7 days (D<sub>3</sub>F<sub>1</sub>) resulted in significantly higher plant height (47.98 cm and 58.78 cm at 60 and 90 DAT, respectively), plant spread (N-S) i.e., 17.90 26.35 and 31.31 cm at 60 and 90 DAT, respectively, plant spread E-W i.e., 27.10 and 30.35 cm at 60

and 90 DAT, respectively and number of primary branches (7.45 and 10.28 at 60 and 90 DAT, respectively) in pooled data.

Among the different interaction treatments, the plants applied with liquid jeevamrutha @ 1500 liter per hectare at 7 days interval (D<sub>3</sub>F<sub>1</sub>) was observed maximum vegetative parameters like plant height, plant spread and number of primary branches. The higher growth parameters in the above treatment might be attributed to solubilization of nutrients in soil and absorption of the same and further maintenance of good soil moisture level due to the application of jeevamrutha. Similar observation was also made by Yogananda *et al.* [2] in cowpea and Biradar [3] in French bean. Further higher doses of liquid jeevamrutha applied at regular interval stimulate the activities of micro organisms to release the nitrogen in a synchronous manner, which might have stirred the cellular activity. Further, more the presence of growth promoting hormones viz., IAA and GA<sub>3</sub> in jeevamrutha might have favored rapid cell division and multiplication contributing to increased growth parameters among the different levels of liquid jeevamrutha which is in accordance with the results of Palekar [4] and Neelima and Sreenivas, [5]. It also enhances the microbial activity in soil and ultimately ensuring the availability and uptake of nutrients by the crop. These results are in accordance with the findings of earlier study done by Kumbar and Deavakumar [6] in French bean and Gangadhar *et al.* [7] in chilli (Table 2).

When interaction treatments were compared with RPP (recommended package of practice) and

INM (integrated nutrient management) it was revealed that, INM recorded significantly highest plant (30.31, 68.43 and 77.80 cm at 30, 60 and 90 DAT, respectively), plant spread N-S *i.e.*, 30.76, 35.73 and 42.66 cm at 30, 60 and 90 DAT, respectively, plant spread (E-W) *i.e.*, 24.97, 35.98 and 40.97 cm at 30, 60 and 90 DAT, respectively, number of primary branches (5.51, 10.08 and 13.37 at 30, 60 and 90 DAT, respectively) and also in RPP plant height (28.37, 63.68 and 74.61 cm at 30, 60 and 90 DAT, respectively), plant spread in both North-South (23.72, 33.67 and 38.26 cm at 30, 60 and 90 DAT, respectively), East-West (23.82, 34.71 and 36.10 cm at 30, 60 and 90 DAT, respectively) directions and number of primary branches per plant (5.01, 9.42 and 12.71 at 30, 60 and 90 DAT, respectively) in pooled data. In INM and RPP, significantly maximum values recorded for all the growth parameters could be attributed to the higher uptake of nutrients, particularly nitrogen [8]. Improved growth parameters in the present study might be due to the fact that FYM besides supplying N, P and K improves the soil condition which enhances the source to sink relationship and also makes unavailable sources of elemental nitrogen, bound phosphates, micronutrients and decomposed plant residues into available form to facilitate the plants to absorb the nutrients. These results are in agreement with the findings of earlier works done by Singh *et al.* [9] and Siddappa [10].

Flower yield parameters like number of flowers per plant and flower yield differed significantly. Significantly maximum number of flowers per plant and flower yield (32.78, 93.09 g/plant, 4.52 kg/plot and 6.70 t/ha, respectively) were recorded in D<sub>3</sub> (liquid jeevamrutha applied @ 1500 l/ha), F<sub>1</sub> (frequency of liquid jeevamrutha applied at 7 days interval) *i.e.*, 35.95, 105.13 gram per plant, 4.76 kg per plot and 7.07 tonnes per hectare and in interaction treatment D<sub>3</sub>F<sub>1</sub> (liquid jeevamrutha applied @ 1500 l/ha at an interval of 7 days) *i.e.*, 38.42, 111.39 gram per plant, 5.41 kg per plot and 8.06 tonnes per hectare, respectively in pooled data (Table 3). This might be due to favourable effects of macro and micronutrients, which helps in better availability of nutrients throughout the crop growth which might be the result of improved microbial activity in the soil. These findings are in accordance with Kasbe *et al.* [11]. The effects of interaction of dosage and frequencies of liquid

jeevamrutha was significant and was observed that higher dosage of liquid jeevamrutha applied @ 1500 liter per hectare at lesser intervals (7 days interval) produced more China aster flower yield than other levels. The higher microbial activities in the rhizosphere soil and further higher enzymatic activities might have resulted in higher release of adsorbed nutrients and their uptake and also higher vegetative growth by crop might be the reason for higher flower yield in the above treatments. This jeevamrutha might have helped in decomposition of organic matter and release of available nutrients for easy uptake and utilization by the plants. These findings are in agreement with earlier studies done by Fazeel *et al.* [12], Jhade *et al.* [13] and Chaithra and sujith [14].

The interaction treatments were compared with RPP and INM showed that, significantly maximum number of flowers per plant and flower yield were recorded in INM (49.00, 147.82 g/plant, 7.27 kg/plot and 10.76 t/ha, respectively) and RPP (47.11, 137.86 g/plant, 6.84 kg/plot and 10.09 t/ha, respectively) in pooled data. In INM more yield was due to indirect effect of more number of branches as estimated and developed by the influence of inorganic fertilizers along with organic manures and biofertilizers and also may be due to proper nitrogen, phosphorus and potassium assimilation from the NPK in association with more nitrogen fixing and phosphorus solubilizing proficiency and secretion of hormones by the biofertilizers similar results were obtained by Kumar *et al.* [15] in China aster. The increased yield in RPP was due to the supply of nutrients through FYM and chemical fertilizer might be attributed to the quick release and availability of nutrients in required quantity with the application of fertilizers. Further, FYM acts as store house for various micro and macro nutrients that are released during the process of mineralization. The results are supported with the findings of Mahapatra *et al.* [16] and Gorabal [17].

The dosage and frequency of liquid jeevamrutha also influenced the flower diameter (cm) and individual flower weight (g) significantly. The dosage D<sub>3</sub> (liquid jeevamrutha applied @ 1500 l/ha) *i.e.*, 5.10 cm and 2.86 g, respectively, frequency F<sub>1</sub> (frequency of liquid jeevamrutha applied at 7 days interval) *i.e.*, 5.51 cm and 2.97 g, respectively and their interaction D<sub>3</sub>F<sub>1</sub> (liquid

**Table 2. Growth parameters in China aster var. AAC-1 at different growth stages as influenced by dosage and frequency of liquid jeevamrutha**

Treatment	Pooled data											
	Plant height (cm)			Plant spread (cm) (North-South)			Plant spread (cm) (East-West)			Number of primary branches		
	30 DAT	60 DAT	90 DAT	30 DAT	60 DAT	90 DAT	30 DAT	60 DAT	90 DAT	30 DAT	60 DAT	90 DAT
<b>Factor A: Dosage of liquid jeevamrutha (D)</b>												
D <sub>1</sub> : Jeevamrutha@500 l/ha	17.58	40.68 <sup>c</sup>	50.95 <sup>c</sup>	15.34	22.56 <sup>c</sup>	28.36 <sup>b</sup>	11.50	19.70 <sup>c</sup>	21.47 <sup>c</sup>	3.01	5.48 <sup>c</sup>	8.04 <sup>b</sup>
D <sub>2</sub> : Jeevamrutha@1000 l/ha	17.85	40.74 <sup>b</sup>	52.70 <sup>b</sup>	15.80	23.65 <sup>b</sup>	28.48 <sup>a</sup>	12.79	22.99 <sup>b</sup>	24.12 <sup>b</sup>	3.11	5.94 <sup>b</sup>	8.73 <sup>ab</sup>
D <sub>3</sub> : Jeevamrutha@1500 l/ha	19.00	43.28 <sup>a</sup>	56.27 <sup>a</sup>	16.57	24.92 <sup>a</sup>	28.51 <sup>a</sup>	13.50	24.19 <sup>a</sup>	25.51 <sup>a</sup>	3.16	6.23 <sup>a</sup>	9.34 <sup>a</sup>
<b>S.Em±</b>	<b>1.01</b>	<b>0.18</b>	<b>0.41</b>	<b>0.75</b>	<b>0.12</b>	<b>0.05</b>	<b>0.71</b>	<b>0.14</b>	<b>0.15</b>	<b>0.11</b>	<b>0.03</b>	<b>0.06</b>
<b>C.D. @ 5%</b>	<b>NS</b>	<b>0.52</b>	<b>1.21</b>	<b>NS</b>	<b>0.35</b>	<b>0.16</b>	<b>NS</b>	<b>0.40</b>	<b>0.44</b>	<b>NS</b>	<b>0.08</b>	<b>0.18</b>
<b>Factor B: Frequency of liquid jeevamrutha application (F)</b>												
F <sub>1</sub> : Once in a week (7 days)	19.92	46.13 <sup>a</sup>	56.64 <sup>a</sup>	17.48	25.19 <sup>a</sup>	31.86 <sup>a</sup>	14.14	24.81 <sup>a</sup>	26.68 <sup>a</sup>	3.74	7.03 <sup>a</sup>	10.23 <sup>a</sup>
F <sub>2</sub> : Once in two weeks (15 days)	18.50	42.73 <sup>b</sup>	53.63 <sup>b</sup>	16.38	24.38 <sup>b</sup>	28.90 <sup>b</sup>	13.08	22.56 <sup>b</sup>	24.09 <sup>b</sup>	3.19	5.75 <sup>b</sup>	8.72 <sup>b</sup>
F <sub>3</sub> : Once in three week (21 days)	17.33	39.29 <sup>c</sup>	51.92 <sup>c</sup>	15.77	23.13 <sup>c</sup>	27.70 <sup>b</sup>	11.91	21.52 <sup>c</sup>	22.78 <sup>c</sup>	2.83	5.40 <sup>c</sup>	8.21 <sup>c</sup>
F <sub>4</sub> : Once in a month (30 days)	16.83	38.14 <sup>d</sup>	51.16 <sup>c</sup>	13.99	22.16 <sup>d</sup>	26.23 <sup>c</sup>	11.24	20.29 <sup>d</sup>	21.25 <sup>d</sup>	2.64	5.37 <sup>d</sup>	7.68 <sup>d</sup>
<b>S.Em±</b>	<b>1.17</b>	<b>0.20</b>	<b>0.48</b>	<b>1.12</b>	<b>0.14</b>	<b>0.53</b>	<b>0.85</b>	<b>0.16</b>	<b>0.17</b>	<b>0.30</b>	<b>0.03</b>	<b>0.07</b>
<b>C.D. @ 5%</b>	<b>NS</b>	<b>0.60</b>	<b>1.39</b>	<b>NS</b>	<b>0.41</b>	<b>1.59</b>	<b>NS</b>	<b>0.46</b>	<b>0.51</b>	<b>NS</b>	<b>0.11</b>	<b>0.21</b>
<b>Interaction (D x F)</b>												
D <sub>1</sub> F <sub>1</sub>	18.90	44.96 <sup>abc</sup>	54.50 <sup>abc</sup>	17.64	23.77 <sup>b</sup>	27.95 <sup>a</sup>	12.67	22.38 <sup>e</sup>	24.12 <sup>de</sup>	3.61	6.63 <sup>cde</sup>	8.45 <sup>bc</sup>
D <sub>1</sub> F <sub>2</sub>	17.90	39.74 <sup>bc</sup>	51.15 <sup>de</sup>	15.96	23.79 <sup>cde</sup>	26.90 <sup>bc</sup>	12.25	19.35 <sup>g</sup>	22.12 <sup>g</sup>	3.06	5.05 <sup>de</sup>	8.40 <sup>bc</sup>
D <sub>1</sub> F <sub>3</sub>	17.42	37.11 <sup>c</sup>	50.40 <sup>de</sup>	14.90	21.94 <sup>f</sup>	27.24 <sup>cdef</sup>	10.98	18.98 <sup>g</sup>	20.52 <sup>h</sup>	2.70	4.81 <sup>e</sup>	7.87 <sup>cd</sup>
D <sub>1</sub> F <sub>4</sub>	16.12	35.80 <sup>d</sup>	47.61 <sup>e</sup>	13.41	20.80 <sup>f</sup>	24.80 <sup>def</sup>	10.08	18.08 <sup>h</sup>	19.12 <sup>i</sup>	2.70	4.72 <sup>e</sup>	7.78 <sup>d</sup>
D <sub>2</sub> F <sub>1</sub>	20.15	45.79 <sup>a</sup>	56.49 <sup>a</sup>	17.38	25.46 <sup>a</sup>	29.80 <sup>a</sup>	14.20	24.93 <sup>b</sup>	26.57 <sup>b</sup>	3.61	7.00 <sup>a</sup>	9.39 <sup>a</sup>
D <sub>2</sub> F <sub>2</sub>	17.94	40.00 <sup>ab</sup>	54.34	16.96	24.22 <sup>ab</sup>	27.15 <sup>b</sup>	13.20	23.80 <sup>cd</sup>	24.82 <sup>cd</sup>	3.37	6.21 <sup>ab</sup>	7.66 <sup>bc</sup>
D <sub>2</sub> F <sub>3</sub>	16.76	39.20 <sup>ab</sup>	51.09 <sup>ab</sup>	16.30	22.81 <sup>cd</sup>	27.43 <sup>bcd</sup>	12.03	22.37 <sup>e</sup>	23.58 <sup>ef</sup>	2.83	5.98 <sup>bc</sup>	7.21 <sup>cd</sup>
D <sub>2</sub> F <sub>4</sub>	16.55	38.41 <sup>bc</sup>	48.36 <sup>bcd</sup>	14.57	22.13 <sup>f</sup>	24.94 <sup>bc</sup>	11.72	20.87 <sup>f</sup>	21.52 <sup>g</sup>	2.66	5.95 <sup>c</sup>	6.35 <sup>cd</sup>
D <sub>3</sub> F <sub>1</sub>	20.69	47.98 <sup>a</sup>	58.78 <sup>a</sup>	17.90	26.35 <sup>a</sup>	31.31 <sup>a</sup>	15.57	27.10 <sup>a</sup>	30.35 <sup>a</sup>	3.90	7.45 <sup>a</sup>	10.28 <sup>a</sup>
D <sub>3</sub> F <sub>2</sub>	19.67	43.54 <sup>ab</sup>	55.40 <sup>abc</sup>	17.36	26.26 <sup>a</sup>	28.21 <sup>bcd</sup>	13.80	24.53 <sup>bc</sup>	25.33 <sup>c</sup>	3.13	5.81 <sup>bc</sup>	7.93 <sup>bc</sup>
D <sub>3</sub> F <sub>3</sub>	17.82	41.14 <sup>abc</sup>	54.45 <sup>bcd</sup>	16.63	26.34 <sup>ef</sup>	27.16 <sup>ef</sup>	12.72	23.20 <sup>d</sup>	24.25 <sup>de</sup>	2.95	5.38 <sup>cd</sup>	7.65 <sup>d</sup>
D <sub>3</sub> F <sub>4</sub>	17.83	40.49 <sup>abc</sup>	51.89 <sup>cde</sup>	14.39	23.54 <sup>def</sup>	25.29 <sup>f</sup>	11.93	21.92 <sup>e</sup>	23.12 <sup>f</sup>	2.58	5.03 <sup>de</sup>	6.50 <sup>d</sup>
<b>S.Em±</b>	<b>2.02</b>	<b>0.80</b>	<b>0.82</b>	<b>1.75</b>	<b>0.24</b>	<b>0.43</b>	<b>1.95</b>	<b>0.27</b>	<b>0.30</b>	<b>0.40</b>	<b>0.11</b>	<b>0.28</b>
<b>C.D. @ 5%</b>	<b>NS</b>	<b>2.56</b>	<b>2.41</b>	<b>NS</b>	<b>0.70</b>	<b>1.29</b>	<b>NS</b>	<b>0.79</b>	<b>0.88</b>	<b>NS</b>	<b>0.35</b>	<b>0.84</b>
INM	30.16	68.43	77.80	30.76	35.73	42.66	24.97	35.98	40.97	5.51	10.08	13.37
RPP	28.37	63.68	74.61	23.72	33.67	38.26	23.82	34.71	36.10	5.01	9.42	12.71
<b>S.Em±</b>	<b>1.88</b>	<b>1.65</b>	<b>1.30</b>	<b>1.43</b>	<b>0.32</b>	<b>0.27</b>	<b>1.30</b>	<b>0.38</b>	<b>1.67</b>	<b>0.22</b>	<b>0.40</b>	<b>0.16</b>
<b>C.D. @ 5%</b>	<b>5.47</b>	<b>4.80</b>	<b>3.90</b>	<b>4.16</b>	<b>0.94</b>	<b>0.78</b>	<b>3.79</b>	<b>1.11</b>	<b>5.01</b>	<b>0.66</b>	<b>1.20</b>	<b>0.47</b>

Note: INM: Integrated nutrient management (50% RDN through FYM (9 t/ha) + 50% RDN through inorganic fertilizer (45 kg N/ha) + 120 kg P/ha + K @ 60 Kg/ha + 20 t FYM/ha + Azospirillum @ 2.5 kg/ha + PSB @ 2.5 kg/ha RPP: Recommended package of practice (NPK @ 90:120:60 kg/ha + 20 t FYM/ha) DAT: Days after transplanting NS: Non significant

**Table 3. Yield parameters in China aster var. AAC-1 as influenced by dosage and frequency of liquid jeevamrutha**

Treatment	Number of flowers/plant			Flower yield/plant (g)			Flower yield/ plot (kg)			Flower yield/ hectare (t)		
	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled
<b>Factor A: Dosage of liquid jeevamrutha (D)</b>												
D <sub>1</sub> : Jeevamrutha@ 500 l/ha	31.09 <sup>c</sup>	28.01 <sup>c</sup>	29.55 <sup>c</sup>	82.64 <sup>c</sup>	79.76 <sup>b</sup>	81.20 <sup>c</sup>	3.82 <sup>c</sup>	3.46 <sup>c</sup>	3.65 <sup>c</sup>	5.67 <sup>c</sup>	5.13 <sup>c</sup>	5.40 <sup>c</sup>
D <sub>2</sub> : Jeevamrutha@1000 l/ha	33.25 <sup>b</sup>	29.27 <sup>b</sup>	31.26 <sup>b</sup>	93.14 <sup>b</sup>	88.72 <sup>a</sup>	90.93 <sup>b</sup>	4.28 <sup>b</sup>	3.70 <sup>b</sup>	4.00 <sup>b</sup>	6.38 <sup>b</sup>	5.92 <sup>b</sup>	6.15 <sup>b</sup>
D <sub>3</sub> : Jeevamrutha@1500 l/ha	35.80 <sup>a</sup>	29.77 <sup>a</sup>	32.78 <sup>a</sup>	97.78 <sup>a</sup>	89.01 <sup>a</sup>	93.09 <sup>a</sup>	4.71 <sup>a</sup>	4.31 <sup>a</sup>	4.52 <sup>a</sup>	6.98 <sup>a</sup>	6.41 <sup>a</sup>	6.70 <sup>a</sup>
<b>S.Em±</b>	<b>0.17</b>	<b>0.28</b>	<b>0.14</b>	<b>0.91</b>	<b>1.12</b>	<b>1.01</b>	<b>0.07</b>	<b>0.05</b>	<b>0.05</b>	<b>0.09</b>	<b>0.10</b>	<b>0.07</b>
<b>C.D. @ 5%</b>	<b>0.49</b>	<b>0.83</b>	<b>0.41</b>	<b>2.67</b>	<b>3.29</b>	<b>2.97</b>	<b>0.20</b>	<b>0.15</b>	<b>0.13</b>	<b>0.25</b>	<b>0.29</b>	<b>0.20</b>
<b>Factor B: Frequency of liquid jeevamrutha application (F)</b>												
F <sub>1</sub> : Once in a week (7 days)	37.77 <sup>a</sup>	34.13 <sup>a</sup>	35.95 <sup>a</sup>	107.21 <sup>a</sup>	103.06 <sup>a</sup>	105.13 <sup>a</sup>	4.92 <sup>a</sup>	4.60 <sup>a</sup>	4.76 <sup>a</sup>	7.29 <sup>a</sup>	6.84 <sup>a</sup>	7.07 <sup>a</sup>
F <sub>2</sub> : Once in two weeks (15 days)	34.24 <sup>b</sup>	28.49 <sup>b</sup>	31.36 <sup>b</sup>	91.72 <sup>b</sup>	86.04 <sup>b</sup>	88.88 <sup>b</sup>	4.37 <sup>b</sup>	3.84 <sup>b</sup>	4.11 <sup>b</sup>	6.49 <sup>b</sup>	5.70 <sup>b</sup>	6.09 <sup>b</sup>
F <sub>3</sub> : Once in three week (21 days)	31.79 <sup>c</sup>	27.07 <sup>c</sup>	29.43 <sup>c</sup>	85.32 <sup>c</sup>	78.32 <sup>c</sup>	81.82 <sup>c</sup>	4.08 <sup>c</sup>	3.53 <sup>c</sup>	3.81 <sup>c</sup>	6.07 <sup>c</sup>	5.51 <sup>c</sup>	5.79 <sup>c</sup>
F <sub>4</sub> : Once in a month (30 days)	29.73 <sup>d</sup>	26.39 <sup>d</sup>	28.06 <sup>c</sup>	80.52 <sup>d</sup>	75.10 <sup>c</sup>	77.81 <sup>d</sup>	3.74 <sup>d</sup>	3.34 <sup>d</sup>	3.54 <sup>d</sup>	5.56 <sup>d</sup>	5.26 <sup>c</sup>	5.41 <sup>d</sup>
<b>S.Em±</b>	<b>0.19</b>	<b>0.33</b>	<b>0.16</b>	<b>1.05</b>	<b>1.30</b>	<b>1.17</b>	<b>0.08</b>	<b>0.06</b>	<b>0.05</b>	<b>0.10</b>	<b>0.12</b>	<b>0.08</b>
<b>C.D. @ 5%</b>	<b>0.56</b>	<b>0.95</b>	<b>1.47</b>	<b>3.08</b>	<b>3.80</b>	<b>3.43</b>	<b>0.23</b>	<b>0.17</b>	<b>0.15</b>	<b>0.29</b>	<b>0.34</b>	<b>0.23</b>
<b>Interaction (D x F)</b>												
D <sub>1</sub> F <sub>1</sub>	34.73 <sup>bc</sup>	31.97 <sup>a</sup>	33.35 <sup>bc</sup>	96.57 <sup>b</sup>	93.11 <sup>b</sup>	95.51 <sup>ab</sup>	4.23 <sup>de</sup>	3.90 <sup>bc</sup>	4.07 <sup>de</sup>	6.27 <sup>de</sup>	5.78 <sup>bc</sup>	6.02 <sup>de</sup>
D <sub>1</sub> F <sub>2</sub>	30.33 <sup>cd</sup>	27.70 <sup>bc</sup>	29.02 <sup>cd</sup>	78.89 <sup>cde</sup>	76.00 <sup>c</sup>	77.45 <sup>bc</sup>	4.10 <sup>def</sup>	3.63 <sup>cd</sup>	3.87 <sup>e</sup>	6.12 <sup>def</sup>	5.68 <sup>cd</sup>	5.75 <sup>e</sup>
D <sub>1</sub> F <sub>3</sub>	29.90 <sup>def</sup>	26.80 <sup>c</sup>	28.35 <sup>d</sup>	77.73 <sup>ef</sup>	75.61 <sup>d</sup>	76.67 <sup>cd</sup>	3.77 <sup>f</sup>	3.30 <sup>de</sup>	3.53 <sup>f</sup>	5.90 <sup>f</sup>	5.40 <sup>de</sup>	5.65 <sup>f</sup>
D <sub>1</sub> F <sub>4</sub>	29.40 <sup>f</sup>	25.60 <sup>c</sup>	27.50 <sup>fg</sup>	76.39 <sup>f</sup>	74.00 <sup>d</sup>	75.20 <sup>cd</sup>	3.20 <sup>g</sup>	3.03 <sup>ef</sup>	3.12 <sup>g</sup>	4.74 <sup>g</sup>	4.49 <sup>ef</sup>	4.62 <sup>g</sup>
D <sub>2</sub> F <sub>1</sub>	38.58 <sup>b</sup>	33.60 <sup>b</sup>	36.09 <sup>ab</sup>	110.61 <sup>a</sup>	106.40 <sup>a</sup>	108.51 <sup>a</sup>	5.00 <sup>b</sup>	4.60 <sup>b</sup>	4.80 <sup>b</sup>	7.41 <sup>b</sup>	6.81 <sup>a</sup>	7.11 <sup>b</sup>
D <sub>2</sub> F <sub>2</sub>	34.17 <sup>c</sup>	29.00 <sup>bc</sup>	31.59 <sup>def</sup>	96.22 <sup>b</sup>	93.45 <sup>b</sup>	94.95 <sup>ab</sup>	4.20 <sup>de</sup>	3.80 <sup>bc</sup>	4.00 <sup>de</sup>	6.22 <sup>de</sup>	5.63 <sup>bc</sup>	5.93 <sup>de</sup>
D <sub>2</sub> F <sub>3</sub>	30.47 <sup>cde</sup>	27.13 <sup>cd</sup>	28.80 <sup>cde</sup>	85.56 <sup>def</sup>	79.77 <sup>c</sup>	82.67 <sup>bc</sup>	3.96 <sup>gh</sup>	3.27 <sup>fg</sup>	3.60 <sup>h</sup>	5.97 <sup>gh</sup>	5.67 <sup>eg</sup>	5.82 <sup>h</sup>
D <sub>2</sub> F <sub>4</sub>	29.95 <sup>de</sup>	27.37 <sup>cd</sup>	28.66 <sup>d</sup>	80.18 <sup>f</sup>	75.26 <sup>d</sup>	77.72 <sup>cd</sup>	3.95 <sup>gh</sup>	3.19 <sup>g</sup>	3.57 <sup>h</sup>	5.96 <sup>h</sup>	5.60 <sup>d</sup>	5.78 <sup>h</sup>
D <sub>3</sub> F <sub>1</sub>	40.00 <sup>a</sup>	36.83 <sup>a</sup>	38.42 <sup>a</sup>	113.44 <sup>a</sup>	109.33 <sup>a</sup>	111.39 <sup>a</sup>	5.53 <sup>a</sup>	5.30 <sup>a</sup>	5.41 <sup>a</sup>	8.20 <sup>a</sup>	7.92 <sup>a</sup>	8.06 <sup>a</sup>
D <sub>3</sub> F <sub>2</sub>	38.21 <sup>b</sup>	28.77 <sup>bc</sup>	33.49 <sup>def</sup>	100.04 <sup>b</sup>	88.67 <sup>b</sup>	94.36 <sup>ab</sup>	4.80 <sup>bc</sup>	4.10 <sup>b</sup>	4.45 <sup>c</sup>	7.11 <sup>bc</sup>	6.07 <sup>b</sup>	6.59 <sup>c</sup>
D <sub>3</sub> F <sub>3</sub>	35.00 <sup>c</sup>	27.27 <sup>c</sup>	31.14 <sup>cde</sup>	92.67 <sup>c</sup>	79.56 <sup>c</sup>	86.12 <sup>bc</sup>	4.50 <sup>cd</sup>	4.03 <sup>bc</sup>	4.27 <sup>cd</sup>	6.67 <sup>cd</sup>	5.98 <sup>bc</sup>	6.32 <sup>cd</sup>
D <sub>3</sub> F <sub>4</sub>	30.00 <sup>ef</sup>	26.21 <sup>cd</sup>	28.11 <sup>efg</sup>	85.00 <sup>cd</sup>	76.04 <sup>c</sup>	80.52 <sup>bc</sup>	4.03 <sup>ef</sup>	3.83 <sup>bc</sup>	3.93 <sup>e</sup>	5.98 <sup>ef</sup>	5.68 <sup>bc</sup>	5.83 <sup>e</sup>
<b>S.Em±</b>	<b>0.33</b>	<b>0.56</b>	<b>0.28</b>	<b>1.82</b>	<b>2.24</b>	<b>2.02</b>	<b>0.13</b>	<b>0.10</b>	<b>0.09</b>	<b>0.17</b>	<b>0.20</b>	<b>0.13</b>
<b>C.D. @ 5%</b>	<b>0.97</b>	<b>1.65</b>	<b>0.82</b>	<b>5.34</b>	<b>6.58</b>	<b>5.93</b>	<b>0.39</b>	<b>0.30</b>	<b>0.27</b>	<b>0.50</b>	<b>0.58</b>	<b>0.39</b>
INM	53.00	45.00	49.00	152.32	143.32	147.82	7.52	7.02	7.27	11.12	10.40	10.76
RPP	51.00	43.21	47.11	145.36	130.36	137.86	7.15	6.53	6.84	10.38	9.67	10.09
<b>S.Em±</b>	<b>0.55</b>	<b>1.69</b>	<b>0.79</b>	<b>1.86</b>	<b>2.10</b>	<b>1.95</b>	<b>0.22</b>	<b>0.17</b>	<b>0.13</b>	<b>0.16</b>	<b>0.19</b>	<b>0.25</b>
<b>C.D. @ 5%</b>	<b>1.59</b>	<b>4.92</b>	<b>2.30</b>	<b>5.40</b>	<b>6.11</b>	<b>5.66</b>	<b>0.63</b>	<b>0.35</b>	<b>0.37</b>	<b>0.47</b>	<b>0.55</b>	<b>0.76</b>

Note: INM: Integrated nutrient management (50% RDN through FYM (9 t/ha) + 50% RDN through inorganic fertilizer (45 kg N/ha) + 120 kg P/ha + K @ 60 kg/ha + 20 t FYM/ha + Azospirillum @ 2.5 kg/ha + PSB @ 2.5 kg/ha RPP: Recommended package of practice (NPK @ 90:120:60 kg/ha +20 t FYM/ha)

**Table 4. Flower diameter, individual flower weight, shelf life and vase life in China aster var. AAC-1 as influenced by dosage and frequency of liquid jeevamrutha**

Treatment	Flower diameter (cm)			Individual flower weight (g)			Shelf life (Days)			Vase life (Days)		
	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled
<b>Factor A: Dosage of liquid jeevamrutha (D)</b>												
D <sub>1</sub> : Jeevamrutha@500 l/ha	4.58 <sup>b</sup>	5.13 <sup>b</sup>	4.85 <sup>b</sup>	2.52 <sup>c</sup>	2.56 <sup>c</sup>	2.54 <sup>c</sup>	2.62	2.72 <sup>b</sup>	2.66 <sup>c</sup>	7.34 <sup>c</sup>	7.21 <sup>c</sup>	7.28 <sup>c</sup>
D <sub>2</sub> : Jeevamrutha@1000 l/ha	4.75 <sup>a</sup>	5.27 <sup>a</sup>	5.09 <sup>a</sup>	2.64 <sup>b</sup>	2.77 <sup>b</sup>	2.71 <sup>b</sup>	2.57	2.79 <sup>b</sup>	2.68 <sup>b</sup>	7.44 <sup>b</sup>	7.67 <sup>b</sup>	7.55 <sup>b</sup>
D <sub>3</sub> : Jeevamrutha@1500 l/ha	4.95 <sup>a</sup>	5.27 <sup>a</sup>	5.10 <sup>a</sup>	2.77 <sup>a</sup>	2.96 <sup>a</sup>	2.86 <sup>a</sup>	2.62	3.17 <sup>a</sup>	2.89 <sup>a</sup>	7.79 <sup>a</sup>	7.91 <sup>a</sup>	7.85 <sup>a</sup>
<b>S.Em±</b>	<b>0.10</b>	<b>0.04</b>	<b>0.06</b>	<b>0.02</b>	<b>0.02</b>	<b>0.01</b>	<b>0.10</b>	<b>0.05</b>	<b>0.04</b>	<b>0.03</b>	<b>0.08</b>	<b>0.04</b>
<b>C.D. @ 5%</b>	<b>0.29</b>	<b>0.13</b>	<b>0.20</b>	<b>0.06</b>	<b>0.07</b>	<b>0.04</b>	<b>NS</b>	<b>0.15</b>	<b>0.11</b>	<b>0.09</b>	<b>0.24</b>	<b>0.11</b>
<b>Factor B: Frequency of liquid jeevamrutha application (F)</b>												
F <sub>1</sub> : Once in a week (7 days)	5.40 <sup>a</sup>	5.62 <sup>a</sup>	5.51 <sup>a</sup>	2.82 <sup>a</sup>	3.11 <sup>a</sup>	2.97 <sup>a</sup>	2.89 <sup>a</sup>	3.57 <sup>a</sup>	3.23 <sup>a</sup>	8.23 <sup>a</sup>	8.40 <sup>a</sup>	8.32 <sup>a</sup>
F <sub>2</sub> : Once in two weeks (15 days)	5.15 <sup>b</sup>	5.39 <sup>b</sup>	5.27 <sup>b</sup>	2.72 <sup>b</sup>	2.81 <sup>b</sup>	2.76 <sup>b</sup>	2.67 <sup>a</sup>	3.00 <sup>b</sup>	2.83 <sup>b</sup>	7.61 <sup>b</sup>	7.53 <sup>b</sup>	7.57 <sup>b</sup>
F <sub>3</sub> : Once in three week (21 days)	4.67 <sup>c</sup>	5.26 <sup>b</sup>	4.96 <sup>c</sup>	2.59 <sup>c</sup>	2.64 <sup>c</sup>	2.62 <sup>c</sup>	2.51 <sup>b</sup>	2.57 <sup>c</sup>	2.54 <sup>c</sup>	7.32 <sup>c</sup>	7.44 <sup>b</sup>	7.38 <sup>c</sup>
F <sub>4</sub> : Once in a month (30 days)	3.82 <sup>d</sup>	4.63 <sup>c</sup>	4.22 <sup>d</sup>	2.45 <sup>d</sup>	2.51 <sup>d</sup>	2.48 <sup>d</sup>	2.35 <sup>b</sup>	2.45 <sup>c</sup>	2.40 <sup>d</sup>	6.95 <sup>d</sup>	7.04 <sup>c</sup>	7.00 <sup>d</sup>
<b>S.Em±</b>	<b>0.11</b>	<b>0.05</b>	<b>0.07</b>	<b>0.02</b>	<b>0.03</b>	<b>0.02</b>	<b>0.12</b>	<b>0.06</b>	<b>0.04</b>	<b>0.04</b>	<b>0.09</b>	<b>0.04</b>
<b>C.D. @ 5%</b>	<b>0.33</b>	<b>0.15</b>	<b>0.20</b>	<b>0.07</b>	<b>0.08</b>	<b>0.05</b>	<b>0.35</b>	<b>0.17</b>	<b>0.13</b>	<b>0.11</b>	<b>0.28</b>	<b>0.13</b>
<b>Interaction (D x F)</b>												
D <sub>1</sub> F <sub>1</sub>	5.09 <sup>ab</sup>	5.53 <sup>abc</sup>	5.31 <sup>ab</sup>	2.76 <sup>a</sup>	3.01 <sup>b</sup>	2.88 <sup>a</sup>	2.77	3.23 <sup>bcd</sup>	3.00 <sup>b</sup>	8.02 <sup>abc</sup>	7.77 <sup>ab</sup>	7.89 <sup>ab</sup>
D <sub>1</sub> F <sub>2</sub>	5.03 <sup>abc</sup>	5.29 <sup>bc</sup>	5.16 <sup>abcd</sup>	2.57 <sup>abc</sup>	2.56 <sup>d</sup>	2.57 <sup>cd</sup>	2.60	2.90 <sup>de</sup>	2.75 <sup>bc</sup>	7.35 <sup>cde</sup>	6.87 <sup>cde</sup>	7.11 <sup>de</sup>
D <sub>1</sub> F <sub>3</sub>	4.53 <sup>abcd</sup>	5.05 <sup>c</sup>	4.80 <sup>def</sup>	2.42 <sup>bc</sup>	2.36 <sup>d</sup>	2.38 <sup>d</sup>	2.57	2.50 <sup>ef</sup>	2.53 <sup>cd</sup>	7.20 <sup>cde</sup>	7.27 <sup>cdef</sup>	7.23 <sup>e</sup>
D <sub>1</sub> F <sub>4</sub>	3.68 <sup>cde</sup>	4.63 <sup>f</sup>	4.16 <sup>ef</sup>	2.35 <sup>bc</sup>	2.32 <sup>d</sup>	2.34 <sup>d</sup>	2.53	2.25 <sup>f</sup>	2.39 <sup>cd</sup>	6.83 <sup>e</sup>	6.97 <sup>efg</sup>	6.90 <sup>e</sup>
D <sub>2</sub> F <sub>1</sub>	5.22 <sup>abc</sup>	5.62 <sup>ab</sup>	5.44 <sup>a</sup>	2.84 <sup>a</sup>	3.11 <sup>a</sup>	2.97 <sup>a</sup>	2.83	3.67 <sup>a</sup>	3.25 <sup>a</sup>	8.05 <sup>ab</sup>	8.50 <sup>abc</sup>	8.28 <sup>b</sup>
D <sub>2</sub> F <sub>2</sub>	5.01 <sup>bc</sup>	5.42 <sup>cde</sup>	5.21 <sup>abc</sup>	2.76 <sup>ab</sup>	2.80 <sup>cd</sup>	2.79 <sup>ab</sup>	2.70	3.00 <sup>de</sup>	2.85 <sup>bc</sup>	7.53 <sup>cde</sup>	7.72 <sup>abcd</sup>	7.63 <sup>abcd</sup>
D <sub>2</sub> F <sub>3</sub>	4.92 <sup>abcd</sup>	5.37 <sup>bc</sup>	5.11 <sup>cdef</sup>	2.65 <sup>abc</sup>	2.70 <sup>cd</sup>	2.66 <sup>c</sup>	2.57	2.30 <sup>ef</sup>	2.43 <sup>cd</sup>	7.30 <sup>de</sup>	7.53 <sup>bcd</sup>	7.43 <sup>bcd</sup>
D <sub>2</sub> F <sub>4</sub>	3.83 <sup>e</sup>	4.62 <sup>d</sup>	4.23 <sup>f</sup>	2.35 <sup>abc</sup>	2.50 <sup>bc</sup>	2.43 <sup>c</sup>	2.20	2.20 <sup>f</sup>	2.19 <sup>cd</sup>	6.88 <sup>e</sup>	6.93 <sup>g</sup>	6.91 <sup>g</sup>
D <sub>3</sub> F <sub>1</sub>	5.67 <sup>a</sup>	5.89 <sup>a</sup>	5.76 <sup>a</sup>	2.87 <sup>a</sup>	3.20 <sup>a</sup>	3.04 <sup>a</sup>	3.07	3.80 <sup>ab</sup>	3.43 <sup>a</sup>	8.63 <sup>a</sup>	8.94 <sup>a</sup>	8.79 <sup>a</sup>
D <sub>3</sub> F <sub>2</sub>	5.43 <sup>ab</sup>	5.46 <sup>ab</sup>	5.44 <sup>abcd</sup>	2.83 <sup>a</sup>	3.07 <sup>a</sup>	2.94 <sup>a</sup>	2.70	3.10 <sup>bcd</sup>	2.90 <sup>ab</sup>	7.95 <sup>abc</sup>	8.00 <sup>abc</sup>	7.98 <sup>abc</sup>
D <sub>3</sub> F <sub>3</sub>	4.53 <sup>bcd</sup>	5.34 <sup>abc</sup>	4.93 <sup>bcd</sup>	2.71 <sup>a</sup>	2.87 <sup>cd</sup>	2.80 <sup>ab</sup>	2.40	2.90 <sup>de</sup>	2.65 <sup>cd</sup>	7.46 <sup>cde</sup>	7.50 <sup>def</sup>	7.48 <sup>def</sup>
D <sub>3</sub> F <sub>4</sub>	3.93 <sup>de</sup>	4.56 <sup>d</sup>	4.27 <sup>def</sup>	2.65 <sup>abc</sup>	2.70 <sup>cd</sup>	2.68 <sup>ab</sup>	2.33	2.90 <sup>de</sup>	2.62 <sup>d</sup>	7.15 <sup>de</sup>	7.23 <sup>fg</sup>	7.19 <sup>fg</sup>
<b>S.Em±</b>	<b>0.12</b>	<b>0.38</b>	<b>0.13</b>	<b>0.04</b>	<b>0.05</b>	<b>0.03</b>	<b>0.20</b>	<b>0.10</b>	<b>0.07</b>	<b>0.06</b>	<b>0.16</b>	<b>0.08</b>
<b>C.D. @ 5%</b>	<b>0.36</b>	<b>1.13</b>	<b>0.38</b>	<b>0.12</b>	<b>0.14</b>	<b>0.08</b>	<b>NS</b>	<b>0.30</b>	<b>0.22</b>	<b>0.19</b>	<b>0.48</b>	<b>0.23</b>
INM	6.91	7.21	7.06	3.30	3.42	3.36	2.90	3.10	3.00	7.47	8.17	7.87
RPP	6.81	7.10	6.95	3.25	3.39	3.32	2.30	2.80	2.55	7.00	7.80	7.40
<b>S.Em±</b>	<b>0.12</b>	<b>0.37</b>	<b>0.14</b>	<b>0.05</b>	<b>0.04</b>	<b>0.03</b>	<b>0.20</b>	<b>0.10</b>	<b>0.07</b>	<b>0.06</b>	<b>0.15</b>	<b>0.07</b>
<b>C.D. @ 5%</b>	<b>0.34</b>	<b>1.06</b>	<b>0.41</b>	<b>0.13</b>	<b>0.13</b>	<b>0.09</b>	<b>0.59</b>	<b>0.30</b>	<b>0.21</b>	<b>0.17</b>	<b>0.45</b>	<b>0.21</b>

Note: INM: Integrated nutrient management (50% RDN through FYM (9 t/ha) + 50% RDN through inorganic fertilizer (45 kg N/ha) + 120 kg P/ha + K @ 60 Kg/ha + 20 t FYM/ha + Azospirillum @ 2.5 kg/ha + PSB @ 2.5 kg/ha RPP: Recommended package of practice (NPK @ 90:120:60 kg/ha +20 t FYM/ha)

jeevamrutha applied @ 1500 l/ha at an interval of 7 days) i.e., 5.76 cm and 3.04 g, respectively, produced significantly maximum flower diameter and individual flower weight respectively than other levels and frequencies of liquid jeevamrutha application. This might be due to direct response of crop to organic manures which may promote cell proliferation efficiently. Cell division and cell enlargement are accelerated by ample supply of nitrogen which initiates meristematic activity in crop. Similar results were obtained by Chaupoo and Kumar [18] in marigold.

When INM and RPP were compared with other interaction treatments, it was found that significantly maximum flower diameter and individual flower weight was recorded in INM (7.06 cm and 3.36 g, respectively) and RPP (6.95 cm and 3.32 g, respectively) than rest of the interaction treatments (Table 4). In INM Inoculation of *Azospirillum* and PSB, enhanced the cell division and enlargement and also produced growth hormones, which is possible reason for increased floral characters. The beneficial effect of nitrogen and phosphorus nutrient on flower size and weight of flower might be due to the fact that nitrogen increases the protein synthesis, thus promote the development of floral primordia, while phosphorus found involved in formation of floral premodial resulting increased flower diameter and weight. The results were in conformity with Chaitra and Patil [19] in China aster.

Shelf life and vase life of flower is an important parameter which decides the durability and marketability of flowers. In the present investigation jeevamrutha application @ 1500 liter per hectare once in 7 days led to higher shelf life (3.43 days) and vase life (8.79 days) of flower than other levels and frequencies of application in pooled data (Table 4). The higher shelf life and vase life in flowers might be due to application of organic manures which influenced flower longevity due to the increased nutrient uptake by plant and greater development of water conducting tissues. Similar results were reported by Sendhilnathan *et al.* [20] in marigold.

#### 4. CONCLUSION

From the present study it can be concluded that, among the jeevamrutha treatments, application of liquid jeevamrutha @ 1500 litre per hectare once in 7 days registered significantly the highest vegetative growth, floral characters and shelf life

with more flower yield per hectare. However, INM and RPP registered significantly highest vegetative growth with more flower yield per hectare than all the natural farming treatments. Hence, for cultivation of China aster, soil application of liquid jeevamrutha @ 1500 litre per hectare once in 7 days was found better among all the natural farming treatments.

#### ACKNOWLEDGEMENT

Authors are thankful to Department of Floriculture and Landscape Architecture, K. R. C. College of Horticulture, Arabhavi and University of Horticultural Sciences, Bagalkot

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Anonymous, Area and production of horticulture crops (2020-21), Department of Horticulture, Lal bagh, Bangaluru. 2022;61-72.
2. Yogananda SB, Devakumar N, Shruti MK, and Ningaraju, Growth and yield of cowpea as influenced by different sources of organic manures. Natl. Symp. Organic Agric. Sustain. Food Security: Challenges and Opportunities. Tamil Nadu, India, 2015;113-120.
3. Biradar N, Effect of organic liquid formulations on growth and yield of organic French bean (*Phaseolus vulgaris* L.). M.Sc. Thesis, 2016; Univ. Agric. Sci., Bengaluru (India).
4. Palekar S, Text book on Shoonyabandovalada naisargika krushi, published by Swamy Anand, Agri Prakashana, Bangalore; 2006.
5. Neelima SG, Sreenivas MN, Influence of liquid organic manures on growth, nutrient content and yield of tomato (*Lycopersicon esculentum* L.) in the sterilized soil. Karnataka J. Agric. Sci. 2011;24(2):153-157.
6. Kumbar B, Devakumar N, Effect of jeevamrutha and panchagavya on growth, yield and microbial population of French bean (*Phaseolus vulgaris* L.). Adv. Life Sci. 2016;5(9):3619-3623.
7. Gangadhar K, Devakumar N, Vishwajith, Lavanya G, Growth, yield and quality



- parameters of chilli (*Capsicum annuum* L.) as influenced by application of different organic manures and decomposers. Int. J. Chem. Stud. 2020;8(1):473-482.
8. Nalina L, Kumar N, Soorianathasundram K, Jeyakumar P, Effect of different nutrient levels on growth and development of tissue cultured banana cv. Robusta (AAA). Indian J. Hort. 2009;66(2):169-174.
  9. Singh M, Dwivedi DH, Kumar M, Efficiency of organic and biodynamic manures on growth and flowering in marigold (*Tagetes patula* L.). Prog. Agric. 2015;15(1): 134-137.
  10. Siddappa, Effect of liquid organic manures on growth and yield of field bean. M. Sc. (Agri.) Thesis, 2015, Univ. Agric. Sci., Bangalore (India).
  11. Kasbe SS, Mukund J, Bhaskar S, Characterization of farmers jeevamrutha formulations with respect to aerobic rice. Mysore J. Agric. Sci., 2009;43(3):570-573.
  12. Fazeel MLM, Sutharsan S, Srikrishnah S, Effect of different levels of nitrogen fertilizer and jeevamrita application on growth and yield of *Abelmoschus esculentus*. Int. Symp. Promoting Multidisciplinary Academic Res. and Innovation., 27-28 November, 2019, South Eastern University of Sri Lanka. 2019;310-319.
  13. Jhade RK, Ambulkar PL, Shrivastava DC, Alawa SL, Pannase SK, Efficacy of INM with jeevamrutha on growth, yield and quality of papaya (*Carica papaya* L.) cv. Taiwan Red Lady under satpura plateau region of Madhya Pradesh district Chhindwara. Int. J. Curr. Microbiol. App. Sci. 2020;1:3050-3056.
  14. Chaithra M, Sujith G, Influence of farmyard manure and jivamrutha on growth and yield of sunflower. Int. J. Environ. 2021;11(11):135-141.
  15. Kumar SB, Prasanth P, Sreenivas M, Gouthami P, Sathish G, Effect of NPK, biofertilizers and biostimulants on flowering, flower quality and yield of China aster Cv. 'Arka Kamini'. The Pharma Innov. J. 2022;11(5):1239-1243.
  16. Mahapatra SS, Sunita N, Reddiramu Y, Prasanthi Y, Efficacy of different organic nutrient management practices on growth and yield of finger millet. Andhra Pradesh J. Agric. Sci. 2017;3(3):165-170.
  17. Gorabal DA, Effect of different levels of jeevamrutha and ghanajeevamrutha on yield and quality of rainfed groundnut. M.Sc. Thesis, Univ. Agric. Hort. Sci. Shivamogga (India); 2020.
  18. Chaupoo AS, Kumar S, Integrated nutrient management in marigold (*Tagetes erecta* L.) cv. Pusa Narangi Gaiinda. Int. J. Curr Microbiol. Appl. Sci. 2020;9(5): 2927-2939.
  19. Chaitra R, Patil VS, Integrated nutrient management studies in China aster (*Callistephus chinensis* (L.) Nees). Karnataka J. Agric. Sci. 2007;20 (3):689-690.
  20. Sendhilnathan R, Velmurugan V, Manimaran P, Effect of bio salicylic acid in increasing yield of marigold (*Tagetes erecta* L.). Adva. Plant Sci. 2017;(22): 119-121.

© 2023 Hegde et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/97136>