



Association between the ABO Type, Rh Factor and the 2D:4D Ratio: A Cross-Sectional Study in Ghana

Moses Banyeh ^{a*}, Martha Nyewie ^b, Abdul-Rafik Abdulai ^c, Charles Nkansah ^d, Gabriel Abbam ^d, Thea Kangkpi ^e, Kervin Edinam Zogli ^f, Shafiat Omotoyosi Shittu ^g, Augusta S. Kolekang ^h, Muniru Mohammed Tanko ^f, Peter Eugene Atuahene ^a, Aisha Yaaba Iddrisu ^a, Christine Ekor ^a and Emmanuel Osei Akoto ^a

^a Department of Biomedical Laboratory Science, University for Development Studies, Tamale, Ghana.

^b Department of Public Health, Community Nurses Training School, Ministry of Health, Tamale, Ghana.

^c Department of Histology and Histotechnology, University for Development Studies, Tamale, Ghana.

^d Department of Haematology, University for Development Studies, Tamale, Ghana.

^e Department of Infectious Diseases, University for Development Studies, Tamale, Ghana.

^f Department of Immunology and Immunodiagnostics, University for Development Studies, Tamale, Ghana.

^g Department of Population and Reproductive Health, University for Development Studies, Tamale, Ghana.

^h Department of Epidemiology, Biostatistics and Disease Control, University for Development Studies, Tamale, Ghana.

Authors' contributions

This work was carried out in collaboration among all authors. Authors MB, MN, ARA, CN, GA, TK, and ASK contributed to the conception, methodology, supervision, and writing-review of the manuscript. Authors KEZ, SOS, MMT, PEA, AYI, CE, and EOA contributed to experimentation, data collection, and writing-review of the manuscript. Authors MB and CN contributed to data analysis, statistical interpretation, and writing-draft of the manuscript. All authors read the work and approved its content.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/120945>

*Corresponding author: E-mail: mosesbanyeh@gmail.com;

Cite as: Banyeh, Moses, Martha Nyewie, Abdul-Rafik Abdulai, Charles Nkansah, Gabriel Abbam, Thea Kangkpi, Kervin Edinam Zogli, Shafiat Omotoyosi Shittu, Augusta S. Kolekang, Muniru Mohammed Tanko, Peter Eugene Atuahene, Aisha Yaaba Iddrisu, Christine Ekor, and Emmanuel Osei Akoto. 2024. "Association Between the ABO Type, Rh Factor and the 2D:4D Ratio: A Cross-Sectional Study in Ghana". *Asian Hematology Research Journal* 7 (3):115-25. <https://www.journalahrj.com/index.php/AHRJ/article/view/176>.

ABSTRACT

Aims: The second-to-fourth digit ratio (2D:4D) is the putative marker of prenatal androgen exposure. The 2D:4D ratio tends to be lower in males than females; however, there are population variabilities. Previous studies have suggested a possible association between the ABO and Rhesus blood group systems with the 2D:4D ratio. This observation has not been tested in a Ghanaian population.

Study Design: The study was cross-sectional.

Place and Duration of Study: The study was conducted in Tamale between May and June 2021.

Methodology: There were 206 participants comprising males (n=94) and females (n=112), aged 18 to 32 years. The right digit ratio (2D:4DR), the left digit ratio (2D:4DL) and the right-left difference in digit ratio (Dr-l) were measured using computer-assisted analysis. The ABO blood type and the Rh factor were determined using monoclonal antibodies. The effect sizes were reported as either Hedge's *g* for a *t*-test or partial Eta-squared (η_p^2) for the one-way ANOVA test.

Results: Males' 2D:4DL was lower than females with a small effect size ($P= 0.033$, $g: 0.28$). The differences in digit ratios by the ABO blood type had small to medium effect sizes in both males and females ($P \geq 0.050$, $\eta_p^2: 0.00-.08$). In females, the 2D:4DR of participants with Rh- was higher while the 2D:4DL and the Dr-l were lower than participants with the Rh+ with medium to large effect sizes ($P < 0.050$, $g: 0.56, 0.84, 2.72$ respectively). In males, the 2D:4DL and the Dr-l were higher in participants with Rh+ than the Rh- participants with large effect sizes ($P < 0.050$, $g: 1.00, 2.52$ respectively).

Conclusion: The expression of the Rhesus factor may affect the expression of the 2D:4D ratio in the study population.

Keywords: 2D:4D ratio; prenatal androgens; ABO blood type; Rh factor; Ghana.

ABBREVIATIONS

2D:4D	: Second-to-fourth Digit Ratio
AR	: Androgen Receptor
BMI	: Body Mass Index
CI	: Confidence Interval
EFNA1	: Ephrin A1
FLI1	: Friend leukaemia integration 1 Transcription Factor
GLI3	: Zinc Finger Protein
GLIS1	: Glis Family Zinc Finger 1
HOX	: Homeo Box
L	: Left
LDAH	: Lipid Droplet-associated Hydrolase
LIN28B	: Lin-28 Homolog B
M	: Mean
OLA1	: Olg-like ATPase 1
R	: Right
SALL1	: Spalt-like 1
UDS	: University for Development Studies

1. INTRODUCTION

The ratio of the length of the fore-finger to the size of the ring-finger is termed the 2D:4D ratio.

It has been suggested that the 2D:4D ratio is a putative marker of prenatal androgen exposure. The ratio is sexually dimorphic with males showing lower ratios than females [1,2]. Previous twin and familial studies have demonstrated that the 2D:4D ratio is heritable due to the observation of significant additive genetic effects [3,4]. Also, the 2D:4D ratio is influenced by assortative mating [5]. The heritability of the 2D:4D ratio may be complex as several genes may be involved. However, the Homeobox and the androgen receptor genes have been suggested to play a role in the heritability of the 2D:4D ratio. Recent advances in molecular biology have also identified novel gene loci that may be associated with the 2D:4D ratio [6].

The ABO and the Rh are the two major blood group systems from the over 30 known blood group systems in humans [7]. The gene that controls the expression of the ABO system is located on chromosome 9 with locus at q34.2 while the genes for Rh are located at p36.11 on the short arm of chromosome 1 [8]. The ABO

system has four main phenotypes; A, B, AB and O while the Rh system has several phenotypes but the RhD is the most immunogenic [9]. Some genes that are associated with the 2D:4D ratio may be located around the vicinity of the ABO and Rh loci and may be inherited together or there may be some level of pleiotropism between these genes [8,6].

Previous studies have attempted to draw an association between the 2D:4D ratio and many human phenotypes [10,11]. However, the work of Voracek [8] has been the single most significant study that sought to determine an association between the 2D:4D ratio and human blood group systems. This study sought to determine the association between the ABO system, the Rh factor and the 2D:4D ratio in a Ghanaian population because there are variabilities in the 2D:4D ratio and the distribution of human blood group systems within and between populations [9,12].

2. MATERIALS AND METHODS

2.1 Study Design and Settings

The study was cross-sectional from May to June 2021 at the University for Development Studies (UDS), Tamale. UDS is a multicampus, multidiscipline and multinational university that is located in the Northern region of Ghana. The Tamale campus is the largest and a host to academic programs in the fields of Medical Laboratory Sciences, Nursing, Nutritional Sciences, Medical Imaging, Pharmaceutical Sciences, Medicine and Education [13].

2.2 Participants

The study involved 206 adults (males: 94 and females: 112) within the age range of 18 to 32 years. The target study population was stratified by sex before a nonprobability sampling technique was used to select the study participants. The participants were devoid of fractures or deformities to their fingers or upper extremities. Participation in the study was voluntary and no restrictions were placed on a participant's cultural group, the program of study or religious affiliation.

2.3 Variables

The 2D:4D ratios and their derivatives (M2D:4D and Dr-l) were the independent variables. The ABO type and Rh factor of the participants were the dependent variables. The participants' age, cultural group and body mass index (BMI) were the additional variables.

2.4 Data collection

2.4.1 Digit measurements

The digit or finger lengths of all the participants were measured using a computer-assisted program. The participant was asked to remove all objects such as rings from their fingers before the palmar surface of each hand was placed on the flatbed surface of a Hp 2620 series Desk jet scanner (HP Inc. CA 94304 United States). The participant's second to fifth fingers were held parallel and the tip of the middle finger aligned with the wrist and elbow [14]. The palm and fingers were then scanned, together with the participant's study unique identifier, at a resolution of 150 dpi. The scanned images were then exported to GIMP (v 2.10.22), an image manipulation program (www.gimp.org). Finger lengths were measured from the mid-point of the most proximal flexion crease to the tip of each finger using a mouse-assisted calliper. Measurements were taken twice by one observer at a week's interval. The intraclass correlation coefficients (two-way mixed, single measures with absolute agreement) were found to be 0.98 and 0.97 respectively for the right (2D:4DR) and the left (2D:4DL) digit ratios. The two measurements were then averaged to obtain the final value. The mean of the left and right 2D:4D ratio was calculated (M2D:4D) and also the right-left difference or directional asymmetry (Dr-l).

2.4.2 Blood sample collection and typing

Venous blood was collected into an EDTA anticoagulant tube. The tubes were gently inverted about eight times to ensure proper mixing of the blood with the anticoagulant. The blood samples were centrifuged at 1500 rpm for five minutes to obtain the red cells [15]. The red cells were washed three times with saline before typing by agglutination using anti-A, anti-B and anti-D monoclonal antibodies (Immucor Inc., Norcross, GA, USA).

2.5 Statistical Analysis

The data were collected and then entered into an Excel spreadsheet (RRID: SCR_016137) before statistical analysis in SPSS (v23) (RRID: SCR_019096) and GraphPad Prism (v8) (RRID: SCR_002798). The continuous data were checked for outliers and normality using the ROUT and Kolmogorov-Smirnov tests, respectively. Descriptive statistics were performed and parametric variables were presented as mean \pm SD and frequency (%) for

categorical variables. The differences between male and female variables were determined using the student t-test (unpaired, 2-tailed). For the categorical variables, the sex variable was dummy coded (female=zero, male=one) before logistic regression analysis. The ABO type was dummy coded (O=zero, A=one, B=two and AB=three) and also the Rh factor (dummy coded Rh+ =0 and Rh- =1). The differences in means of the digit ratios by the participant ABO blood types were compared using the one-way ANOVA test followed by the Post hoc test (Bonferroni). The mean digit ratios by participants' Rh factor were compared using the student t-test (unpaired, 2-tailed). The Hedge's *g* method was used to estimate the effect sizes for the t-test due to the differences in male and female sample sizes while the partial Eta-squared (η_p^2) was used to estimate the effect sizes in the one-way ANOVA analysis [16].

To determine the relationship between digit ratios and the distribution of the ABO type and Rh factor among the participants, digit ratios \leq mean were regarded as low (dummy coded as zero) and digit ratios $>$ mean were regarded as high (dummy coded as one). Participants with a right-left difference in digit ratio ($Dr-l$) $<$ 0.00 were regarded as having a more leftward asymmetry (dummy coded as zero) while those with $Dr-l \geq 0.00$ were considered as having a

more rightward asymmetry (dummy coded as one). Logistic regression analysis was then performed to establish associations. The ABO type, the Rh factor and the cultural group variables were entered simultaneously into the same model for all logistic regression analyses because of possible pleiotropic effects and cultural variabilities. The results were then reported as adjusted odds ratios (AOR) with their 95% confidence intervals (CI). All analyses were 2-tailed and a probability value $<$ 0.050, was considered statistically significant.

3. RESULTS

3.1 The General Characteristics of the Study Population

The general characteristics of the study population are summarized in Table 1. The study involved 206 participants with females constituting 54.4% (112/206) and males, 45.6% (94/206). The study participants were aged between 18 to 32 years with a mean \pm SD age of 22.6 \pm 2.61 years. The ABO type O was the most common (48.4%) and the AB type was the least common (4.4%). Also, the Rh+ formed the majority (89.4%) while the Rh- constituted the remainder (10.4%).

Table 1. General characteristics of the study population

Variable	Statistic
Age(years)	22.6 \pm 2.61
Sex	
Female	112(54.4)
Male	94(45.6)
Cultural group	
Mole-Dagomba	61(29.7)
Akan	63(30.4)
Others	82(39.9)
ABO type	
O	100(48.4)
A	43(20.9)
B	55(26.4)
AB	8(4.4)
Rh type	
Rh+	185(89.6)
Rh-	21(10.4)
BMI (Kg/m ²)	22.0 \pm 2.95
2D:4DR	0.94 \pm 0.034
2D:4DL	0.94 \pm 0.036
M2D:4D	0.94 \pm 0.033
Dr-l	-0.001 \pm 0.027

Note. The results were presented as mean \pm SD for continuous variables and frequency (%) for categorical variables

Table 2. Comparison of male and female blood types and anthropometric variables

Variable	Female	Male	AOR (95%CI)/P-value	<i>g</i>
ABO type				
O	57(57.0)	43(43.0)	1	
A	21(48.7)	22(51.3)	1.400(0.646-3.035)	
B	34(61.5)	21(38.5)	0.823(0.374-1.815)	
AB	4(50.0)	4(50.0)	1.367(0.316-5.919)	
Rh type				
Rh+	103(55.5)	82(44.5)	1	
Rh-	12(58.8)	9(41.2)	0.830(0.295-2.334)	
2D:4DR	0.94±0.035	0.93±0.033	0.103	0.293
2D:4DL	0.94±0.036	0.93±0.034	0.033	0.285
M2D:4D	0.94±0.034	0.94±0.031	0.035	0.000
Dr-l	-0.003±0.026	-0.000±0.027	0.460	0.113

Table 3. Comparison of digit ratios by ABO blood type in males and females

Variable	O	A	B	AB	P-value	η_p^2
Female						
2D:4DR	0.94±0.035	0.94±0.032	0.95±0.035	0.98±0.034	0.062	0.081
2D:4DL	0.94±0.037	0.95±0.035	0.95±0.038	0.98±0.037	0.262	0.045
M2D:4D	0.94±0.035	0.94±0.030	0.95±0.033	0.98±0.033	0.110	0.067
Dr-l	-0.004±0.024	0.009±0.030	0.003±0.029	0.003±0.010	0.538	0.025
Male						
2D:4DR	0.94±0.029	0.94±0.032	0.92±0.026	0.94±0.084	0.384	0.044
2D:4DL	0.94±0.033	0.94±0.030	0.92±0.033	0.94±0.028	0.155	0.074
M2D:4D	0.94±0.029	0.94±0.030	0.92±0.026	0.95±0.056	0.216	0.063
Dr-l	-0.001±0.024	-0.002±0.018	0.003±0.032	-0.003±0.062	0.959	0.004

Note. Results were presented as mean ± SD. The one-way ANOVA test with Post hoc (Bonferroni) analysis was performed to determine the differences between the means. The effect sizes were presented as partial Eta-squared (η_p^2): small ($0.01 < \eta_p^2 \leq 0.06$), moderate ($0.06 < \eta_p^2 \leq 0.14$), large ($\eta_p^2 > 0.14$)

Table 4. Comparison of digit ratios by Rh blood type in males and females

Variable	Rh+	Rh-	P-value	<i>g</i>
Female				
2D:4DR	0.93±0.036	0.95±0.033	0.010	0.560
2D:4DL	0.96±0.036	0.93±0.033	<0.001	0.840
M2D:4D	0.95±0.036	0.94±0.032	0.502	0.281
Dr-l	-0.025±0.015	0.016±0.016	<0.001	2.715
Male				
2D:4DR	0.92±0.029	0.94±0.034	0.250	0.678
2D:4DL	0.95±0.030	0.92±0.031	<0.001	0.997
M2D:4D	0.94±0.029	0.93±0.032	0.238	0.342
Dr-l	-0.026±0.017	0.017±0.018	<0.001	2.516

Note. Results were presented as mean ± SD. The differences between the means were determined using the student t-test (unpaired, 2-tailed). The effect sizes were presented as Hedge's *g*: similar ($g < 0.20$), small ($0.20 \leq g < 0.50$), moderate ($0.50 \leq g < 0.80$), large ($g \geq 0.80$)

3.2 Comparison of Male and Female Variables

From Table 2, the mean 2D:4DL of males was significantly lower than females ($P=0.033$), but the effect size was small ($g=0.28$). However, there were no significant differences in the distribution of the ABO type and Rh factor between the male and female participants.

3.3 Differences in Digit Ratio by the ABO Type

The digit ratios, their mean and the difference were stratified by the ABO type in both females and males (Table 3). Although there were no significant differences in digit ratios by the ABO blood type, individuals with the AB type had the highest left, right and mean 2D:4D ratio while

Table 5. The association between the 2D:4D ratio and the distribution of the ABO and Rh blood types

Variable	Female				Male			
	≤Mean	>Mean	AOR (95%CI)	P-value	≤Mean	>Mean	AOR (95%CI)	P-value
2D:4DR								
O	37(59.1)	20(40.9)	1		21(48.5)	22(51.5)	1	
A	11(52.6)	10(47.4)	1.295(0.438-3.829)	0.641	9(40.0)	13(60.0)	1.472(0.473-4.577)	0.504
B	14(41.7)	20(58.3)	2.037(0.738-5.619)	0.170	15(73.3)	6(26.7)	0.351(0.092-1.337)	0.125
AB	1(25.0)	3(75.0)	4.279(0.409-44.787)	0.225	3(75.0)	1(25.0)	0.286(0.026-3.155)	0.307
Rh+	53(51.9)	50(48.1)	1		44(53.8)	38(46.2)	1	
Rh-	6(50.0)	6(50.0)	1.101(0.283-4.284)	0.890	4(42.9)	5(57.1)	1.786(0.337-9.454)	0.495
2D:4DL								
O	31(54.5)	26(45.5)	1		16(36.4)	27(63.6)	1	
A	11(52.6)	10(47.4)	1.080(0.367-3.180)	0.889	8(35.0)	14(65.0)	1.158(0.357-3.757)	0.807
B	16(45.8)	18(54.2)	1.419(0.520-3.869)	0.494	11(53.3)	10(46.7)	0.525(0.149-1.849)	0.316
AB	1(25.0)	3(75.0)	3.597(0.344-37.593)	0.285	2(50.0)	2(50.0)	0.456(0.050-4.158)	0.486
Rh+	52(50.6)	51(49.4)	1		35(43.1)	47(56.9)	1	
Rh-	6(50.0)	6(50.0)	1.007(0.261-3.885)	0.992	1(14.3)	8(85.7)	4.966(0.537-45.933)	0.158
M2D:4D								
O	30(52.3)	27(47.7)	1		21(48.5)	22(51.5)	1	
A	12(57.9)	9(42.1)	0.798(0.269-2.367)	0.684	9(40.0)	13(60.0)	1.604(0.507-5.078)	0.507
B	11(33.3)	23(66.7)	2.185(0.773-6.177)	0.140	14(66.7)	7(33.3)	0.496(0.133-1.847)	0.133
AB	1(25.0)	3(75.0)	3.3(0.316-34.499)	0.319	2(50.0)	2(50.0)	0.731(0.077-6.925)	0.077
Rh+	48(46.9)	55(53.1)	1		44(53.8)	38(46.2)	1	
Rh-	6(50.0)	6(50.0)	0.969(0.248-3.777)	0.963	1(14.3)	8(85.7)	7.725(0.844-70.670)	0.844

Note. Results were presented as frequency (%). Male and female blood types were compared by lower (\leq mean) and higher ($>$ mean) digit ratios. Both the ABO and the RhD blood types were entered simultaneously into the same model for each digit ratio and the adjusted odds ratio (AOR) reported along with their 95% confidence interval (CI)

Table 6. The association between digit ratio asymmetry and the blood group type

Variable	<0.00	≥0.00	AOR (95%CI)	P-value
Female				
O	26(45.5)	31(54.5)	1	0.603
A	12(57.9)	9(42.1)	0.620(0.208-1.850)	0.391
B	14(41.7)	20(58.3)	1.115(0.405-3.070)	0.834
AB	1(25.0)	3(75.0)	2.779(0.259-29.843)	0.399
Rh+	46(44.4)	57(55.6)	1	
Rh-	7(60.0)	5(40.0)	0.527(0.132-2.106)	0.365
Male				
O	18(42.4)	25(57.6)	1	
A	8(35.0)	14(65.0)	1.312(0.412-4.177)	0.646
B	7(33.3)	14(66.7)	1.428(0.396-5.153)	0.586
AB	3(75.0)	1(25.0)	0.260(0.024-2.818)	0.268
Rh+	32(38.5)	50(61.5)	1	
Rh-	5(57.1)	4(42.9)	0.549(0.108-2.799)	0.470

Note. Results are presented as frequency (%). The ABO and Rh blood types were entered in the same logistic regression model to obtain the adjusted odds ratios (AOR) and confidence intervals (CI).

individuals with type A had the highest asymmetry (Dr-I) among females. However, in males, participants with the ABO type B had the lowest 2D:4D ratio while those with type AB had the highest M2D:4D. The ABO type O had the lowest asymmetry (Dr-I) among males. The effect sizes were small to medium (η_p^2 : 0.00 - 0.08)

3.4 Differences in Digit Ratio by the Rh Factor

Digit ratios were compared by the Rh factor in both males and females (Table 4). Among the females, the 2D:4DR was significantly lower while the 2D:4L was significantly higher in those with Rh+, relative to the Rh-, with medium to large effect size ($P < 0.050$, g : 0.56 – 2.72). Also, the males with Rh+ had significantly higher 2D:4DL as compared to their counterparts with the Rh- with a large effect size ($P < 0.050$, g : 1.00 – 2.52).

3.5 The Association between the 2D:4D Ratio and Blood Type

To determine the association between prenatal androgen exposure (as indexed by the 2D:4D ratio) and the distribution of the ABO type and Rh factor, the participants' blood types were compared by low (\leq mean) and high ($>$ mean) digit ratios in both males and females (Table 5). Both the ABO type and the Rh factor were entered in the same logistic regression model. No significant associations between the 2D:4D ratios and the ABO type or the Rh factor were observed.

3.6 The Association between Digit Ratio Asymmetry and Blood Type

The ABO and Rh blood types in both males and females were compared by more leftward ($Dr-I < 0.00$) and more rightward ($Dr-I \geq 0.00$) asymmetry (Table 6). No significant associations were observed between the Dr-I and the distribution of the ABO type or the Rh factor.

4. DISCUSSION

The study sought to determine the association between prenatal androgen exposure (as indexed by the 2D:4D ratio) and the ABO type or Rh factor. It was observed that the 2D:4DL of males was significantly lower than females. There were no significant differences in digit ratio by the ABO type but the effect sizes were small to medium. However, females with the ABO type AB and type B had the highest digit ratio and asymmetry (Dr-I) respectively. Also, males with ABO type B had the lowest 2D:4D ratio while those with type O had the lowest asymmetry (Dr-I). But females with Rh- had significantly higher 2D:4DR while those with Rh+ had significantly higher 2D:4DL and Dr-I with medium to large effect sizes, when compared to those with Rh-. Moreover, males with Rh+ had significantly higher 2D:4DL and Dr-I as compared to those with Rh- with large effect sizes. But there was no significant association between the 2D:4D ratio and the distribution of the ABO type or the Rh factor in the study population.

Studies regarding the association between the 2D:4D ratio and blood type are rare. As of the time of writing, the work of Voracek [8] was the most significant study on the subject. The study of Voracek [8] involved two separate samples, I and II: Sample I consisted of 200 males and 200 females while sample II consisted of 395 males and 478 females, drawn from the Austrian population. For both samples, males had a lower 2D:4D ratio in both hands but a higher asymmetry (Dr-l) for sample II. Voracek [8] observed that the differences in the 2D:4D ratio by the ABO or Rh factor were small and not significant in both samples.

There have been many suggestions aimed at the possible explanations for these observed differences in digit ratios by ABO and Rh blood systems. There is evidence from familial studies that the 2D:4D ratio is heritable in humans, just like the ABO and the Rh systems [4]. The genes that regulate the expression of the 2D:4D ratios may be located at loci that are in proximity to genes that control the expression of the ABO and Rh blood systems. For the ABO system, this would be on the long arm of chromosome 9, in the vicinity of the gene loci, q34.2 while for the Rh factor, this would be on the short arm of chromosome 1, around position p36.11 gene loci. Another possible explanation is the seeming pleiotropic effects of either the genes controlling the 2D:4D ratio on the genes for the ABO and Rh system or vice versa [8]. The androgen receptor gene (AR), located on the X-chromosome, have been suggested to be associated with the 2D:4D ratio. Polymorphisms in the AR gene (CAG repeats) have been associated with androgen insensitivity and differences in the 2D:4D ratio, although this is not a universal observation [17,18]. However, the AR gene has pleiotropic effects on other genes whose expression or suppression may be associated with the expression of the 2D:4D ratio [19,20].

In one of the largest meta-analytic genetic association studies regarding the 2D:4D ratio, 9 novel and two previously known loci were associated with the 2D:4D ratio expression in humans. In total, the 11 genes accounted for about 3.8% of the variance in the mean 2D:4D ratio [6]. The Glis Family Zinc Finger 1 (GLIS1) gene (locus; 1p32.3) which is located on the same chromosome as the Rh factor (chromosome 1), plays an important role in embryogenesis by controlling gene expression at different stages in the process [21]. The Ephrin

A1 (EFNA1) gene (locus 1q21-q22), also located on chromosome 1, has been shown to encode for a protein that plays a role in the mediation of developmental events [6]. Other significant genes, that are associated with the 2D:4D ratio but are not located on the same chromosome as the ABO or Rh genes, but may have a pleiotropic effect were also identified. The gene for Rh-associated glycoprotein (RhAG), denoted as RHAG, which is vital for the expression of the Rh polypeptide is located on chromosome 6 (6p11-p21) [7]. The Lipid droplet-associated hydrolase (LDAH), a gene that can be found around locus, 2p24.1, may be involved in steroidogenesis as it mobilizes cholesterol which could be converted into testosterone [22]. Testosterone plays a marked role in the expression of the 2D:4D ratio. Individuals diagnosed with congenital adrenal hyperplasia (CAH), due to 21-hydroxylase deficiency, tend to have androgenized or masculinized 2D:4D ratios while those with androgen insensitivity or Klinefelter's syndromes tend to have feminized digit ratios although the effect sizes may be small [23]. The Obg-like ATPase 1 (OLA1) is a gene that has a locus at 2q31.1. Animal models that lacked the OAL1 gene were observed to have delayed development characterized by stunted growth and immature organs [24]. The 7p14.1 locus, which is linked to the Zinc finger protein (GLI3) has been implicated in craniofacial and limb deformities such as polydactyly and brachydactyly [25]. Similarly, the 11q24.1-q24.3 locus, which is intergenic between the Friend leukaemia integration 1 transcription factor (FLI1) and ETS1 gene has also been suggested to play a role in preaxial polydactyly [26]. A signal at the locus, 16q12.1, in the vicinity of the spalt-like1 (SALL1) gene on chromosome 16, and SALL3 (18q23) have been associated with skeletal abnormalities including hand and genital malformations [27]. There was some level of association between HOXD12 and the 2D:4D ratio. HOXD12 belong to the Homeobox family of genes that play a vital role in morphogenesis, including the development of the limbs and genitalia [6]. There are no direct shreds of evidence that these genes are also associated with the ABO or Rh systems, but there could be some possible pleiotropic interactions between these genes controlling both systems or these genes may be involved either directly or remotely, in steroidogenesis during human foetal development [8].

The current study has many strengths; to the best of our knowledge, this study is the first to be

conducted in Ghana regarding digit ratios and the ABO or the Rh blood group systems. This is significant owing to the variabilities in the 2D:4D ratio and the distribution of blood group systems in human populations. Unlike Voracek [8], results of blood type were obtained from laboratory analysis and not from self-reported data which could have accounted for the differences in outcomes between the two studies. In this study, digit ratios were measured by computer-assisted analysis which is more precise compared to measurements from photocopies, physical measurements and printed scanned images [14]. Also, both the ABO and the Rh variables were entered simultaneously into the same regression model to reduce confounding in case there are any pleiotropic effects between genes controlling both systems. Moreover, aside from P-values, effect sizes (Hedge's *g* and partial Eta-squared) were presented to add clarity to the observed differences. However, some limitations exist in the current study: there was neither genetic studies regarding the inheritance of the 2D:4D ratio, ABO or the Rh factor nor participants detailed data regarding their ancestry [8]. Also, the sample size was smaller, relative to a previous study [8], which may not allow for the generalization of the findings. In conclusion, there are significant differences in the 2D:4D ratio between the Rh+ and Rh- participants. We, however, recommend further studies with a larger sample size in which the minor blood group systems will also be studied. Further studies should also include genetic and detailed ancestral studies of the participants.

5. CONCLUSION

The second-to-fourth digit ratio, a marker of prenatal hormone exposure may be associated with blood type, particularly the Rhesus blood type. This, however, requires further studies to validate the observations of the current study.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Authors hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

CONSENT AND ETHICAL APPROVAL

The study followed all guidelines regarding human studies as stipulated in the 1964 Declaration of Helsinki and its later amendments. The study was approved by the

institutional review board of the University for Development Studies, Tamale. All participants gave their written informed consent before the study.

ACKNOWLEDGEMENTS

We wish to acknowledge Christine Ekor and Akoto Osei Emmanuel for assisting in the data collection. We acknowledge all the participants for consenting to the study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Lutchmaya S, Baron-Cohen S, Raggatt P, Knickmeyer R, Manning JT. 2nd to 4th digit ratios, fetal testosterone and estradiol. *Early Human Development*. 2004;77(1-2):23-28.
2. Manning Scutt D, Wilson J, Lewis-Jones DI. The ratio of 2nd to 4th digit length: A predictor of sperm numbers and concentrations of testosterone, luteinizing hormone and oestrogen. *Human Reproduction (Oxford, England)*. 1998;13(11):3000-3004.
3. Gobrogge KL, Breedlove SM, Klump KL. Genetic and environmental influences on 2D:4D finger length ratios: A study of monozygotic and dizygotic male and female twins. *Arch Sex Behav*. 2008;37(1):112-118. DOI: 10.1007/s10508-007-9272-2
4. Richards G, Bellin W, Davies W. Familial digit ratio (2D: 4D) associations in a general population sample from Wales. *Early Human Development*. 2017;112:14-19.
5. Voracek M, Dressler SG, Manning JT. Evidence for assortative mating on digit ratio (2D:4D), a biomarker for prenatal androgen exposure. *J Biosoc Sci*. 2007;39(4):599-612. DOI: 10.1017/s0021932006001647
6. Warrington NM, Shevroja E, Hemani G, Hysi PG, Jiang Y, Auton A, Evans DM. Genome-wide association study identifies nine novel loci for 2D:4D finger ratio, a putative retrospective biomarker of testosterone exposure in utero. *Hum Mol Genet*. 2018;27(11):2025-2038. DOI: 10.1093/hmg/ddy121

7. Flegel WA. Molecular genetics and clinical applications for RH. *Transfusion and Apheresis Science*. 2011;44(1):81-91.
8. Voracek M. Digit ratio (2D:4D), ABO blood type, and the Rhesus factor. *Percept Mot Skills*. 2008;107(3):737-746.
DOI:10.2466/pms.107.3.737-746
9. Doku GN, Agbozo WK, Annor RA, Kisseh GD, Owusu MA. Frequency of ABO/Rhesus (D) blood groupings and ethnic distribution in the Greater-Accra region of Ghana, towards effective blood bank inventory. *Int J Immunogenet*. 2019;46(2):67-73.
DOI: 10.1111/iji.12412
10. Butovskaya ML, Lazebny OE, Vasilyev VA, Dronova DA, Karelin DV, Mabulla AZ, Ryskov AP. Androgen receptor gene polymorphism, aggression, and reproduction in Tanzanian foragers and pastoralists. *PLoS One*. 2015;10(8).
11. Stibbard-Hawkes DNE. No association between 2D:4D ratio and hunting success among hadza hunters. *Hum Nat*. 2020;31(1):22-42.
DOI: 10.1007/s12110-019-09359-z
12. Manning. *Digit ratio: A pointer to fertility, behavior, and health*: Rutgers University Press; 2002.
13. UDS. Profile of the university for development studies; 2020.
Available:<https://uds.edu.gh/about-us/> [Assessed 19-04-2020].
Available:<https://uds.edu.gh/about-us/>
14. Allaway HC, Bloski TG, Pierson RA, Lujan ME. Digit ratios (2D:4D) determined by computer-assisted analysis are more reliable than those using physical measurements, photocopies, and printed scans. *Am J Hum Biol*. 2009;21(3):365-370.
DOI: 10.1002/ajhb.20892
15. Canizalez-Román A, Campos-Romero A, Castro-Sánchez JA, López-Martínez MA, Andrade-Muñoz FJ, Cruz-Zamudio CK, Alcántar-Fernández J. Blood groups distribution and gene diversity of the ABO and Rh (D) Loci in the Mexican population. *Biomed Res Int*, 2018;1925619.
DOI: 10.1155/2018/1925619
16. Fritz CO, Morris PE, Richler JJ. Effect size estimates: current use, calculations, and interpretation. *Journal of experimental psychology: General*. 2012;141(1):2.
17. Butovskaya ML, Vasilyev VA, Lazebny OE, Burkova VN, Kulikov AM, Mabulla A, Ryskov AP. Aggression, digit ratio, and variation in the androgen receptor, serotonin transporter, and dopamine D4 receptor genes in African foragers: the Hadza. *Behavior genetics*. 2012;42(4):647-662.
18. Zhang K, Yang X, Zhang M, Wang C, Fang P, Xue M, Gong P. Revisiting the relationships of 2D:4D with androgen receptor (AR) gene and current testosterone levels: Replication study and meta-analyses. *J Neurosci Res*. 2020;98(2):353-370.
DOI: 10.1002/jnr.24502
19. Aurilio G, Cimadamore A, Mazzucchelli R, Lopez-Beltran A, Verri E, Scarpelli M, Montironi R. Androgen receptor signaling pathway in prostate cancer: From Genetics to Clinical Applications. *Cells*. 2020;9(12).
DOI: 10.3390/cells9122653
20. Li Q, Liu B, Chao HP, Ji Y, Lu Y, Mehmood R, Tang DG. LRIG1 is a pleiotropic androgen receptor-regulated feedback tumor suppressor in prostate cancer. *Nat Commun*. 2019;10(1):5494.
DOI: 10.1038/s41467-019-13532-4
21. Li L, Chen K, Wang T, Wu Y, Xing G, Chen M, Liu X. Glis1 facilitates induction of pluripotency via an epigenome-metabolome-epigenome signalling cascade. *Nat Metab*. 2020;2(9):882-892.
DOI: 10.1038/s42255-020-0267-9
22. Goo YH, Son SH, Kreienberg PB, Paul A. Novel lipid droplet-associated serine hydrolase regulates macrophage cholesterol mobilization. *Arterioscler Thromb Vasc Biol*. 2014;34(2):386-396.
DOI: 10.1161/atvbaha.113.302448
23. Richards G, Browne WV, Aydin E, Constantinescu M, Nave G, Kim MS, Watson SJ. Digit ratio (2D: 4D) and congenital adrenal hyperplasia (CAH): Systematic literature review and meta-analysis. *BioRxiv*; 2020.
24. Ding Z, Liu Y, Rubio V, He J, Minze LJ, Shi ZZ. OLA1, a translational regulator of p21, maintains optimal cell proliferation necessary for developmental progression. *Mol Cell Biol*. 2016;36(20):2568-2582.
DOI: 10.1128/mcb.00137-16
25. Al-Qattan MM, Shamseldin HE, Salih MA, Alkuraya FS. GLI3-related polydactyly: A review. *Clin Genet*. 2017;92(5):457-466.
DOI:10.1111/cge.12952
26. Lettice LA, Williamson I, Wiltshire JH, Peluso S, Devenney PS, Hill AE, Hill RE. Opposing functions of the ETS factor

- family define Shh spatial expression in limb buds and underlie polydactyly. Dev Cell. 2012;22(2):459-467.
DOI: 10.1016/j.devcel.2011.12.010
27. Kohlhase J, Hausmann S, Stojmenovic G, Dixkens C, Bink K, Schulz-Schaeffer W, Engel W. SALL3, a new member of the human spalt-like gene family, maps to 18q23. Genomics. 1999;62(2):216-222.
DOI: 10.1006/geno.1999.6005

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/120945>