



## **Influence of Potassium Levels and Spacing on Growth and Yield of Summer Groundnut (*Arachis hypogaea* L.)**

**Kaithi Adithya Reddy <sup>a\*</sup>, Umesha C. <sup>a#</sup> and Mannepu Venkata Sai Ganesh <sup>a\*</sup>**

<sup>a</sup> Department of Agronomy, Sam Higginbottom University of Agriculture, Technology and Sciences, India.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/IJECC/2022/v12i1030863

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/87094>

**Original Research Article**

**Received 20 March 2022**

**Accepted 30 May 2022**

**Published 01 June 2022**

## **ABSTRACT**

**Background:** Groundnuts are immensely rich in potassium, calcium, phosphorus and B vitamins which offers you with a host of health benefits. Groundnuts are a great blend of healthy fats, protein and fibre that curbs your appetite, lowers the risk of heart disease and regulates blood glucose levels.

**Objectives:** Effects of potassium levels and spacing on growth and yield of summer groundnut.

**Methods:** With the goal of studying the effect of potassium and spacing on growth and yield of groundnut under a Randomized block design with 9 treatments (T1-T9) The experimental results revealed that 120 kg N/ha + 60 kg P/ha produced maximum plant height (49.7 cm) No. of Nodules/Plant (106) plant dry weight (22.9) No of pods (19.3) no of kernals (2.7) pod yield (3.62 ta/ha) Halum yield (4.65 ta/ha).

**Conclusion:** The combination of 40 kg potassium and 30x10 proved to be the most advantageous to farmers, resulting in, plant height (49.7 cm) No. of Nodules/Plant(106) plant dry weight (22.9) No of pods (19.3) no of kernals (2.7) pod yield (3.62 ta/ha) Halum yield (4.65 ta/ha) respectively.

**Keywords:** Potassium; spacing; yield.

<sup>o</sup>Msc Scholar;

<sup>#</sup>Assistant Professor;

<sup>\*</sup>Corresponding author: E-mail: adithyareddy8800@gmail.com;

## 1. INTRODUCTION

Groundnut (*Arachis hypogea* L.) is an herbaceous crop of Leguminosae family. The plant has central stem upright with many branches which are different from prostrate to almost erect based on the variety. Groundnut, also known as “The king of oilseeds”. In India, it is produced mostly as oilseed crop, covers an area of 40-50% and gives a production of 60 to 70%. It is first major oil seed crop among all the other oil seed harvests. In India, it covers an area of 85 lakh hectares and produced 84 lakh tones from southern states (Andhra Pradesh, Tamil Nadu, Karnataka), from western and central parts (Gujarat, Maharashtra, Madhya Pradesh), from northern states (Uttar Pradesh, Rajasthan), north western part of the subcontinent i.e., Punjab and from eastern India (Orissa). However, the maximum yield and area of 84% is covered by southern and western regions [1].

Potassium is a multifunctional versatile nutrient, indispensable for plants. Among the three major nutrients, potassium (K) has a special position as evident by its role in increasing the crop yield by adding tolerance to various biotic and abiotic stresses [2] and [3]. The potassium application improves the kernel size of the groundnut, test weight and shelling percentage. Groundnut crop well for potassium and play role in maintaining balance in enzymatic, stomatal activity (water use), transport of sugars, water and nutrient and synthesis of protein, photosynthesis and starch thus K application increases growth and yield attributes in groundnut [4] Planting density is one of the main factors that play an important role on growth, yield and quality of [5] reported that plant dry matter, dry weight/plant and branching were found to be maximum along with yield attributes (pod/plant, yield/plant and 1000-grain weight).

However, thin spacing lower the production of the crop plants because of the competitiveness of the crop plants for nutrients and moisture [6]. Likewise, the row alignment too influences photosynthetic competence and canopy temperature for a better light interception and greater photosynthetic efficiency, a sustained and uniform orientation as well as crop distribution is needed [7] Proper application of spacing along with suitable nutrients will help in attaining maximum productivity within India. All these points are the basis of the present study which was laid out in Zaid 2021.

## 2. MATERIALS AND METHODS

The experiment was carried out during summer season of 2021 at CRF (Crop Research Farm), Department of Agronomy, Naini Agricultural Institute, SHUATS, Prayagraj (UP). The farm is geographically situated at 25° 24' N latitude and 81° 51' Longitude. The experiment was carried out in Randomized Block Design (RBD) with nine treatments which are replicated thrice. The experiment consist of nine. 20 kg K<sub>2</sub>O/ha + 20 cm x 10 cm, (T1) 20 K<sub>2</sub>O/ha + 30 cm x 10 cm (T2) , 20 K<sub>2</sub>O/ha + 40 cm x 10 cm (T3) , K<sub>2</sub>O/ha + 20 cm x 10 cm, (T4) K<sub>2</sub>O/ha + 30 cm x 10 cm (T5) , K<sub>2</sub>O/ha a + 40 cm x 10 cm (T6), 40 K<sub>2</sub>O/ha + 20 cm x 10 cm, (T7) 40 K<sub>2</sub>O/ha + 30 cm x 10 cm, (T8) 40 K<sub>2</sub>O/ha + 40 cm x 10 cm (T9) The mean (maximum and minimum) temperature was 37.98°C and 24.21°C respectively, mean (maximum and minimum) relative humidity was 82.16 percent and 45.26 percent during the crop growing season. The experimental soil was sandy loam in texture, nearly neutral in soil reaction (pH 7.1), low in organic carbon (0.32) edium in available N (70 Kg/ha), p (12.50 Kg/ha) and k (216.10 Kg/ha). Fertilizers were applied in the form of urea, single super phosphate and murate of potash, respectively. The Shelling was done manually, seeds were winnowed and cleaned and seed weight per net lot was recorded on hectare basis and expressed in kg /ha. The observation regarding yield were recorded after harvesting of crop. The recording with references of Gomez KA, Gomez AA [8].

## 3. RESULTS AND DISCUSSION

### 3.1 Growth Attribute

#### 3.1.1 Plant height

At Harvest recorded that significantly maximum plant height (61.20 cm) was recorded with application of 40 Kg k/ha+ 30 x 10 cm. However, treatment with application of 30 Kg k/ha+ 30 x 10 cm (59.40 cm), and 40 Kg k/ha+ 40 x 10 cm (58.60 cm) were statistically at par with 40 Kg k/ha+ 30 x 10 cm compared to other treatments. Applying of spacing 30 x10 cm give less competition for the plants so nutrient uptake will me more so plant height will be increases similar finding are seen in [6].

#### 3.2 Number of Nodules

At harvest recorded that significantly maximum number of nodules (65.21) recorded with

application of 40 Kg k/ha + 30 x 10 cm which was significantly superior over rest of the treatments. However, treatment with application of 30 Kg k/ha + 30 x 10 cm (62.94) is statically at par with 40 Kg k/ha + 30 x 10 cm as compared to other treatments.

The rhizobium inoculation along with fertilizer application including MOP significantly increased the number nodules and nodule dry weight compared with uninoculated control in mungbean cultivars [2].

### 3.3 Plant Dry Weight

At harvest recorded that significantly maximum plant dry weight (31.93) was observed with application of 40 Kg k/ha+ 30 x 10 cm. However, treatment with application of 30Kg k/ha+ 30 x 10 cm (32.38), 30Kg k/ha+ 40 x 10 cm (32.26) and 40 Kg k/ha+ 40 x 10 cm (32.48) were statistically on par with 40 Kg k/ha + 30 x 10 cm compared to other treatments. By less competition there will good in take of food and so photosynthesis done will and the dry weight of the plant will be increases. [7] by applying potassium ADP and energy in tissues will be increases so it also increases the photosynthetic rate [2].

### 3.4 Yield Attributes

#### 3.4.1 Number of pods/plant

Number of pods/plants recorded maximum was obtained with application of 40Kg k/ha+ 30 x 10 cm (19.30). However, treatment with application of 30Kg k/ha+ 30 x 10 cm (19.00) and 40Kg k/ha+ 40 x 10 cm (18.80) were statistically on par with application of 40Kg k/ha+ 30 x 10 cm compared to other treatments. when there is

good growth by spacing and potassium the no of pods per plant will be increases [2] and [7].

### 3.5 Number of Kernels/Pod

Number of kernels/plants recorded maximum was obtained with application of 40Kg k/ha+ 30 x 10 cm (2.00). However, treatment with application of 30Kg k/ha+ 30 x 10 cm (2.00) and 40Kg k/ha+ 40 x 10 cm (2.00) were statistically on par with application of 40Kg k/ha+ 30 x 10 cm compared to other treatments. Similar finding was recorded in the [7].

### 3.6 Pod Yield (t/ha)

Pod yield, recorded maximum with application of 40Kg k/ha+ 30 x 10 cm (3.62 t/ha) significantly superior over rest of the treatments. However, treatment with application of 30 Kg k/ha+ 30 x 10 cm (3.48 t/ha) were statistically at par with application of 40Kg k/ha+ 30 x 10 cm compared to other treatments. When there free uptake nutrient by the spacing and free energy to new tissue by potassium there will good growth in the pod yield [3] and [7].

### 3.7 Halum Yield (t/ha)

Halum yield, recorded maximum with application of 40Kg k/ha+30 x 10 cm (44.76 t/ha) significantly superior over rest of the treatments. However, treatments with application of 30Kg t/ha+ Rhizobium (43.19 t/ha) were statistically at par compared to other treatment combination.by apply ing potassium and free spacing there will lot of improvement in nutrient uptake, energy in new tissues and photosynthesis so the halum yield will be increases [3].

**Table 1. Influence of potassium levels and spacing on growth parameters**

S. No	Treatments	Plant height (cm)	No. of Nodules/Plant	Plant dry weight (g/plant)
1	K 20 kg/ha + 20 X 10 cm	44.2	95.7	17.53
2	K 20 kg/ha + 30 X 10 cm	44.9	98.8	18.99
3	K 20 kg/ha + 40 X 10 cm	46.2	103.3	20.21
4	K 30 kg/ha + 20 X 10 cm	45.7	101.7	19.44
5	K 30 kg/ha + 30 X 10 cm	46.9	106	22.21
6	K 30 kg/ha + 40 X 10 cm	47.2	103.9	21.21
7	K 40 kg/ha + 20 X 10 cm	46.3	101.9	19.55
8	K 40 kg/ha + 30 X 10 cm	49.7	106.9	22.99
9	K 40 kg/ha + 40 X 10 cm	48.3	104.6	21.55
	F-test	S	S	S
	SE(±)	0.54	0.64	0.28
	CD (P=0.05)	1.61	1.92	0.83

**Table 2. No. of pods/plant, No. of kernels/pod, Seed index (g), seed yield (kg/ha), halum yield (kg/ha), harvest index (%)**

S. No	Treatments	No. Pods/plant	No. of kernels/pod	Seed index (g)	Pod yield (g)	Halum yield (g)	Harvest index (%)
1	K 20 kg/ha + 20 X 10 cm	16.9	1.4	35.18	2.82	3.79	42.60
2	K 20 kg/ha + 30 X 10 cm	17.5	1.5	35.4	2.85	3.93	41.94
3	K 20 kg/ha + 40 X 10 cm	18	1.8	38.5	3.1	4.21	42.77
4	K 30 kg/ha + 20 X 10 cm	17.5	1.5	35.98	3.03	4.04	42.88
5	K 30 kg/ha + 30 X 10 cm	19	2.3	39.95	3.51	4.56	43.28
6	K 30 kg/ha + 40 X 10 cm	18.3	1.8	39.6	3.14	4.39	41.70
7	K 40 kg/ha + 20 X 10 cm	17.9	1.5	36.9	2.99	4.13	42.00
8	K 40 kg/ha + 30 X 10 cm	19.3	2.7	41.5	3.62	4.65	43.72
9	K 40 kg/ha + 40 X 10 cm	18.8	2	39.17	3.3	4.51	43.14
	F-test	S	S	S	S	S	NS
	SE(±)	0.21	0.06	0.58	0.07	0.05	0.82
	CD (P=0.05)	0.64	0.53	7.74	0.22	0.17	-

#### 4. CONCLUSION

The combination of 40 kg potassium and 30x10 proved to be the most advantageous to farmers, resulting in, plant height (49.7 cm) No. of Nodules/Plant (106) plant dry weight (22.9) No of pods (19.3) no of kernals (2.7) pod yield (3.62 ta/ha) Halum yield (4.65 ta/ha) respectively. Hence we can recommend for the farmers.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Abdzad GA, Noorhosseini NSA. Effects of Potassium fertilizers on yield and yield components of peanut (*Arachis hypogaea* L.). American-Eurasian Journal of Agricultural and Environmental Sciences. 2010;9(3):256-262.
2. Anonymous. Estimates of Production of Food grains, Directorate of Economics and Statistics, Department of Agriculture, Cooperation and Farmers Welfare Government of India. New Delhi; 2020-21.
3. Bhagiya SR, Polara JV. Effect of potassium on yield and quality and nutrient absorption by groundnut. Advances in Plant Sciences. 2005;18(11): 803-806.
4. Caliskan S, Ozkaya I, Caliskan ME, Arslan M. The effect of nitrogen and iron fertilizer on growth, yield and fertilizer use efficiency of soybean in Mediterranean-type soil. Field Crop Research. 2008;108(2):126-132.
5. Duyingqiong Q, Liao Xinrong, He Jianghua, Hiang Zhoyao, Zhou Xiaohong. Effect of potassium on the growth, development and yield of peanut. Plant Nutrition. Fertilizer: Sci. 2002;8(2):233-235.
6. Kumar SN. Effect of plant density and weed management practices on production potential of groundnut (*Arachis hypogaea* L.).

- Indian Journal of Agricultural Research. 2009;43(1):57-60.
7. Parameshwarareddy R, Angadi SS, Yenagi BS. Influence of plant spacing and fertilizer levels on growth and yield of summer groundnut. Journal of Pharmacognosy and Phytochemistry. 2019;8(3):3745-3748.
8. Gomez KA, Gomez AA. Statistical Procedures for Agricultural Research, edn. 2, Johns & Sons, New York. 1984; 591.

---

© 2022 Reddy et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/87094>