



Unintended Inhalational Organophosphate Poisoning Presenting as Angioedema: A Rare Case Report

Sanyukta Hepat ^{a*}, Abhijit Wadekar ^{a[≡]}, Gaurav Jagtap ^{a[≡]}, Vijay Kota ^{a[≡]}
and Anil Wanjari ^{a^o}

^a Department of Medicine, Jawaharlal Nehru Medical College, Datta Meghe Institute of Medical Sciences (Deemed to be University), Wardha, Maharashtra, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i52A33574

Editor(s):

(1) Dr. Paola Angelini, University of Perugia, Italy.

Reviewers:

(1) Yasser Abu Safieh, An-Najah University, Palestine.

(2) Rachna Wadhwa, G. B. Pant Hospital, GIPMER, Delhi University, India.

(3) Pallavi Tiwari, Dr. A.P.J. Abdul Kalam Technical University, India.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:
<https://www.sdiarticle5.com/review-history/76851>

Case Study

Received 10 September 2021

Accepted 20 November 2021

Published 29 November 2021

ABSTRACT

Organophosphorus compounds are chemical agents in widespread use throughout the world, mainly in agriculture. In developing countries organophosphorus (OP) poisoning is a commonly encountered problem. Here we present a rare case of OP poisoning with angioedema as the presenting feature. Prompt treatment of these cases will help in reducing the morbidity and mortality.

Keywords: Organophosphate poisoning; angioedema; atropine.

1. INTRODUCTION

In Indian population, especially, the rural areas, one of the major public health concerns is

suicidal poisoning by organophosphorus pesticide, due to its cheap and easy accessibility. Since organophosphorus poisoning is associated with high mortality and morbidity it is of prime

[≡] Post Graduate Resident

^o Professor

*Corresponding author: E-mail: sanyukta211194@gmail.com;

importance to the clinicians [1]. Early diagnosis should be made to initiate prompt treatment in order to save the patient's life [2]. Sometimes the presentation is atypical and onset of symptoms are delayed. In such cases, even when the symptoms are mild in initial stages, still the patients should be observed for longer periods of time [3]. There have been no previous case reports of angioedema seen in organophosphate poisoning patient.

2. CASE REPRESENTATION

A 22-year-old male (weight 56 kg) was admitted in our hospital because of acute organophosphate (OP) poisoning. He had history of inhalational poisoning with monocrotophos and profenophos while spraying insecticide in a farm. He had multiple episodes of vomiting the following day and was brought to the hospital by his mother. He had swelling of both eyes and upper lip since one day. He had difficulty in deglutition suggestive of laryngeal edema and ryle's tube was inserted at the time of admission. At the time of hospitalization, he was conscious and oriented. He had no history of diabetes mellitus, hypertension and psychological problems. On admission, his vital signs revealed a pulse rate of 68/min, blood pressure of 110/70 mmHg, respiratory rate of 18/min, afebrile, and O₂ saturation of 97%. Neurological examination was done and Glasgow Coma Scale was 15/15. On local examination he had edema around both eyelids and swelling of the upper lip (Figure 1). Pupils were miotic on examination. On respiratory examination bilateral coarse crepitations were present. Cardiovascular

examination revealed normal heart sounds. Gastric lavage via nasogastric tube was done in the casualty.

Laboratory investigations revealed his haemoglobin as 14.9g/dL, total leukocyte count 10,400/cumm and platelet count 2,32,000/cumm. Peripheral smear showed predominantly normocytic, normochromic RBC's, PT INR was 1.07, PT was 12.8 (PT control was 11.9) and APTT was 30.6 (APTT control was 29.5). His serum sodium was 143mg/dL, potassium was 4.6 mg/dL. Liver enzymes was normal (Aspartate aminotransferase of 30mg/dL and Alanine aminotransferase of 27mg/dl, total bilirubin 0.9mg/dL, conjugated bilirubin 0.2mg/dl, unconjugated bilirubin 0.7mg/dL, albumin 4.4g/dL, globulin 3.3g/dL). Serum creatinine was 0.6mg/dL, urea was 13mg/dL. Electrocardiogram, and chest X-ray were normal. There was decreased serum cholinesterase in our patient of level 0.2 mg/dL. Fluid therapy was started, and Inj. Atropine 0.6mg was administered intramuscularly in lateral thigh. He did not have hypotension or respiratory distress. In view of angioedema of face IV hydrocortisone 100 mg and chlorpheniramine 10 mg were given. The general condition of the patient improved after management.

Oralcetirizine was also given. To improve the clinical manifestations of OP poisoning, pralidoxime infusion was also started for the patient. After 5 days of hospitalization, he was discharged from the hospital in stable condition.



Fig. 1. Angioedema around eyelids and swelling of upper lip

3. DISCUSSION

In organophosphorus poisoning the symptom onset depends on the route of exposure. The common routes are oral, inhalation, occasionally parenteral and through skin [4]. The onset of symptoms is most rapid in inhalational poisoning occurring over a few minutes. In farmers, while spraying insecticides, the direction of wind may result in accidental inhalation exposure. When poisoning occurs through skin exposure the symptoms onset will depend on few factors like the skin intactness, solubility and amount of organophosphorus compound [5,6]. In ingestional OP poisoning, the time lag period for symptom onset would depend on the amount consumed and the absorption characteristics of the OP compound. Usually the symptoms occur within few minutes to hours. In India OP pesticides are commonly used for suicide purposes particularly in the rural areas [7]. Another important mode of poisoning is occupational exposure. The outcome of OP poisoning depends on the dose, route of exposure and the time interval before starting treatment [8]. The clinical features can be broadly classified as secondary to (i) muscarinic effects, (ii) nicotinic effects and (iii) central receptor effects. Three different types of paralysis are recognized largely based on the time of occurrence and their differing pathophysiology (i) Type I paralysis or acute paralysis, (ii) Type II paralysis or Intermediate syndrome and (iii) Type III paralysis or Organophosphate induced delayed neuropathy [9]. Cholinesterase estimation is the only useful biochemical tool for confirming exposure to OP. In ingestional poisoning gastric lavage should be given and the sample collected must be sent for analysis and medico legal purposes. Atropine is the specific antidote for muscarinic effects and has no effect on nicotinic symptoms. It helps in reversing life threatening complications that may result in death such as central respiratory depression, bronchospasm, severe bradycardia and hypotension [10,11]. Current guidelines recommend use of incremental dose regimen followed by setting up an infusion to follow these end points. Oximes are also administered and their mechanism of action is by reactivating acetylcholinesterase that has been bound to the OP molecule [12].

4. CONCLUSION

This case highlights the unusual presentation of OP poisoning. Eliciting a proper and detailed

history and other clues found on examination will help in making an early diagnosis. The physicians have to be aware of the various intoxication syndromes caused by OP so that prompt and appropriate treatment can be provided. The various OP compounds have different outcomes due to its unique characteristics. All this should be kept in mind when approaching a patient with OP poisoning.

CONSENT

Written and oral consent was obtained by the patient before writing case report.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kumar S, Agarwal S, Raisinghani N, Khan S. Leukocyte count: a reliable marker for the severity of organophosphate intoxication? *J Lab Physicians*. 2018; 10(02):185-188.
2. Khan S, Kumar S, Agrawal S, Bawankule S. Correlation of serum cholinesterase and serum creatine phosphokinase enzymes with the severity and outcome of acute organophosphate poisoning: study in rural central India. *World J of Pharm and Pharmaceutical Sciences* 2016;5(4): 1365-1373.
3. Kumar S, Divan SK, Dubey S. Myocardial infarction in organophosphorus poisoning: Association or just chance? *J Emerg Trauma Shock* 2014;7:131-132
4. Pandit V, Seshadri S, Rao S N, Samarasinghe C, Kumar A, Valsalan R. A case of organophosphate poisoning presenting with seizure and unavailable history of parenteral suicide attempt. *J Emerg Trauma Shock* 2011;4:132-134
5. Patil A, Kumar S, Inamdar A, Acharya S, Wanjari A, Bawankule S, et al. Impact of Serum Amylase Level in the Outcome of Acute Organophosphorus Poisoning: 2-Year Cross-Sectional Study at Rural Teaching Hospital. *Journal of Laboratory Physicians*; 2021.

6. Agarwal A, Bhagawati J, Acharya S, Shukla, S. Organophosphate Poison Induced Delayed Polyneuropathy (Opidn) After Inhalational Poisoning- A Rare Sequel. International Journal of Medical Science and Clinical Invention. 2017;4(11):3271–3273.
7. Acharya S, Shukla S, Malpani V. An unusual case of triazophos poisoning presenting with New-Onset refractory status epilepticus. Toxicology International. 2015;22(1):172.
8. Batra AK, Keoliya AN, Jadhav GU. Poisoning: an unnatural cause of morbidity and mortality in rural India. J Assoc Physicians India. 2003;51: 955-959.
9. Aguilera L, Martinez-Bourio R, Cid C, Arino JJ, Saez de Eguilaz JL, Arizaga A, et al. Anaphylactic reaction after atropine. Anaesthesia. 1988;43:955-957.
10. Peter JV, Sudarsan TI, Moran JL. Clinical features of organophosphate poisoning: A review of different classification systems and approaches. Indian J Crit Care Med. 2014;18(11):735-45.
11. Eddleston M, Buckley NA, Eyer P, Dawson AH. Management of acute organophosphorus pesticide poisoning. Lancet. 2008;371(9612):597-607.
12. Zafar R, Munawar K, Nasrullah A, et al. Acute Renal Failure due to Organophosphate Poisoning: A Case Report. Cureus. 2017;9(7).

© 2021 Hepat et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/76851>*