



## Antibiotics Susceptibility Profile of *Listeria* Species Isolated from Poultry Wastes and Fishpond Water from Private and Institutional Farms in Ibadan, Nigeria

Olutayo Israel Falodun<sup>1\*</sup> and Moturayo Janet Amusan<sup>1</sup>

<sup>1</sup>Department of Microbiology, University of Ibadan, Ibadan, Nigeria.

### Authors' contributions

This work was carried out in collaboration between both authors. Author OIF designed the study and the protocol. Authors OIF and MJA managed literature search, data acquisition and wrote the draft. Both authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/JABB/2017/34566

#### Editor(s):

(1) Cosmas Nathanailides, Dept Fisheries and Aquaculture Technology, Technological Educational Institute of West Greece, Greece.

#### Reviewers:

(1) James K. Prah, University Of Cape Coast, Ghana.

(2) Nagahito Saito, Hokkaido University, Japan.

Complete Peer review History: <http://www.sciencedomain.org/review-history/19849>

Original Research Article

Received 31<sup>st</sup> May 2017  
Accepted 27<sup>th</sup> June 2017  
Published 4<sup>th</sup> July 2017

### ABSTRACT

**Introduction:** Untreated waste being discharged into the environment due to proliferation of poultry and fish farms can constitute a public health threat to human. *Listeria*, an emerging pathogen is commonly associated with food. This study aimed at determining the antibiotic resistant pattern of *Listeria* species isolated from poultry droppings and fish pond water in Ibadan.

**Materials and Methods:** Poultry waste and fishpond water samples were collected between April and July, 2016. *Listeria* Selective Agar was used to isolate *Listeria* species and identified using conventional methods. Antimicrobial susceptibility testing was done using the Kirby-Bauer disk diffusion method against ampicillin, cloxacillin, amoxicillin, streptomycin, ceftriazone, chloramphenicol, ciprofloxacin, ofloxacin, sulfamethoxazole-trimethoprim and tetracycline.

**Results:** A total of forty samples were collected and 105 *Listeria* spp. isolated; 62.9% from poultry waste and 37.1% from fishpond water. From the institutional fish pond and poultry waste, 18.1% and 51.4% isolates were obtained while from private fish pond and poultry waste, 19.1% and 11.4% isolates respectively were obtained. The isolates were *Listeria monocytogenes* (27.6%), *L. innocua*

\*Corresponding author: E-mail: oi.falodun@ui.edu.ng, falod2013@gmail.com;

(8.6%), *L. ivanovii* (16.2%) and other *Listeria* spp. (47.6%). All the isolates were resistant to ampicillin and ciprofloxacin; all *L. monocytogenes*, *L. innocua* and *L. ivanovii* were resistant to ceftriaxone, but 89.3% of the other *Listeria* spp. showed resistance. Furthermore, all (100%) the isolates obtained from private poultry farm were resistant to chloramphenicol while 53.3% were resistant to trimethoprim/sulfamethoxazole. Also, 12.2% of the isolates were found to be resistant to a combination of six antibiotics including: Ampicillin, amoxicillin, ceftriaxone, streptomycin and trimethoprim/sulfamethoxazole.

**Conclusion:** This study revealed that poultry waste and fish pond water from the selected farms could be a potential source for the transmission of multi-drug resistant bacteria to humans. Proper treatment of poultry waste and fish pond water should be ensured before discharge into the environment.

**Keywords:** *Listeria* species; antibiotics resistance; fish pond water, poultry waste.

## 1. INTRODUCTION

*Listeria* species are members of the family Listeriaceae, with eleven species of *Listeria* within the phylogeny and include *L. monocytogenes*, *L. ivanovii*, *L. grayi*, *L. innocua*, *L. welshmeri*, *L. seeligeri*, *L. marthii*, *L. maris*, *L. rocourtiae*, *L. fleischmannii* and *L. weihenstephanensis* [1]. Some *Listeria* spp. are pathogenic causing opportunistic infections in both human and animals leading to Listeriosis; among these, *Listeria ivanovii* and *L. monocytogenes* cause infections in animal and human [2]. Listeriosis is an emerging infectious disease of major public health concern worldwide because it is associated with food-borne outbreak with significant risk of morbidity and mortality [3]. The main route of entry of this pathogen is the intestinal mucosa of the host after ingestion. The clinical manifestations of listeriosis depends on the host and may be either invasive, when *L. monocytogenes* penetrates the blood-brain-barrier causing severe infections of the brain or non invasive, when the placental is penetrated leading to infection of the foetus. Invasive form of listeriosis affects high-risk people, including immunocompromised individuals, neonatal, and elderly persons usually present with septicemia or meningitis [4]. While non-invasive listeriosis, also referred to as febrile listerial gastroenteritis is a milder form of infection; It manifests symptoms such as diarrhea, fever, headache and muscle pain [5].

The most important aspect of this organism in food hygiene is the ability of the bacteria to survive in a wide range of temperatures and biofilms formation on various environmental surfaces, which serve as natural habitats or reservoirs [6]. The raising of food animals with antibiotics either as growth promoters or for infection control results in harboring significant

populations of antibiotic-resistant bacteria, which can be transmitted to humans through direct contact with the animals or through their products such as meat, eggs, and milk [7]. Aquatic ecosystems such as fish ponds and rivers have been recognized as a reservoir of antibiotic resistant bacteria and antibiotic resistance genes [8]. An outbreak of listeriosis was reported in a herd of cattle [9] from the south-western city of Ibadan while in a recent study conducted also in Ibadan on abattoir wastewater, *Listeria* spp. isolated from the wastewater include *L. monocytogenes* (26.8%), *L. innocua* (20.7%), *L. ivanovii* (18.3%) and other *Listeria* species (34.2%) [10].

A number of studies have reported isolation of *Listeria* species in Nigeria and elsewhere from food animals and their products such as pork, cattle and fish [11,12] as well as from environmental samples including municipal water, and abattoir wastewater [10,13]. However, there is a dearth of information from poultry waste and fish pond water especially in Nigeria. There are reports of resistance of *Listeria* spp. to antibiotics such as chloramphenicol and ampicillin [14]. In a study conducted on human samples and slaughterhouse, it was observed that *Listeria* species exhibited a high level of resistance to antibiotics such as clindamycin, daptomycin, fluoroquinolone as well as to the third and fourth generation antibiotics cephalosporins [15]. Also in a recent study carried out in Ibadan on samples collected from abattoir slaughter slab and drainage, all the *Listeria* spp. were found to be resistant to ampicillin together with a high resistant rate to ceftriaxone, amoxicillin/clavulanate and tetracycline [10]. This present study was therefore carried out to determine the occurrence and antibiotic resistant pattern of *Listeria* species isolated from poultry droppings and fish pond

water from private and tertiary farms in Ibadan metropolis.

## 2. MATERIALS AND METHODS

### 2.1 Study Area

The study sites were private poultry farm in Moniya, private fish farm in Ojoo and University of Ibadan poultry farm and fish pond. Ibadan is located in South-western part of Nigeria, it was founded in 1829; it is the largest city in West Africa. Ibadan is located at 70 24'N; 30 54'E; 234 m above sea level [16].

### 2.2 Sample Collection

Samples of poultry waste and fish pond water were aseptically collected into sterile sampling bottles. Samples were transported immediately to the laboratory in ice packs for microbiological analyses. Samples were collected weekly over a period of ten weeks between April and July, 2016.

### 2.3 Isolation and Identification of *Listeria* Species

Isolation of *Listeria* species was carried out according to the method of Akano et al. [17]. Analysis of the poultry wastes and fish pond water was carried out using the *Listeria* Selective Agar Base (Oxoid). The media was prepared according to manufacturers' instruction and supplemented with a vial of *Listeria* Selective Supplement. The poultry wastes and fish pond water were serially diluted and the standard pour plate technique was used by plating out 1 ml of the appropriate dilutions on the *Listeria* selective agar. The cultured plates were incubated at 35°C for 24-48 hours. Colonies with gray colour and dark background, were sub-cultured on *Listeria* Selective Agar media to obtain pure isolates and the isolates were identified using standard biochemical tests.

### 2.4 Antibiotics Susceptibility Test of the Isolates

Using the standard disk diffusion technique on Mueller-Hinton agar, antibiotics susceptibility test of the isolates was carried out based on the recommendation of Clinical Laboratory Standards Institute [18]. A total of ten antibiotics obtained from Oxoid, U.K. were used which include: tetracycline (30 µg), streptomycin (10

µg), cloxacillin (5 µg), ofloxacin (5 µg), amoxicillin (5 µg), ceftriaxone (30 µg), sulfamethoxazole-trimethoprim (25 µg), chloramphenicol (30 µg), ampicillin (10 µg) and ciprofloxacin (5 µg). The isolates were sub cultured and colonies of 18-24 hour old culture was picked and suspended in a tube containing sterile normal saline with turbidity adjusted to 0.5 McFarland standards. The suspension was uniformly spread over already prepared solidified Mueller Hinton agar plates using a sterile swab stick. Sterile forceps was used to aseptically place the antibiotics disc on the inoculated plates and incubated at 37°C for 18-24 hours. After the incubation period, the zones of inhibition were measured, recorded and interpreted using the CLSI standards.

## 3. RESULTS

A total of 105 *Listeria* spp. were isolated from poultry waste and fishpond water samples comprising of 66 (62.9%) and 39 (37.1%) isolates from poultry waste and fishpond water samples respectively. The occurrence of the isolates showed that 29 (27.6%) were *Listeria monocytogenes*; 17 (16.2%) were *L. ivanovii*, 9 (8.6%) were *L. innocua* while 50 (47.6%) belongs to other *Listeria* species (Table 1).

The antibiotic susceptibility test results revealed that all (100%) the *Listeria* isolates were resistant to ampicillin and ciprofloxacin. Likewise, all (100%) the *L. ivanovii* were resistant to cloxacillin and similarly, all (100%) the *L. ivanovii*, *L. innocua* and *L. monocytogenes* were resistant to ceftriaxone. Furthermore, the results of the susceptibility test showed that resistant to ceftriaxone, cloxacillin and sulfamethoxazole-trimethoprim were 97.1%, 83.8% and 53.3% respectively while it was observed that among the isolates that showed resistant to sulfamethoxazole-trimethoprim, the highest was among the *L. ivanovii* (58.8%) followed by *L. monocytogenes* (58.6%), the other *Listeria* spp. (50.0%) and *L. innocua* (44.4%). However, a low resistant rate was observed among the isolates to tetracycline (29.5%), chloramphenicol (15.2%) and streptomycin (32.4%). Moreover, none of the *L. ivanovii*, *L. innocua* and *L. monocytogenes* showed any resistance to ofloxacin (Table 2).

In addition, all (100%) the isolates obtained from the fish pond water exhibited resistant to ceftriaxone while it was 94.4% resistance for isolates from the poultry waste. While the *Listeria* spp. isolated from the poultry waste showed

1.5%, 22.7% and 39.4% resistance to ofloxacin, chloramphenicol and tetracycline respectively, none of the isolates obtained from the fish pond water samples showed any resistant to the antibiotics. However, all (100%) the isolates obtained from the institution fish pond water samples were resistant to sulfamethoxazole-trimethoprim while those isolated from the private fishpond water showed resistance of 50% to the antibiotic. Similarly, the rate of resistance of the *Listeria* isolates from the private poultry farm to tetracycline (100%), sulfamethoxazole-trimethoprim (100%) and chloramphenicol (91.7%) were high while the ones isolated from the institutional fishpond showed a lower resistance of 25.9%, 29.6% and 7.4% to the antibiotics respectively (Table 3).

The result of the phenotypes of resistance of the isolates to antibiotics from the fish pond water and the poultry waste is as shown in Table 4 and Table 5 respectively. The result showed that from fishpond water, all (100%) the isolates were resistant to the combination of three antibiotics that included ampicillin, ceftriaxone and ciprofloxacin while 74.4% of the isolates were resistant to another set of three drugs which include: sulfamethoxazole-trimethoprim, ceftriaxone and ciprofloxacin. It was also observed that 41.0% of the isolates were resistant to the following four drugs: streptomycin, sulfamethoxazole-trimethoprim, ceftriaxone and ciprofloxacin while 35.9% of the isolates were resistant to the combination of another set of four drugs including amoxicillin, sulfamethoxazole-trimethoprim, ceftriaxone and ciprofloxacin. Furthermore, among the isolates obtained from the poultry wastes, 72.7% were resistant to the combination of ampicillin, ceftriaxone and ciprofloxacin while it was observed that 10.5% were resistant to a combination of six antibiotics including ampicillin, streptomycin, sulfamethoxazole-trimethoprim, chloramphenicol, ciprofloxacin and tetracycline while 1.5% of the isolates were resistant to a combination of another set of six antibiotics including ampicillin, tetracycline, amoxicillin, ciprofloxacin, sulfamethoxazole-trimethoprim and chloramphenicol (Table 5).

#### 4. DISCUSSION

In Nigeria, little is known about the true state of listeriosis especially in relation to the environment because there have not been comprehensive studies carried out on *Listeria* in the country. Although, listeriosis is an important

disease of public health and animals coupled with the possibility of the environment being a route of transmission, there has been little or no attention given to it. Isolation of *L. monocytogenes* and *L. ivanovii* from both the fish pond water and the poultry waste studied is an indication that the possibility of these environmental samples serving as route of transmission of the pathogens to both human and animals cannot be ruled out because these *Listeria* species has been reported to be responsible for illness in human and animals. While *L. monocytogenes* is principally a pathogen of human and animals, *L. ivanovii* is a pathogen of animals as well as being implicated in human infection occasionally [2]. Different species of *Listeria* were isolated from the environmental samples studied which corroborates previous reports in which *Listeria* species were isolated from environmental samples including abattoir wastewater, butchers tables, fresh water samples and municipal waste effluents [1,10,15,19]. It was observed from this study that the occurrence of the isolates was 27.6% (*L. monocytogenes*), 16.2% (*L. ivanovii*) and 8.6% (*L. innocua*); This is similar to the rate of occurrence of the isolates in a recent study conducted on abattoir wastewater in which *L. monocytogenes* had the highest rate of occurrence compared to the other species [10]. However, from the fish pond, the occurrence of the isolates as observed in this study was lower compared to same isolates from the poultry droppings. The reason for this may be as a result of the direct introduction of antibiotics into the fish pond water which might have possibly eliminated the susceptible strains thereby reducing the microbial load.

The resistance of the isolates obtained in this study to ampicillin (100%) is comparable to the (total resistance) reported from previous studies which was carried out on homemade white cheese and fishes respectively [20,21]. In a study carried out in South Africa [13], resistance of *Listeria* to ciprofloxacin (91%) was high but a bit lower compared to the 100% observed in this study, the little disparity may be due to the studied samples. The South African study was on municipal wastewater. However, this observation is not in agreement with 0% ciprofloxacin resistance reported from a study on ready to eat foods in South Africa as well as another study from Mangalore, India on clinical and food samples [22,23]. Generally, resistance of the isolates in this study to chloramphenicol (15.2%), tetracycline (29.5%)

and ofloxacin (0.9%) was lower compared to the resistance reported on abattoir wastewater to the same set of antibiotics which were 41.5%, 65.5% and 6.1% for chloramphenicol, tetracycline and ofloxacin respectively [10]. However, the latter study was on abattoir wastewater as against the poultry waste and fish pond water in this study.

**Table 1. Number and percentage of occurrence of *Listeria* spp. isolated from the environmental samples n(%)**

Source	<i>Listeria monocytogenes</i>	<i>Listeria ivanovii</i>	<i>Listeria innocua</i>	<i>Listeria spp.</i>	Total
Poultry waste (Institution)	12 (11.4)	12 (11.4)	2 (1.9)	28 (26.7)	54 (51.4)
Poultry waste (Private)	7 (6.7)	1 (1.0)	0 (0.0)	4 (3.8)	12 (11.4)
Fish pond water (Institution)	6 (5.7)	2 (1.9)	3 (2.9)	8 (7.6)	19 (18.1)
Fish pond water (Private)	4 (3.8)	2 (1.9)	4 (3.8)	10 (9.5)	20 (19.1)
Total	29 (27.6)	17 (16.2)	9(8.6)	50 (47.6)	105 (100.0)

**Table 2. Antibiotic resistance patterns of all the *Listeria* spp. isolated from the environmental samples n(%)**

Antibiotics	<i>L. monocytogenes</i> N=29	<i>L. ivanovii</i> N=17	<i>L. innocua</i> N=9	<i>Listeria spp.</i> N=50	Total N=105
Ampicillin (10µg)	29 (100.0)	17 (100.0)	9 (100.0)	50 (100.0)	105 (100.0)
Streptomycin (10µg)	11 (37.8)	5 (29.4)	1 (11.1)	17 (34)	34 (32.4)
Ceftriazone (30µg)	29 (100.0)	17 (100.0)	9 (100.0)	47 (94.0)	102 (97.1)
Chloramphenicol (30µg)	7 (24.1)	2 (11.7)	0 (0.0)	7 (14.0)	16 (15.2)
Sulfamethoxazole	17 (58.6)	10 (58.8)	4 (44.4)	25 (50.0)	56 (53.3)
Trimethoprim (25µg)					
Cloxacillin (5µg)	26 (89.7)	17 (100.0)	7 (77.8)	38 (76.0)	88 (83.8)
Tetracycline (30µg)	8 (27.6)	5 (29.4)	1 (11.1)	17 (34.0)	31 (29.5)
Ciprofloxacin (5µg)	29 (100.0)	17 (100.0)	9 (100.0)	50 (100.0)	105 (100.0)
Ofloxacin (30µg)	0 (0.0)	0 (0.0)	0 (0.0)	1 (2.0)	1 (0.9)
Amoxicillin (25µg)	7 (24.1)	6 (35.3)	0 (0.0)	14 (28.0)	27 (54.0)

**Table 3. Antibiotic resistance patterns of the *Listeria* spp. isolated from different sources**

Antibiotics	FPI n=19	FPP n=20	PWI n=54	PWP n=12	FPW n=39	PW n=66
Ampicillin (10µg)	19 (100.0)	20 (100.0)	54 (100.0)	12 (100.0)	39 (100.0)	66 (100.0)
Streptomycin (10µg)	5 (26.3)	12 (60.0)	12 (22.2)	5 (41.7)	17 (43.6)	17 (25.8)
Ceftriazone (30µg)	19 (100.0)	20 (100.0)	51 (94.4)	12 (100.0)	39 (100.0)	63 (95.5)
Chloramphenicol (30µg)	0 (0.0)	0 (0.0)	4 (7.4)	11 (91.7)	0 (0.0)	15 (22.7)
Sulfamethoxazole	19 (100.0)	10 (50.0)	16 (29.4)	12 (100.0)	29 (74.4)	28 (42.4)
Trimethoprim(25µg)						
Cloxacillin (5µg)	17 (89.5)	16 (80.0)	48 (88.9)	7 (58.3)	33 (84.6)	55 (83.3)
Tetracycline (30µg)	0 (0.0)	0 (0.0)	14 (25.9)	12 (100.0)	0 (0.0)	26 (39.4)
Ciprofloxacin (5µg)	19 (100.0)	20 (100.0)	54 (100.0)	12 (100.0)	39 (100.0)	66 (100.0)
Ofloxacin (30µg)	0 (0.0)	0 (0.0)	1 (1.9)	0 (0.0)	0 (0.0)	1 (1.5)
Amoxicillin (25µg)	10 (50.0)	10 (50.0)	8 (14.8)	0 (0.0)	20 (51.3)	8 (12.1)

Keys FPI: Fishpond water from the institution, FPP: Fish pond water from the private farm  
 PWI: Poultry waste from the institution, PWP: Poultry waste from private farm  
 FPW: Total from both fish pond water, PW: Total from both poultry waste

**Table 4. Antibiotypes resistance pattern of *Listeria* species isolated from fish pond water samples. n(%)**

Antibiotypes	<i>L. monocytogenes</i> (n=10)	<i>L. ivanovii</i> (n=4)	<i>L. innocua</i> (n=7)	<i>Listeria</i> spp. (n=18)	Total (n=39)
AMX-STR	4 (10.3)	1 (2.6)	0 (0.0)	2 (5.1)	7 (10.3)
STR-SXT	5 (12.8)	1 (2.6)	4 (10.3)	6 (15.4)	16 (41.0)
SXT-AMX	4 (10.3)	3 (7.7)	0 (0.0)	7 (10.3)	14 (35.9)
AMP-CRO-CIP	10 (25.6)	4 (10.3)	7 (18.0)	18 (46.2)	39 (100)
AML-SXT-STR	4 (10.3)	1 (2.6)	0 (0.0)	2 (5.1)	7 (10.3)
SXT-CIP-CRO	8 (20.5)	3 (7.7)	4 (10.3)	14 (35.9)	29 (74.4)
STR-CIP-CRO	5 (12.8)	2 (5.1)	4 (10.3)	9 (23.1)	20 (51.3)
AMX-CIP-CRO	4 (10.3)	3 (7.7)	0 (0.0)	8 (20.5)	15 (38.5)
SXT-CRO-CIP-AMX	4 (10.3)	3 (7.7)	0 (0.0)	7 (10.3)	14 (35.9)
STR-AMX-CRO-CIP	4 (10.3)	1 (2.6)	0 (0.0)	2 (5.1)	7 (10.3)
SXT-STR-CIP-CRO-AMP	5 (12.8)	1 (2.6)	4 (10.3)	6 (15.4)	16 (41.0)

**Table 5. Antibiotypes resistance pattern of *Listeria* species isolated from poultry waste n(%)**

Antibiotypes	<i>L. monocytogenes</i> n= 19	<i>L. ivanovii</i> n=13	<i>L. innocua</i> n=2	<i>Listeria</i> spp. n=32	Total n =66
CRO-CIP	12 (18.2)	11 (16.7)	2 (3.0)	23 (34.8)	48 (72.7)
AMX-STR	0 (0.0)	1 (1.5)	0 (0.0)	0 (0.0)	1 (1.5)
STR-SXT	1 (1.5)	3 (4.5)	0 (0.0)	3 (4.5)	7 (10.5)
SXT-AMX	0 (0.0)	1 (1.5)	2 (3.0)	0 (0.0)	3 (4.5)
AMP-CRO-CIP	12 (18.2)	11 (16.7)	0 (0.0)	23 (34.8)	48 (72.7)
AML-SXT-STR	0 (0.0)	1 (1.5)	0 (0.0)	0 (0.0)	1 (1.5)
SXT-CIP-CRO	2 (3.0)	3 (4.5)	0 (0.0)	4 (6.0)	9 (13.5)
STR-CIP-CRO	3 (4.5)	2 (3.0)	0 (0.0)	2 (3.0)	7 (10.5)
AMX-CIP-CRO	3 (4.5)	3 (4.5)	0 (0.0)	2 (3.0)	8 (12.0)
SXT-CRO-CIP-AMX	0 (0.0)	1 (1.5)	0 (0.0)	0 (0.0)	1 (1.5)
STR-AMX-CRO-CIP	0 (0.0)	1 (1.5)	0 (0.0)	0 (0.0)	1 (1.5)
SXT-STR-CIP-CRO	1 (1.5)	2 (3.0)	0 (0.0)	1 (1.5)	4 (6.0)
TET-STR-CIP-CRO	0 (0.0)	2 (3.0)	0 (0.0)	1 (1.5)	3 (4.5)
AMP-STR-SXT-C-CIP-TET	3 (4.5)	0 (0.0)	0 (0.0)	4 (6.0)	7 (10.5)
AMP-TET-AMX-CIP-SXT-CHL	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.5)	1 (1.5)

Keys: TET- Tetracycline, STR- Streptomycin, CHL- Chloramphenicol, CRO- Ceftriaxone, CIP- Ciprofloxacin, SXT- Sulfamethoxazole-trimethoprim, AMP- Ampicillin, AMX- Amoxicillin

However, the results from this present study also revealed that the isolates from the fish pond water samples had a higher resistance (74.4%) to sulfamethoxazole-trimethoprim than that of the poultry waste samples (42.4%). This may be as a result of direct release of antibiotics into the fish pond. In addition, the total resistance to sulfamethoxazole-trimethoprim observed in both the institutional fish pond isolates and the private poultry farm isolates were much higher than 50% and 29.6% resistance which was observed in the isolates from the private fish pond water and institution poultry farm respectively. The reason for these differences may be as a result of misuse of this antibiotic on different farms sampled. This observation however is not in agreement with the resistance rate (17.1%) reported from a study on frozen burger patties in

Malaysia [24]. Similarly, the resistance observed for *L. monocytogenes* (100%), *L. ivanovii* (100%), *L. innocua* (100%) and the other *Listeria* spp. (94%) to ceftriaxone in this study is a bit higher than the ones reported by Falodun *et al.* [10] where resistance to the same antibiotic was 90.9% (*L. monocytogenes*), 82.4% (*L. innocua*), 73.3% (*L. ivanovii*) and 92.9% (other *Listeria* spp.). The observation from the result of the susceptibility test in this study showed that resistance of the isolates obtained from the fish pond water samples (100%) and poultry waste samples (95.5%) to ceftriaxone were high; the high resistant to this third generation antibiotic may be as a result of the development of natural resistance to the antibiotic. Moreover, resistance of the isolates to chloramphenicol (0%) as observed in this study for both the institutional

and private fish pond water samples, is in agreement with (0%) resistance reported in Malaysia and Mangalore, India [23,25]. Whereas, the resistance (91.7%) of the isolates from the poultry waste of the private farm to chloramphenicol is much higher than the 7.4% resistance obtained from the poultry waste of the institutional farm. This may be due to possible misuse of this antibiotic in the private poultry farm, while it may be that there is possibility of proper regulation of antibiotics usage in the Institutional farm.

## 5. CONCLUSION

In conclusion, the result of this study has further confirmed a high occurrence of *Listeria* species in environmental samples; *L. monocytogenes* is prevalent in the studied samples as well as a sizeable percentage of occurrences of *L. ivanovii*. The prevalence of these two species which have been implicated as human and animal pathogens could lead to the outbreak of listeriosis, which may results into death in poultry and fishes. The high antibiotic resistance exhibited by *L. monocytogenes* and *L. ivanovii* in these two environmental samples to ampicillin, ciprofloxacin and sulfamethoxazole-trimethoprim is suggestive of possibility of misuse of antibiotics either for therapeutic or other purposes in both fish pond and the poultry. This may lead to emergence, selection and dissemination of antibiotic resistant pathogenic *Listeria* species. Similarly, the high resistant exhibited by the *Listeria* spp. in this study to the third generation cephalosporins (ceftriaxone) which is the last resort for the treatment of listeriosis portend a difficult scenario for the treatment of listeriosis infection in the study area if there is outbreak of infection. Hence, adequate measures are to be put in place to regulate the use of antibiotics in animal husbandry and treatment of effluents and waste generated before they are discharged into the environment.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

- Halter EL, Neuhaus K, Scherer S. *Listeria weihenstephanensis* spp. isolated from the water plant *Lemna trisulca* taken from a freshwater pond. Int J Syst Evo Microbiol. 2013;63:641-647.
- Low JC, Donachie W. A review of *Listeria monocytogenes* and listeriosis. Vet J. 1997;153:9-29.
- Evans MR, Swaminathan B, Graves LM, Altermann E, Klaenhammer TR, Fink RC. Genetic markers unique to *Listeria monocytogenes* serotype 4b differentiate epidemic clone II (hot dog outbreak strains) from other lineages. Appl Environ Microbiol. 2004;70:2383-90. DOI: 10.1128/AEM.70.4.2383-2390
- Barry AL, Thornsberry C. Susceptibility tests: Diffusion test procedures. In manual of clinical microbiology, edited by Lennette EH, Balows A, Hausler WJ, Shadomy HJ. Washington DC: Am Soc Microbiol; 1997.
- Ooi ST, Lorber B. Gastroenteritis due to *Listeria monocytogenes*. Clin Infect Dis. 2005;40:1327-32.
- Mafu A, Roy D, Goulet J, Magny P. Attachment of *Listeria monocytogenes* to stainless steel, glass, polypropylene and rubber surfaces after short contact times. J Food Prot. 1990;53:742-746.
- Marshal BM, Levy SB. Food animals and antimicrobials impacts on human health. Clin Microbiol Rev. 2011;24:718-733.
- Baquero F, Martinez JL, Canton R. Antibiotics and resistance in water environments. Curr Opin Biotechnol. 2008;19(3):260-265.
- Akpavie SO, Ikheloa JO. An outbreak of listeriosis in cattle in Nigeria. Revue d'Elevage et de Med Vet des Pays Trop. 1992;45:263-264.
- Falodun OI, Rabi AG, Fagade OE. Antibiotics susceptibility profile of *Listeria* species isolated from untreated abattoir wastewater in Akinyele, Ibadan, Nigeria and its implication on public health. British Biomed Res J. 2016;16(6):1-10. DOI: 10.9734/bmrj/2016/28066
- Atil E, Ertas HB, Ozbey G. Isolation and molecular characterization of *Listeria* spp. from animals, food and environmental samples. Vet Med. 2011;56(8):386-94.
- Abhay VR, Swapnil PD, Krupali VP, Ajay P, Saroj B, Barbuddhe SB. Isolation and genotypic characterization of *Listeria monocytogenes* from pork and pork products. Int J Curr Microbiol Appl Sci. 2015;4(1):788-798.
- Nwaiwu O. Presence of *Listeria* spp. in fish samples, fish products and sea products. Int Food Res J. 2015;22(2):455-464.

14. Odjadjare EEO, Obi CL, Okoh AI. Municipal wastewater effluents as a source of listerial pathogens in the aquatic milieu of the Eastern Cape Province of South Africa: A concern of public health importance. *Int J Environ Res Public Health*. 2010;7:2376-239.
15. Moreno LZ, Paixão R, Gobbi DDS, Raimundo DC, Ferreira TP, Moreno AM, et al. Characterization of antibiotic resistance in *Listeria* spp. isolated from slaughterhouse environments, pork and human infections. *J Infect Dev Countries*. 2014;8(4):416-423.
16. Lloyd PC, Mabogunje AL, Awe B. *The City of Ibadan*. Cambridge University Press, Cambridge; 1967.
17. Akano SO, Moro DD, Agboola AM, Oluwadun A. Public health Implication of *Listeria species* and other bacteria isolates of abattoir effluent in Lagos, Nigeria. *Int Res J Microbiol*. 2013;4(7):162-167.
18. Clinical and Laboratory Standards Institute (CLSI). *Performance standards for antimicrobial disk susceptibility tests; 9<sup>th</sup> ed.; Document M2-A9; Clinical and Laboratory Standards Institute (CLSI): Wayne, PA, USA; 2011.*
19. Nwachukwu NC, Orji FA. Studies on the Isolation of *Listeria monocytogenes* from food, water, and animal droppings: Environmental Health Perspective, *Environmental Health – Emerging Issues and Practice*. 2012;978-953-307-854.
20. Arslan S, Ozdemir F. Prevalence and antimicrobial resistance of *Listeria* species in homemade white cheese. *Food Control*. 2008;19:360-363.
21. Issa ZM, Mustakim M, Mohamed SAS, Muda NM, Yen LH, Radu S. Antibigram Profiles of *Listeria monocytogenes* isolated from foods. *Int Conf Biotechnol Food Sci*. 2011;7:133-137.
22. Nyenje ME, Ndip RN, Tanih NF, Green E. Current status of antibiograms of *Listeria ivanovii* and *Enterobacter cloacae* isolated from ready-to-eat foods in Alice, South Africa. *Int J Environ Res Public Health*. 2010;9:3101-3114.
23. Dhanashree B, Otta SK, Karunasagar I, Goebel W, Karunasagar I. Incidence of *Listeria* spp. in clinical and food samples in Mangalore, India. *Food Microbiol*. 2003;20: 447–453.
24. Wong WC, Pui CF, Tunung R, Ubong A, Noor HMS, Farinazleen MG. Antibigram Pattern among Cultures of *Listeria monocytogenes* isolated from frozen burger patties in Malaysia. *J Trop Agric Sci* 2003;35:793-804.
25. Marian MN, Aminah SM, Zuraini MI, Son R, Maimunah M, Lee HY. “MPN-PCR detection and antimicrobial resistance of *Listeria monocytogenes* isolated from raw and ready-to-eat foods in Malaysia. *Food Control*. 2012;28:309-14.

© 2017 Falodun and Amusan; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<http://sciedomain.org/review-history/19849>