



Common Causes of Neonatal Death in Calabar, South-South Nigeria: An Autopsy Study

Ugbem Theophilus Ipeh ^a, Akaba Kingsley ^{b*},
Ushie David Edem ^a, Juliet Venn ^c, Nnoli Martin ^a,
Ebughe Godwin Abeng ^a and Oshatuyi Olukayode ^d

^a Department of Pathology, University of Calabar Teaching Hospital, Calabar, Nigeria.

^b Department of Haematology and Blood Transfusion, University of Calabar Teaching Hospital, Calabar, Nigeria.

^c Department of Paediatrics, University of Calabar Teaching Hospital, Calabar, Nigeria.

^d Department of Medical Microbiology and Parasitology, University of Calabar, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ARRB/2023/v38i430578

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/92841>

Original Research Article

Received: 28/10/2022

Accepted: 31/12/2022

Published: 06/05/2023

ABSTRACT

Background: Neonatal deaths are deaths of live born babies occurring before 28 completed days. The vast majority occur in low-income countries like Nigeria with a high neonatal mortality rate. There is paucity of autopsy studies due to refusal of family to give consent for such procedures.
Aim: To identify the commonest causes of neonatal death in 53 neonatal autopsies in Calabar, Nigeria.
Materials and Methods: Detailed postmortem was carried out using lettuces techniques and bits taken for histological analysis to ascertain the cause of death. Other relevant contributory factors

*Corresponding author: E-mail: akaba_kingsley@yahoo.com;

such as the gestational age, mode of delivery, place of birth, antemortem cause of deaths and maternal obstetric history were obtained from the medical records and autopsy request forms.

Results: In the one-year retrospective study of 53 neonatal autopsies, male: female ratio was 1: 0.83 and mean age at death was 6.5 + 7.3 days, ranging from 1 to 28 days. The commonest cause of neonatal death was severe birth asphyxia seen in 10 cases (18.9%), followed by kernicterus in 6 cases (11.3%), birth trauma seen in 6 cases (11.3%), congenital heart disease seen in 5 cases (9.4%), and prematurity seen in 5 cases (9.4%).

Conclusion: The study confirms the usefulness of neonatal autopsy in ascertaining the definitive cause of death. Severe birth Asphyxia was identified as the commonest cause of death in the neonatal period followed by birth trauma, kernicterus and congenital heart diseases.

Keywords: Neonatal death; neonatal autopsy; cause of death; severe birth asphyxia.

1. INTRODUCTION

“Neonatal period is defined as the time between birth and 28 days of life. It is a highly vulnerable period of life when a neonate may develop certain serious problems which may lead to death” [1]. “Early neonatal deaths are deaths of live born babies during the first seven completed days after birth, and late neonatal deaths occur after 7 completed days and before 28 completed days” [1]. “The vast majority of neonatal deaths occur in the low-income countries where standards of both maternal and newborn care are low and of the approximately 130 million infants born worldwide each year, it is estimated that four million infants die during the first month of life” [2–4]. “Children who die within the first 28 days of birth suffer from conditions and diseases associated with lack of quality care at birth or skilled care and treatment immediately after birth and in the first days of life (WHO). Nigeria as of 2019 fall within the first 10 countries with the highest number of newborn deaths” [5–7]. “The first step in improving early neonatal survival is to identify the common causes and look for modifiable factors and develop policies such as quality neonatal care that is easily available and affordable to the common man within the geopolitical zone. Empowerment and training of community health workers and traditional birth attendants at the community level will help in the reduction of maternal and neonatal mortality. However, few developing countries have vital registration systems that are complete enough to provide accurate estimates of neonatal mortality, or of its early and late components” [8–10].

2. MATERIALS AND METHODS

A retrospective study of 53 neonatal autopsies conducted in the Department of Pathology,

University of Calabar Teaching Hospital within a period of a year. Detailed postmortem was carried out using lettuces techniques and bits taken for histological analysis to ascertain the cause of death. Other relevant contributory factors such as the gestational age, mode of delivery, place of birth, antemortem cause of deaths and maternal obstetric history were obtained from the medical records and autopsy request forms. The data obtained was transferred into SPSS version 21.0 and statistical analysis carried out.

3. RESULTS

In the one-year retrospective study of 53 neonatal autopsies, male : female ratio was 1: 0.83 and mean age at death was 6.5 + 7.3 days, ranging from 1 to 28 days, with a little less than one-fifth [10] i.e. 18.9% dying within 24 hours and one-half (26) i.e. 49.1%, dying within 2-7 days of life, respectively. The commonest cause of neonatal death was severe birth asphyxia seen in 10 cases (18.9%), followed by kernicterus in 6 cases (11.3%), birth trauma seen in 6 cases (11.3%), congenital heart disease seen in 5 cases (9.4%), and prematurity seen in 5 cases (9.4%). As shown in Table 7 below. Table 1 presents the sociodemographic data of the mothers, antenatal care, fetal presentation and the places of delivery. The type of labour, mode of delivery and clinical presentations are presented in Table 2. Clinical diagnosis, wards and length of stay are presented in Table 3. Findings from the skin, scalp, and brain are presented in Table 4. That cardiac and lungs findings are presented in Table 5. Findings from the liver, spleen, GIT, adrenal gland, and the urinary system are presented in Table 6.

Table 1. Sociodemographic variables

Variable	Frequency	Percentage
Maternal Age Group		
16-20 years	8	15.1
21-25 years	13	24.5
26-30 years	17	32.1
31-35 years	12	22.6
36-40 years	2	3.8
41-45 years	1	1.9
Maternal Educational Level		
Primary	8	15.1
Secondary	32	60.4
Tertiary	11	20.8
No formal Education	2	3.8
Others	0	0.0
Antenatal Care		
UCTH	21	39.6
General Hosp	9	17.0
Private Hosp	4	7.5
Other Hosp	2	3.8
Unbooked	17	32.1
Foetal Presentation		
Cephalic	48	90.6
Breech	3	5.7
Transverse	2	3.8
Others	0	0.0
Place of Delivery		
UCTH	26	49.1
General Hosp	9	17.0
Private Hosp	3	5.7
Other Hosp	2	3.8
Traditional Birth Facility	4	7.5
Church	4	7.5
Home	5	9.4
Other	0	0.0

Table 2. Showing type of labour, mode of delivery and clinical presentations

Variable	Frequency	Percentage
Labour		
Spontaneous	30	56.6
Induced	16	30.2
Others	7	13.2
Labour Progress		
Normal	16	30.2
Prolonged	32	60.4
Others	5	9.4
History of Foetal Distress		
Yes	38	73.1
No	14	26.9
Liquor Colour		
Colourless	30	56.6
Meconium-stained	22	41.5
Others	1	1.9
Mode of Delivery		
SVD(Normal)	40	75.5

Variable	Frequency	Percentage
Forceps Delivery	0	0.0
Caesarean Section	11	20.8
Breech	2	3.8
Others	0	0.0
Gender	29	54.7
Male		
Female	24	45.3
Clinical presentation		
Low birth weight	2	3.8
prematurity	13	24.5
postdate	11	20.8
cord around neck	2	3.8
true knot	0	0.0
jaundice	3	5.7
spasms	2	3.8
Fever	3	5.7
Fast breathing	5	9.4
convulsion	0	0.0
poor cry	5	9.4
Abdominal mass	1	1.9
poor suck	4	7.5
ruptured omphalocele	1	1.9
cellulitis	1	1.9

Table 3. Showing clinical diagnosis, wards and length of stay

Variable	Frequency	Percentage
Clinical Diagnosis		
Severe Birth Asphyxia	22	41.5
Acute Respiratory Distress Syndr	1	1.9
Prematurity	6	11.3
Neonatal Sepsis	9	17.0
Neonatal Jaundice (Kernicterus)	3	5.7
Neonatal Tetanus	3	5.7
Meconium Aspiration	1	1.9
Congenital Heart Disease	3	5.7
Congenital Pneumonia	3	5.7
Congestive Cardiac Failure	1	1.9
Posterior Urethral Valve	1	1.9
Lung Collapse	0	0.0
Ward	35	66.0
Sick Baby Unit (SBU)		
Special Care Baby Unit (SCBU)	14	26.4
Labour Ward	4	7.5
Others	0	0.0
Age Group	36	67.9
1-7 days		
8-14 days	11	20.8
15-21 days	2	3.8
22-28 days	4	7.5
Length of Stay	43	81.1
1-7 days		
8-14 days	4	7.5
15-21 days	2	3.8
22-28 days	4	7.5

Table 4. Showing the findings from the skin, scalp, and brain

Variable	Frequency	Percentage
Skin Findings	2	3.8
Cellulitis		
Lividity	24	45.3
Jaundice	10	18.9
Meconium staining	11	20.8
Haemangiomas	0	0.0
cafe au lait spots	0	0.0
Pustules	0	0.0
scaly lesions	0	0.0
petechiae haemorrhage	6	11.3
Scalp Findings	3	5.7
Subgaleal hemorrhage		
Cephalhaematoma	30	56.6
Laceration	4	7.5
puncture needle wounds	12	22.6
caput succedenum	4	7.5
Brain Findings	2	3.8
Epidural haemorrhage		
Subdural haemorrhage	0	0.0
Subarachnoid haemorrhage	4	7.5
Intraventricular haemorrhage	1	1.9
Intraparenchymal haemorrhage	0	0.0
Meningitis	0	0.0
Kernicterus	6	11.3
Tonsillar herniation	3	5.7
cerebral oedema	19	35.8
cerebral congestion	18	34.0
staining of basal ganglia and ventricles	0	0.0

Table 5. Showing the cardiac and lungs findings

Variable	Frequency	Percentage
Cardiac Findings	14	26.4
Cardiomegaly		
Univentricular heart	1	1.9
ASD	11	20.8
VSD	3	5.7
PDA	0	0.0
TOF	0	0.0
TGA	1	1.9
Left ventricular hypertrophy	3	5.7
Flabby Heart	13	24.5
Haemopericardium	1	1.9
No obvious pathology	6	11.3
Findings in the Lungs	0	0.0
Hypoplasia of lungs		
Bronchial mucosa plug	0	0.0
Consolidation	20	37.7
Pulmonary haemorrhage	5	9.4
Meconium aspiration	1	1.9
Pulmonary embolus	0	0.0
Hyperinflated lungs	0	0.0
Severe venous congestion	8	15.1
Pulmonary edema	17	32.1
Pneumonia	2	3.8


Table 6. Showing findings from the liver, spleen, GIT, adrenal gland, and the urinary system

Variable	Frequency	Percentage
Liver Findings		
Hepatomegaly	5	9.4
Hepatic cyst	1	1.9
Hemangioma	1	1.9
Chronic venous congestion	45	84.9
Viral hepatitis	0	0.0
Fatty Change	1	1.9
Spleen		
Splenomegaly	2	3.8
Malaria spleen	1	1.9
Splenitis	8	15.1
Diffluent spleen	5	9.4
chronic congestion	37	69.8
GIT Findings		
Malrotation	0	0.0
Omphalocele	1	1.9
Imperforate anus	0	0.0
Necrotizing enterocolitis	1	1.9
gangrenous bowel	1	1.9
pneumatosis intestinalis	6	11.3
No Obvious pathology	44	83.0
Adrenal gland		
Adrenal haemorrhage	13	24.5
Adrenal hyperplasia	1	1.9
Adrenal hypoplasia	0	0.0
No obvious pathology	39	73.6
Urinary System		
Shocked kidneys	50	94.3
Renal cyst	1	1.9
Hydronephrosis	1	1.9
Megaureter	0	0.0
Posterior urethra valve	0	0.0
Horse shoe kidney	1	1.9

Table 7. Showing the causes of death

Variable	Frequency	Percentage
Cause of Death		
ARDS	2	3.8
Pulmonary Haemorrhage	2	3.8
Bronchopneumonia	1	1.9
Congestive Cardiac Failure	1	1.9
Congenital Heart Disease	4	7.5
Kernicterus	6	11.3
Birth Trauma	6	11.3
Severe Birth Asphyxia	10	18.9
Aspiration Pneumonitis	1	1.9
Subtotal Lung Collapse	0	0.0
Neonatal Tetanus	1	1.9
Acute Renal Failure	0	0.0
Multicystic Kidney Disease	1	1.9
Prematurity	5	9.4
Major Non-cardiac Congenital Anomalies	0	0.0
Neonatal Sepsis	2	3.8

Variable	Frequency	Percentage
Posterior Urethral valve	1	1.9
Severe Anaemia	1	1.9
Necrotising Enterocolitis	1	1.9
Intracranial Haemorrhage	4	7.5
Meconium Aspiration Syndrome	3	5.7
Birth Trauma	0	0.0
Meningitis	1	1.9



Figs. 1-8. Photographs showing some common causes of Neonatal death in the study
 Fig. 1 (massive subgaleal hematoma), Fig. 2 (Subarachnoid hemorrhage), Figs. 3&4 (Kernicterus),
 Fig. 5 (Aspiration Pneumonitis), Fig. 6 (Adrenal hemorrhages) and Fig. 7 (Disseminated intravascular
 coagulopathy) and Fig. 8 (Congenital Heart defects)

4. DISCUSSION

“The causes of death in the neonatal period in developing world are poorly measured, though major components are believed to be birth asphyxia, severe neonatal infections, complications of prematurity and neonatal tetanus” [8,9,11–13]. “In this study of 53 neonatal autopsies, the commonest cause of neonatal death was also severe birth asphyxia (10, 18.9%), followed by kernicterus (6, 11.3%), birth trauma (6, 11.3%), congenital heart disease (5, 9.4%), prematurity (5, 9.4%), Neonatal sepsis and Bronchopneumonia (3, 5.7%) and Neonatal Tetanus (1,1.9%) (Table 1). These is in variance to earlier study carried in Calabar by Udoma and Udo in 2001 in which they reported that the major causes of morbidity were septicaemia in 35.9 percent, neonatal jaundice in 19 percent, tetanus in 13 percent and birth asphyxia in 10.9 percent. Their findings were based on antemortem

findings rather than postmortem examination. Autopsy studies reveal that misdiagnosis is a relatively common occurrence, even where the diagnosis has been established by sophisticated modern investigative technologies” [14]. “Foetal autopsy is able to provide a definite final diagnosis in 59% (122/206) of cases and this confirms the utility of foetal autopsy in identifying the cause of foetal loss, which will help in the genetic counseling of the couple” [15]. “Asphyxia-specific early neonatal death is defined as death of foetus born at or after 35 completed weeks of gestation or weighing more than 2000g within first 7 days of life if asphyxia is assigned as primary cause of death by the attending neonatologist” [16]. “Hospital based studies in Nepal and south Africa estimated that birth asphyxia accounted for 24% and 14% of perinatal mortality respectively” [17,18]. “The incidence and mortality associated with severe birth asphyxia in Ilesha, Nigeria remained

significantly high over a period of ten years despite changes in social order. The incidence of severe birth asphyxia was 93.7/1000 admissions for the 1994 to 1998 period and 100.2/1000 admissions for the 1999 to 2003 period" [19].

"In Osogbo, Adebani et al, in 2010 reported that out of 59 babies with seizures, 37 (62.7%) were considered to have had birth asphyxia. Babies with seizures and asphyxia had the highest risk of death 59.5% and 48.8% respectively" [20]. "In Lagos, Ekure et al, in 2004 performed autopsy on 115 of the 573 perinatal deaths and reported that 38.7 percent were due to asphyxia conditions and 8.9 percent were attributed to conditions associated with immaturity" [21]. Mukhtar et al, 2007 reported that "birth asphyxia accounted for the highest number of presentations at 27%, with sepsis at 25.3% and Prematurity at 16%". While findings of neonate dying from severe birth asphyxia were mainly intense venous congestion and cyanosis with pronounced lividity, Pulmonary congestion and oedema and Fluidity of the blood. These deliveries were conducted by untrained birth attendant at home and churches with late presentation at the University of Calabar Teaching Hospital, Calabar resulting in prolonged labour. Three out of five were unbooked and two booked in the general hospital Calabar. The two booked were offered caesarean section due to postdatism and contracted pelvis but declined. Their gestational age ranged from 40weeks to 43weeeks. Two out of the five neonates delivered through spontaneous vertex delivery were found dead the next day at the labour ward with massive cephal haematoma, pulmonary haemorrhage and haemoperitoneum. Most deaths on the first day of live were the result of birth asphyxia, birth injury, or complications of preterm delivery. Birth asphyxia or injury (31%) and Preterm (26%) were the most common causes of death on day zero. On day 1 and during the first week of life, the proportion of deaths caused by birth asphyxia or injury was lower than on day 0(<14%) and on days 1-6, the frequent causes of death were preterm birth (30%) and sepsis or pneumonia (25%).

Birth trauma and kernicterus were the second commonest cause of neonatal death from our study (Figs. 3&4). The high cost of obstetric care, ignorance and unwholesome traditional belief by the uneducated and few educated women is an important factor leading to the patronage of traditional birth attendant that are mostly illiterate

and without the basic skill in obstetric care (2). The resultant effect is poor obstetric outcome with babies presenting with Asphyxia due to prolonged obstetric care, massive cephalohematoma as shown in (Fig. 1) and subarachnoid haemorrhages (Fig. 2). Although the federal ministry of health through health ministry and public health physician have taken initiative to educate and trained such high-risk women in the identification of birth challenges and the need for antenatal care with the aim of reducing neonatal mortality. The study revealed five cases of congenital heart disease, which were missed clinically misdiagnosed. None of the neonates had advanced imaging studies such as CT scan, MRI carried out to aid diagnosis due to unavailability of such investigations in our facility and cost implication. The commonest anatomical cardiac findings were large ASD, isolated large ventricular septa defect, univentricular heart chamber, pulmonary stenosis and transposition of the great arteries. The commonest isolated CHD in Nigeria is ventricular septal defect (VSD) which constitutes 27% of the cases followed by patent ductus arteriosus (PDA) 18%, atrial septal defect (ASD) (14.2%) and pulmonary stenosis (12.3%) 20 Congenital malformations of the heart are known leading causes of death in newborn was the third commonest cause of death in our study (Fig. 8).

Intracranial haemorrhage (Fig. 6) as a cause of death was clinically misdiagnosed as severe birth asphyxia (4, 100%). Autopsy findings in these categories of neonate showed diffuse subarachnoid haemorrhage, intraventricular haemorrhage into the falx cerebri and subdural haemorrhage to be the commonest type of intracranial haemorrhage seen and misdiagnosed in these centres. None of the neonates dying from intracranial haemorrhage had advanced imaging study or simple skull x-ray carried out on presentation to aid ante mortem diagnosis. These could be attributed to the length of hospital stay, time at presentation, lack of advanced imaging studies (CT scan, MRI) and the high cost of such facility.

5. CONCLUSION

The study confirms the usefulness of neonatal autopsy in ascertaining the definitive cause of death. Severe birth Asphyxia was identified as the commonest cause of death in the neonatal period followed by birth trauma, kernicterus and congenital heart diseases. Traditional birth attendant plays a major role in the increase rate

of neonatal mortality resulting from birth Asphyxia. Further training is strongly advocated for Traditional birth attendant as they appear to be the first point of care for our pregnant women.

ETHICAL APPROVAL

Ethical approval was obtained from UCTH Ethical Committee.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Buckner T, Blatt J, Smith SV. The autopsy in pediatrics and pediatric oncology: A single-institution experience. *Pediatric and Developmental Pathology*. 2006;9(5):374–80.
2. Okoroiwu HU, Uchendu KI, Essien RA. Causes of morbidity and mortality among patients admitted in a tertiary hospital in southern Nigeria: A 6 year evaluation. *PLoS One* [Internet]. 2020;15(8 August):1–15. Available from: <http://dx.doi.org/10.1371/journal.pone.0237313>
3. Charles NC, Chuku A, Anazodo NM. Childhood mortality in federal medical centre Umuahia, South Eastern Nigeria. *Oman Med J*. 2014;29(5):320–4.
4. Udo A, Ekott M, Ekanem E. Teenage Preganancy and Adverse Birth Outcomes in Calabar, Nigeria. *The Internet Journal of Gynecology and Obstetrics*. 2013;17(2):4–7.
5. NBS, UNICEF. Multiple Indicator Cluster Survey 2016-17, Final Report. Abuja, Nigeria: National Bureau of Statistics and United Nations Children's Fund; 2018.
6. Ekwochi U, Ndu IK, Nwokoye IC, Ezenwosu OU, Amadi OF, Osuorah DIC. Pattern of morbidity and mortality of newborns admitted into the sick and special care baby unit of Enugu State University Teaching Hospital, Enugu state. *Niger J Clin Pract*. 2014;17(3):346–51.
7. John KA, Olabisi FI, Olumuyiwa AA, Olawumi KA, Olubunmi BT, David BB. The Pattern and Causes of Neonatal Mortality in a Tertiary Hospital in the Southwest of Nigeria. *Journal of Kermanshah University of Medical Sciences*. 2020;24(4).
8. Akinyemi JO, Bamgboye EA, Ayeni O. Trends in neonatal mortality in Nigeria and effects of bio-demographic and maternal characteristics. *BMC Pediatr*. 2015;15(1):1–12.
9. Abolodje E. Neonatal mortality: morbidity pattern and risk factors in a resourcelimited centre in North Central Nigeria. *Sri Lanka Journal of Child Health*. 2021;50(3):408–15.
10. Undelikwo VA, Enang EE. Cultural Practices and Infant Mortality in Cross River State, Nigeria: A Sociological Perspective. *Mediterr J Soc Sci*. 2019;9(5):211–20.
11. Agan TU, Monjok E, Akpan UB, Omoronyia OE, Ekabua JE. Trend and causes of maternal mortality in a nigerian tertiary hospital: A 5-year retrospective study (2010-2014) at the university of calabar teaching hospital, Calabar, Nigeria. *Open Access Maced J Med Sci*. 2018;6(6):1153–8.
12. Thomas M, Olusoji O, Awolola N. Spectrum of congenital heart diseases in an African population: A necropsy study. *World J Cardiovasc Dis*. 2013;03(01):34–9.
13. Olufunke B, Ebenezer A, Ayodeji A, Olufunmilayo A, Olaniyi F, Adebami Olusegun. A 5-Year Audit of Neonatal Mortality at Federal Teaching Hospital, Ido-Ekiti, SouthWest Nigeria. *Niger Delta Medical Journal*. 2022;6(2):6–16.
14. FPR DV, Ruhaya M. Students' Opinions on Autopsy and Death. *South African Family Practice*. 2005;47(1):47–50.
15. Maessen M, Matten BC Van Der. Correlation Between Prenatal Test Results and Foetal Autopsy Findings. *Erasmus Journal of Medicine*. 2011;1(2):14–18.
16. Rani, Shikha Chawla, Deepak Huria, Anju and Jain S. Risk Factors for Perinatal Mortality Due to Asphyxia Among Emergency Obstetric Referrals in a Tertiary Hospital. *Indian Pediatrics*. 2010;1:1–7.
17. Lee AC, Mullany LC, Tielsch JM, Katz J, Khattry SK, LeClerg SC. Risk Factors for Neonatal Mortality due to Birth Asphyxia in Southern Nepal: A Prospective, Community-based Cohort Study. *Pediatrics*. 2008;121(5):1–18.
18. Fajolu I, Egri-Okwaji M. Childhood Mortality in Children Emergency Centre of the Lagos University Teaching Hospital.

- Nigerian Journal of Paediatrics. 2011; 38(3):131–135.
19. Adebani OJ. Pattern of neonatal Seizures in Osogbo, South-Western Nigeria. South African Journal of Child Health. 2010;4 (2):46–49.
20. Ekure E, Iroha E, Egri-Okwaji M, Ogedengbe O. Perinatal Mortality at the close of the 20th Century in Lagos University Teaching Hospital. Nigerian Journal of Paediatrics. 2004;31(1):14–18.
21. Baqui AH, Darmstadt GL, Williams EK, Kumar V, Kiran TU, Panwar D et al. Rates, Timing and Causes of Neonatal Deaths in Rural India: Implications for Neonatal Health Programmes. Bull World Health Organization. 2006;84(9): 706-13.

© 2023 Ipeh et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/92841>