



Cadmium Priming Alleviates Salinity Induced Oxidative Stress in Pigeon Pea

**Abhaya Kumar Sahu^a, Beda Saurav Behera^a
and Punam Kumari^{a*}**

^a P. G. Department of Biosciences and Biotechnology, Fakir Mohan University, Vyasa Vihar, Balasore-756089, Odisha, India.

Authors' contributions

This work was carried out in collaboration among all authors. Author AKS prepared the first draft of the manuscript. Author BSB performed the statistical analysis. Author PK designed the study, reviewed and corrected the draft of the manuscript. All authors read, incorporated their ideas, proof read the final paper and approved it for submission.

Article Information

DOI: 10.9734/AJAHR/2023/v10i4264

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/100891>

Original Research Article

Received: 19/04/2023

Accepted: 21/06/2023

Published: 09/07/2023

ABSTRACT

Plants are regularly exposed to an adverse environment due to their sessile nature which has negative impacts on plant development and productivity. In this study, the effect of seeds priming with 50 μM CdCl_2 in ameliorating the salinity (200 mM NaCl) induced oxidative stress in pigeon pea was assessed. Hydrogen peroxide (H_2O_2) and malondialdehyde (MDA) content was reduced in Cd+SS (50 μM CdCl_2 primed followed by 200 mM NaCl treatment) tissues as compared to salt stressed (SS) tissues. The activities of superoxide dismutase (SOD), ascorbate peroxidase (APX), catalase (CAT), and guaiacol peroxidase (GPX) were increased in Cd+SS tissues as compared to SS tissues. Moreover, the concentrations of ascorbic acid (ASC) and proline were also increased in Cd+SS tissues as compared to SS tissues. Thus, a low dose Cd priming provided tolerance to pigeon pea seedlings by activating the antioxidant machinery.

Keywords: *CdCl₂ priming; pigeon pea; salt stress; antioxidants; antioxidative enzymes; proline.*

*Corresponding author: Email: punam.lifescience@gmail.com;

Asian J. Agric. Hort. Res., vol. 10, no. 4, pp. 214-226, 2023

1. INTRODUCTION

Ecological stress places a significant strain on plant sustainability [1]. Salinity is amongst the most frequent abiotic stresses which trigger the plant's proliferation and developmental operations [2]. As a consequence, there is a massive yearly production loss throughout the world [3]. On a worldwide platform, extensive irrigation has enhanced the salinity of agricultural land to the point that 50% of farmland could be reduced to salinity by 2050 [4]. Salt induced farmland deterioration is indeed a big concern that has a global impact on food yields [5,6].

Plant development is negatively affected by salinity worse than by any certain toxic chemical on the planet [7], salt affects a variety of plant physiological and biochemical systems [8]. The stomatal electric density, plant water linkages, and photosynthetic pigments are all reduced during salt stress. Salinity also decreases the speed of transpiration, photosynthesis, growth, and biofuels generation [9]. The principal cause of reactive oxygen species (ROS) formation in chloroplasts is due to inadequate CO₂ uptake caused by insufficient Na⁺ buildup in plants, which leads to hyper-reduction of the electron transport complex (ETC). During stressful situations, hyper-reduction of mitochondrial ETC is another major source of ROS [10]. The primary ROS that triggers oxidative stress in plants is superoxide radical (O₂⁻), hydroxyl radical (OH[•]), singlet oxygen (¹O₂⁻), and H₂O₂ [11].

Plant protection towards enhanced ROS through adverse circumstances is linked to the retention of cellular redox equilibrium, which is primarily provided via non-enzymatic and enzymatic antioxidants like CAT, peroxidase (POD), glutathione reductase (GR), glutathione peroxidase (GSH-Px), SOD, glutathione-S-transferase (GST), monodehydroascorbate reductase (MDAR), APX, and dehydroascorbate reductase (DHAR) [2, 12, 13]. Plant manufacture osmoprotectants including sucrose, glycine betaine, mannitol, trehalose, and proline, which help to sustain water relations, stabilize enzymes, protein complexes, and membranes under saline conditions [14-18].

Pigeon pea [*Cajanus cajan* (L.) Millsp.] is the second-largest legume crop in India, behind chickpea, and is primarily planted mostly around the Kharif season. The pigeon pea has a salt sensitivity to its development. In pigeon pea,

excessive salt induces drought, and persistent and large treatment is fatal.

Seed priming is a technique for controlling the level of moisture in seeds and the metabolic activities needed for germination. Seed priming improved germination, seedling vigor, and mortality of rice seedlings during normal and severe soil and climatic circumstances [19]. Numerous reports described that seed priming employs organic and inorganic compounds such as choline, chitosan, putrescine, ethanol, paclobutrazol, zinc sulphate, potassium sulphate, copper sulphate, and selenium sulphate to promote crop plant development and tolerances against abiotic stresses [20]. Supplementing selenium (Se) at smaller doses prevents plants from ROS-induced oxidative destruction by triggering antioxidative systems [21] and lowering MDA and H₂O₂ concentrations [22]. Cucumber, rapeseed, canola, and parsley were protected against sodium chloride (NaCl) toxicity by Se priming [23]. Metal-induced defense in plants might be achievable, if a specified concentration of metal has little or no damage to the plant than to the pathogen [24].

Previous studies have demonstrated that low dose of 50 µM CdCl₂ priming prevent *Fusarium* infection in wheat [25]. The aim of the study was to unravel the role of 50 µM CdCl₂ priming in alleviation of salinity induced oxidative stress in pigeon pea.

2. MATERIALS AND METHODS

2.1 Seed Priming and NaCl Treatment

Healthy and uniform seeds of pigeon pea (*Cajanus cajan* L.) (PRG-176), were procured from the Center for Pulses Research (CPR), Berhampur, Odisha, India. The seeds were surface sterilized with 0.1% HgCl₂ solution after being rinsed with distilled water. The seeds were divided evenly; the first half was soaked with distilled water (control) and the second half with 50 µM CdCl₂ solution (metal treated). Both the halves were kept in conical flasks and mouths closed with cotton plugs and kept in dark condition at room temperature (RT) for 48 h. The seeds were then transferred into sterilized petri plates with the base covered with muslin fabric, sealed with parafilm, and covered with dark polythene, and maintained in the dark for 48 h before being incubated under light for 12 h [26].

Once they developed two leaves, these were transferred into a test tube for 10 d for further growth and development. Ten days old seedlings of both untreated control (UC) and Cd primed seeds were exposed to salt stress (200 mM NaCl) for 7 d and plants were harvested for further physio-biochemical analyses.

2.2 Chlorophyll Content

The chlorophyll was extracted from fresh leaves of each group by using 80% acetone, the total chlorophyll content was measured in a UV-VIS spectrophotometer following the method proposed by [27].

2.3 Relative Water Content (RWC)

The RWC of plant seedlings from each group was calculated using the method of [28].

$$\text{RWC (\%)} = (\text{FW} - \text{DW}) / (\text{TW} - \text{DW}) \times 100$$

2.4 Assay of Biomass

Fresh weight (FW) was recorded from freshly detached leaves and subjected to oven drying at 70°C for 24 h to record dry weight (DW). The biomass content of plant seedlings from each group was calculated using the method of [29].

$$\text{Total biomass} = [(\text{FW} - \text{DW}) / \text{FW}] \times 100$$

2.5 Assay of Oxidants

2.5.1 Estimation of H₂O₂ Content

The H₂O₂ content was assayed using the method described by [30]. 0.1 g of the shoot tissues of every sample were taken and homogenized independently in a sterile mortar and pestle with 2 ml 0.1% (w/v) TCA before centrifugation at 12,000 rpm for 15 min at 4°C. The intensity was read at 390 nm employing a UV spectrophotometer with 0.5 ml of the supernatant, 0.5 ml of 10 mM phosphate buffer with pH-7.0, and 1 ml of 1M KI in a cuvette. The H₂O₂ content was calculated by an extinction coefficient 0.28 μM⁻¹ cm⁻¹ and was quantified as μM g⁻¹ F.W.

2.5.2 Estimation of MDA Content

The MDA content was assayed using the method describe by [31]. 0.5 ml of 5%TCA was used to homogenize the shoot tissues and centrifuged at 12,000 rpm for 20 min. 2 ml TBA reagent (0.5% TBA in 20% TCA) was added to 0.5 ml supernatants for the MDA estimation. The

absorption of homogenates was evaluated at 532 nm and nonspecific turbidity was corrected by subtracting absorbance at 600 nm, with units of mM⁻¹cm⁻¹.

2.5.3 Histochemical Detection of H₂O₂

The localization of H₂O₂ was carried out histochemically in leaf samples using the method described by [32]. Healthy leaves of each experimental setup were dipped in DAB solution (1 mg ml⁻¹, pH-4.0) in a sterilized glass beaker for 12 h under light at RT. The leaves were then dipped in 95% ethanol and boiled for 20min to decolorize them and after cooling, the presence of H₂O₂ was discovered using a light microscope (brown spots).

2.6 Assay of Enzymatic Antioxidants

2.6.1 Estimation of SOD activity (EC 1.15.1.1)

The SOD activity from each sample was calculated by using the method proposed by [33]. The activity of SOD to prevent the photochemical degradation of nitro blue tetrazolium (NBT) was used to estimate SOD activity. 0.5 g of tissue from each sample was taken and was homogenized with 2ml of 50mM of phosphate buffer containing 1mM EDTA and 2% PVP and then centrifuged at 13,000 rpm for 20 min at 4°C and supernatant was collected.50mM phosphate buffer with pH-7, 0.3 ml of 130 mM methionine, 0.3 ml of 750 μM NBT, 0.3 ml of 10 mM EDTA, 0.3 ml of 20 μM riboflavin, 0.25 ml of distilled water, and 50 μl extracted sample was taken in sterilized test tubes and placed under fluorescent lamp for 10 min. After 10 min the absorbance was checked at 560 nm. The quantity of enzymes necessary to block the photoreduction activity of the NBT by 50% was denoted by one unit (U) of SOD activity. The enzyme performance was expressed as U g⁻¹ f.w. as well as the SOD activity was calculated utilizing provided equation.

$$\% \text{ of inhibition} = [1 - \text{Absorbance of each sample} / \text{Absorbance of the control}] \times 100$$

2.6.2 Estimation of APX activity (EC 1.11.1.11)

An APX activity of every sample was determined using the protocol outlined by [34]. In 2 ml of 50 mM phosphate buffer solution containing 1mM EDTA and 2% PVP, 0.5 g of tissue from each sample was homogenized properly and centrifuged at 13,000 rpm for 20 min at 4°C and

the supernatant was collected into a sterilized eppendorf. In a cuvette, 1 ml of the reaction mixture comprising 600 μ l of 50 mM phosphate buffer solution, 100 μ l of 1 mM EDTA, 100 μ l of 5 mM ascorbic acid, 100 μ l of H_2O_2 , and 100 μ l of the sample extract were taken. A UV-VIS spectrophotometer was employed to measure the intensity at 290 nm for 3 min. The ascorbate extinction coefficient of $2.8 \text{ nm}^{-1} \text{ cm}^{-1}$ was used to calculate the enzyme function by monitoring the changes in absorbance at 290 nm for 3 min. 1 unit of enzyme action ($\text{U g}^{-1} \text{ f.w.}$) was established as the amount of enzyme necessary to catalyze the oxidation of 1 mol of ascorbate in 1 min.

2.6.3 Estimation of CAT activity (EC 1.11.1.6)

The activity of CAT was determined by [35]. 0.5 g tissue was homogenized in 2 ml of 50 mM phosphate buffer solution including 1mM EDTA and 2% PVP. The extracts were centrifuged at 4°C at 13,000 rpm for 20 min. The supernatant was kept on ice for assay. The reaction mixture containing 50 μ l of 30 mM H_2O_2 , 2.9 ml of 50 mM of enzyme extract was taken in a cuvette and the change in absorbance was recorded at 240 nm for 3 min. The activity was measured using extinction coefficient 40 mM cm^{-1} and was expressed as $\text{U g}^{-1} \text{ f.w.}$

2.6.4 Estimation of GPX activity (EC 1.11.1.9)

The GPX activity was assayed by the following method of [36]. 0.5 g of tissue from each sample was taken and homogenized with 50 mM PBS buffer solution containing 1 μ M EDTA and 2% PVP. At 4°C , the supernatants were centrifuged for 20 min at 13,000 rpm. The reaction mixture contained 50 mM PBS buffer, 30 mM guaiacol, and 40 mM H_2O_2 along with enzyme extract was carried out for 30 min and the absorbance was read at 470 nm. One minute of GPX activity equals the quantity of enzyme catalyzing the oxidation of one mole of guaiacol in one minute, and enzyme action was measured in units of $\text{U g}^{-1} \text{ f.w.}$

2.7 Assay of Antioxidants

2.7.1 Estimation of ASC content

The ASC content was measured by the following method of [37]. 0.1 g stored tissue of shoots were taken and homogenized with 4% TCA in sterilized mortar and pestle. Lysates were placed in a 2 ml eppendorf tube and centrifuged for 10 min at 2,000 rpm at 4°C . Supernatants were

collected and a pinch of charcoal was added to it. The samples were kept for 5 min at RT and centrifuged for 10 min at 2,000 rpm at 4°C to remove charcoal. The aliquotes were used for estimation of ascorbate. 0.5 ml charcoal-treated supernatant was transferred to the sterilized test tube. 1.5 ml of 4% TCA, 0.5 ml of 2% dinitrophenyl hydrazine, and 2 drops of 10% thiourea solution was merged with prepared charcoal aggregate and maintained at 37°C for 3 h. Osazone crystal was formed and dissolved in 85% H_2SO_4 under cold conditions. The absorption was read at 540 nm, and measured in $\mu\text{g}^{-1} \text{ f.w.}$

2.7.2 Estimation of Proline content

The protocol of proline was used to determine the quantity by [38]. 1 ml of 3% sulfosalicylic acid was mixed with 0.1 g leaf tissue and centrifuged for 10 min at 4°C at 12,000 rpm. The solution mixture was maintained for 1 h at 100°C in a heated water bath until it became orange, using 500 mM precipitate, glacial acetic acid, and ninhydrin solution. To finish the action, the resulting solution was placed in an ice bath. 1 ml toluene was added and vigorously stirred for 15-20 sec, after which the color changed to pink. At 520 nm, the solution was read and expressed with $\mu\text{mol g}^{-1} \text{ f.w.}$

2.8 Statistical Analysis

All experiments were performed three times repeatedly and independently with standard error mean (SEM). All the biochemical data were statistically examined and Anova tested for significant (* indicates $p \leq 0.05$) difference using GraphPad Prism software.

3. RESULTS AND DISCUSSION

3.1 Morpho-physiological parameters

Plants have established certain quick methods to counteract salinity induced destruction and acclimatize to a saline environment [13]. Micromolecules like phytohormones and signaling molecules, in complement to their escape strategies, let plants readjust to unfavorable environmental circumstances. Cd pigeon pea seedlings showed improved morphological development as compared to UC seedlings. High chlorosis and severe wilting symptoms were observed in UC seedlings than that of Cd primed seedlings when exposed to salt stress (Fig. 1). The Cd primed pigeon pea

seedlings showed a higher amount of total chlorophyll content in the leaf tissues as compared to the UC tissues. The elevated

chlorophyll content was observed in Cd+SS tissues as compared to UC tissues during salt stress (Fig. 2A).




Fig. 1. Comparative assessment of phenotypes in (A) UC, (B) Cd, (C) SS and (D) Cd+SS of pigeon pea seedlings (UC: Untreated control, Cd: 50 µM CdCl₂ pre-treated, SS: Salt-stressed, and Cd+SS: 50 µM CdCl₂ primed followed by 200 mM NaCl treatment)




Fig. 2 (A-C): Physiological analysis (A) Total chlorophyll content, (B) RWC (%), and (C) Biomass (g) of shoot tissues of pigeon pea seedlings (UC: Untreated control, Cd: 50 µM CdCl₂ pre-treated, SS: Salt-stressed, and Cd+SS: 50 µM CdCl₂ primed followed by 200 mM NaCl treatment)

Salinity is amongst the most serious ecological stresses for plants, causing a reduction in growth rate as well as a variety of metabolic disturbances. Even a modest amount of salt can disrupt normal physiological and biochemical pathways, causing the cell to slow down its growth and progression [8, 39]. Plant proliferation is regulated by photosynthetic pigments and photosynthesis rate [13]. The photosynthetic pigments like total chlorophyll content was reduced dramatically in UC seedlings when exposed to salt stress similarly reported in rice [40] and *Pisum sativum* [41]. The aggregation of Na^+ and Cl^- ions enhances with larger salt concentrations, that hampered chlorophyll production by interfering with the Fe^{3+} -containing chlorophyll synthesizing enzymes [42], enhance the levels of the chlorophyll-degrading enzyme chlorophyllase and reactive oxygen species (ROS) [22]. In Cd+SS pigeon pea seedlings, priming with Cd enhanced total chlorophyll content. It was previously reported that Cd^{2+} priming increased total chlorophyll content in *Trigonella foenum-graecum* [43], *Festuca arundinacea* [44], and rice [45].

The RWC in Cd primed pigeon pea seedlings was higher than the UC seedlings. The Cd+SS pigeon pea seedlings showed a significantly high RWC value as compared to the SS seedlings (Fig. 2B). It occurred as a result of osmotic stress, which causes the root to become hard and unable to absorb water [2, 8]. This result was analogous to exogenous Ca^{2+} priming favorably

influencing stomatal performance by maintaining guard cells turgid [46], guaranteeing CO_2 supply, or regulating stomatal conductance [47]. Turgor pressure is essential for cell expansion, however, salinity diminishes turgor pressure, resulting in shorter shoot and root lengths and inhibited growth [48].

The enhanced biomass was observed in UC seedlings as compared to the seedlings under salinity stress. When compared to Cd+SS, SS seedlings had lower biomass (Fig. 2C). [49] reported reduced shoot and root development, and physiological abnormalities (photosynthetic pigment degradation and electron flow impairment) leads to loss of biomass deposition. In our investigation, we found that the biomass was decreases when seedlings were subjected to salt stress. As salinity promotes abscisic acid (ABA) mediated stomatal closure, which slows CO_2 assimilation and disrupts the Calvin cycle's normal electron flow for carbon breakdown, fresh and dry matter synthesis was diminished [50, 51].

3.2 Oxidants

3.2.1 H_2O_2 content

The SS seedlings had a significantly larger H_2O_2 content than the UC, Cd primed and Cd+SS tissues. In contrast to SS tissues, the H_2O_2 content in Cd+SS tissues was lower (Fig. 3).




Fig. 3. H_2O_2 content of shoot tissues of pigeon pea seedlings (UC: Untreated control, Cd: $50 \mu\text{M CdCl}_2$ pre-treated, SS: Salt-stressed, and Cd+SS: $50 \mu\text{M CdCl}_2$ primed followed by 200 mM NaCl treatment)

Salt induces the development of ROS such as H_2O_2 , O_2^- , and OH^\cdot in plants [52]. ROS production could potentially be caused by an abundance of Na^+ in the cells and a lack of Ca^{2+} and K^+ [53]. As a result, the H_2O_2 content in SS tissues was significantly higher than in Cd+SS, Cd primed, and UC tissues in the current investigation, which was consistent with previous results. It had also been reported that H_2O_2 would be the most stable ROS and its excessive generation and aggregation is poisonous and harmful to plant cells during abiotic stress [54] and severely disrupt normal metabolism by peroxidation of membrane lipid [55].

3.2.2 MDA content

The MDA content in the SS tissues was high than the Cd+SS tissues. However, the MDA content was observed minimum in UC and Cd primed tissues (Fig. 4).

Moreover, increased ROS production in plant tissues causes cell damage and poses a serious threat to macromolecules like DNA, proteins, and lipids, leading to lipid peroxidation, protein oxidation, and DNA damage [56]. Moreover, when comparing to UC tissues exposed to salt, a limiting degree of lipid peroxidation was detected in Cd+SS tissues, which leads to less amount of synthesis of MDA as a byproduct of lipid peroxidation [55, 56].

3.2.3 Histochemical analysis of H_2O_2

Salt stress causes a high amount of ROS that induces tissue damage in the plant. The localized H_2O_2 in the pigeon pea leaves was observed by staining the leaves through oxidation of DAB. SS pigeon pea leaves showed higher numbers of dark brown spots which were indicative of accumulated H_2O_2 as compared to the Cd+SS leaves [57] (Fig. 5).




Fig. 4. MDA content of shoot tissues of pigeon pea seedlings (UC: Untreated control, Cd: 50 $\mu\text{M CdCl}_2$ pre-treated, SS: Salt-stressed, and Cd+SS: 50 $\mu\text{M CdCl}_2$ primed followed by 200 mM NaCl treatment)




Fig. 5. Histochemical localization of H_2O_2 content (A) UC, (B) Cd, (C) SS, and (D) Cd+SS leaves of pigeon pea seedlings (UC: Untreated control, Cd: 50 $\mu\text{M CdCl}_2$ pre-treated, SS: Salt-stressed, and Cd+SS: 50 $\mu\text{M CdCl}_2$ primed followed by 200 mM NaCl treatment)

3.3 Assay of Antioxidant Enzymes

The SOD activity was detected higher in the Cd+SS tissues as compared to SS tissues. Similar results found in rice under water deficit condition [58]. Moreover, the Cd+SS tissues showed high SOD activity as compared to the UC and Cd primed tissues (Fig. 6A).

Salinity stress has a big impact on SOD behavior [2]. Under high metal stress, rice varieties showed an increase in SOD activity in the leaves [59]. It could be related to an elevation in O_2^- production, which results in enhanced SOD gene expression [60]. The very first enzyme in the detoxifying process, SOD converts O_2^- radicals to H_2O_2 [61]. The Cd+SS tissues also showed high SOD activity but the Cd^{2+} priming might be ameliorated the H_2O_2 production and it was previously confirmed by [62] on soyabean nodules.

When plants are exposed to salt stress, ROS formation arises [13]. Throughout this condition, antioxidant enzymes are intended to involve a vital part in the plant's protection against ROS [63]. APX activity was lower in the SS tissues than Cd+SS tissues (Fig. 6B) which was previously confirmed by [64] on *Glycine max*. The ascorbate-glutathione (ASC-GSH) cycle is a crucial H_2O_2 detoxifying pathway in plant cells, and APX enzymes play a crucial role in

transforming H_2O_2 to H_2O employing ASC as a primary electron donor [65].

The SS tissues had lower CAT activity than the Cd+SS tissues (Fig. 6C). Rice [66] and *Vigna radiata* had similar results [67]. To eliminate H_2O_2 from plants, CAT converts H_2O_2 to oxygen and water. Hence, the higher CAT activity contributed to better salt tolerance [11].

The Cd+SS tissues showed high GPX activity as compared to SS tissues (Fig. 6D), as also reported by [68] in the roots and leaves of *Oryza sativa*. Our findings showed that increased levels of GPX in seedlings successfully absorb H_2O_2 , which is a key modulator of cellular damage under environmental stress.

3.4 Assay of Antioxidants

ASC is an antioxidant and a critical substrate for the detoxifying of ROS [69]. Through non-enzymatic and enzymatic detoxification pathways, ASC is one of the most effective antioxidants generated from plants, and it performs a vital role in the lowering of excessive ROS [70]. Whenever plants are stressed by drought, salinity, or even other environmental challenges, the non-enzymatic antioxidants ASC and proline accumulate to high levels, according to several reports [71]. ASC concentration was found to be lower in SS tissues when compared to Cd+SS in this study (Fig. 7A).




Fig. 6. (A-D). Antioxidant activity in shoot tissues of pigeon pea seedlings. (A) SOD activity, (B) APX activity, (C) CAT activity, and (D) GPX activity (UC: Untreated control, Cd: 50 μ M $CdCl_2$ pre-treated, SS: Salt-stressed, and Cd+SS: 50 μ M $CdCl_2$ primed followed by 200 mM NaCl treatment)




Fig. 7. (A-B): Antioxidant contents in the shoot tissues of pigeon pea seedlings. (A) ASC content, and (B) Proline content UC: Untreated control, Cd: 50 µM CdCl₂ pre-treated, SS: Salt-stressed, and Cd+SS: 50 µM CdCl₂ primed followed by 200 mM NaCl treatment)

A low dose of Cd could detoxify the ROS content in Cd+SS tissues by accumulating ascorbate content, which might detoxify the ROS to stabilize the redox status. Certain metabolic processes also metabolize MDA and H₂O₂, resulting in the regeneration of reduced ASC in the ASC–GSH cycle [72].

The formation of osmolytes is a normal approach to stress in an attempt to minimize physiological disruption [73]. The increased proline content was observed in Cd+SS tissues as compared to SS tissues (Fig. 7B). The proline content was maximum in Cd+SS tissues which indicates that antioxidants might be acting as ROS detoxifiers at the cellular level and protects the integrity of membranes, stability of enzymes or proteins, and provides tolerance to diverse stimuli [41, 74].

4. CONCLUSION

It was concluded that a low dose CdCl₂ (50µM) pre-treatment conferred tolerance to salinity induced oxidative stress in pigeon pea seedlings through the activation of antioxidants defense machinery.

ACKNOWLEDGMENTS

The authors highly acknowledge the P. G. Department of Biosciences and Biotechnology, Fakir Mohan University, Balasore, Odisha, India for the necessary and laboratory facilities to carry out the study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Mahmood-ur-Rahman Ijaz M, Qamar S, Bukhari SA, Malik K. Abiotic stress signaling in rice crop. In: Hasanuzzaman M (ed) Advances in rice research for abiotic stress tolerance. Woodhead Publishing, Cambridge. 2019;551–569.
2. Safdar H, Amin A, Shafiq Y, Ali A, Yasin R, Shoukat A, Hussan MU, Sarwar MI. A review: impact of salinity on plant growth. Nature and Science.2019;17:34–40.
3. Munns R, Tester M. Mechanisms of salinity tolerance. Annual Review of Plant Biology.2008;59:651–681.
4. Wang W, Vinocur B, Altman A. Plant responses to drought, salinity and extreme temperatures: towards genetic engineering for stress tolerance. Planta. 2003;218(1):1–14.
5. Geist H. The causes and progression of desertification. Routledge, London; 2017.
6. FAO FAOSTAT: online statistical database; 2016.
7. Xiong L, Zhu J. Molecular and genetic aspects of plant responses to osmotic stress. Plant Cell and Environment. 2002;25(2):131–139.
8. Goyal MR, Gupta SK, Singh A. Physiological and biochemical changes in plants under soil salinity stress: a review. In: Gupta SK et al (eds) Engineering practices for management of soil salinity, 1st edn. Apple Academic Press, Palm Bay. 2018;159–200.
9. Morton MJ, Awlia M, Al-Tamimi N, Saade S, Pailles Y, Negrao S, Tester M. Salt stress under the scalpel–dissecting the

- genetics of salt tolerance. *Plant Journal*. 2019;97(1):148–163.
10. Saini P, Gani M, Kaur JJ, Godara LC, Singh C, Chauhan SS, Francies RM, Bhardwaj A, Kumar NB, Ghosh M K. Reactive oxygen species (ROS): a way to stress survival in plants. In: Zargar SM, Zargar MY (eds) *Abiotic stress-mediated sensing and signaling in plants: an omics perspective*. Springer, Singapore. 2018;127–153.
 11. Waszczak C, Carmody M, Kangasja“rvi J. Reactive oxygen species in plant signaling. *Annual Review Plant Biology*. 2018;69:209–236.
 12. Yan K, Shao H, Shao C, Chen P, Zhao S, Brestic M, Chen X. Physiological adaptive mechanisms of plants grown in saline soil and implications for sustainable saline agriculture in coastal zone. *Acta Physiologiae Plantarum*. 2013;35:2867–2878.
 13. Tang X, Mu X, Shao H, Wang H, Brestic M. Global plant responding mechanisms to salt stress: physiological and molecular levels and implications in biotechnology. *Critical Review Biotechnology*. 2015;35(4):425–437.
 14. Slama I, Abdelly C, Bouchereau A, Flowers T, Savoure A. Diversity, distribution and roles of osmoprotective compounds accumulated in halophytes under abiotic stress. *Annals of Botany*. 2015;115(3):433–447.
 15. Iqbal M, Hussain I, Liaqat H, Ashraf MA, Rasheed R, Rehman AU. Exogenously applied selenium reduces oxidative stress and induces heat tolerance in spring wheat. *Plant Physiology & Biochemistry*. 2015;94:95–103.
 16. Sami F, Yusuf M, Faizan M, Faraz A, Hayat S. Role of sugars under abiotic stress. *Plant Physiology and Biochemistry*. 2016;109:54–61.
 17. Kurepin LV, Ivanov AG, Zaman M, Pharis RP, Hurry V, Huner NP. Interaction of Glycine betaine and plant hormones: protection of the photosynthetic apparatus during abiotic stress. In: Hou HJM et al (eds) *Photosynthesis: structures, mechanisms, and applications*. Springer, Cham. 2017;185–202.
 18. Patel TK, Williamson JD. Mannitol in plants, fungi, and plant– fungal interactions. *Trends in Plant Science*. 2016;21(6):486–497.
 19. Hussain S, Khan F, Hussain HA, Nie L. Physiological and biochemical mechanisms of seed priming-induced chilling tolerance in rice cultivars. *Frontier in Plant Science*. 2016;7:116.
 20. Jisha KC, Vijayakumari K, Puthur JT. Seed priming for abiotic stress tolerance: an overview. *Acta Physiologiae Plantarum*. 2013;35(5):1381–1396.
 21. Iqbal N, Umar S, Khan NA. Nitrogen availability regulates proline and ethylene production and alleviates salinity stress in mustard (*Brassica juncea*). *Journal of Plant Physiology*. 2015;178:84–91
 22. Hasanuzzaman M, Alam MM, Rahman A, Hasanuzzaman M, Nahar K, Fujita M. Exogenous proline and glycine betaine mediated upregulation of antioxidant defense and glyoxalase systems provides better protection against salt-induced oxidative stress in two rice (*Oryza sativa* L.) varieties. *Biomed Research International*; 2014.
 23. Habibi G. Selenium ameliorates salinity stress in *Petroselinum crispum* by modulation of photosynthesis and by reducing shoot Na accumulation. *Russian Journal Plant Physiology*. 2017;64:368–374.
 24. Poschenrieder C, Tolra R, Barcelo J. Can metals defend plants against biotic stress? *Trends in Plant Science*. 2006;11(6):287–295.
 25. Mitra B, Ghosh P, Henry SL, Mishra J, Das TK, Ghosh S, Babu CR, Mohanty P. Novel mode of resistance to *Fusarium* infection by mild dose pre-exposure of cadmium to wheat. *Plant Physiology and Biochemistry*. 2004;42:781–787.
 26. Tahjib-UI-Arif M, Roy PR, Sohag AAM, Afrin S, Rady MM, Hossain MA. Exogenous calcium supplementation improves salinity tolerance in BRR1 dhan28; a salt-susceptible high yielding *Oryza sativa* cultivar. *Journal of Crop Science & Biotechnology*. 2018a;21(4): 383–394.
 27. Arnon DI. Copper enzymes in isolated chloroplasts: polyphenol oxidase in *Beta vulgaris*. *Plant Physiology*. 1949;24(1):1–15.
 28. Tahjib-UI-Arif M, Sayed MA, Islam MM, Siddiqui MN, Begum SN, Hossain MA. Screening of rice landraces (*Oryza sativa* L.) for seedling stage salinity tolerance using morpho-physiological and molecular

- markers. *Acta Physiologiae Plantarum*. 2018b;40(4):70.
29. Kumari P, Mahapatro GK, Banerjee N, Sarin NB. Ectopic expression of GroEL from *Xenorhabdus nematophila* in tomato enhances resistance against *Helicoverpa armigera* and salt and thermal stress. *Transgenic Research*. 2015;24(5):859-73.
 30. Noreen Z, Ashraf M. Change in antioxidant enzymes and some key metabolites in some genetically diverse cultivars of radish (*Raphanus sativus* L.). *Environmental and Experimental Botany*. 2009;67(2):395-402.
 31. Basu S, Roychoudhury A, Saha PP. Differential antioxidative responses of indica rice cultivars to drought stress. *Plant Growth Regulation*. 2010;60(51).
 32. Daudi A, O'Brien J. Detection of Hydrogen Peroxide by DAB Staining in *Arabidopsis* Leaves. *Bio-protocol*. 2012;2.
 33. Beyer WF, Fridovich I. Assaying for superoxide dismutase activity: some large consequences of minor changes in conditions. *Analytical Biochemistry*. 1987;161(2):559-66.
 34. Nakano Y, Asada K. Hydrogen peroxide scavenging by ascorbate-specific peroxidase in spinach chloroplast. *Plant and Cell Physiology*. 1981;22:867-880.
 35. Aebi H. Catalase in vitro, *Methods in Enzymology* Academic Press. 1984;105:121-126.
 36. Weydert CJ, Cullen JJ. Measurement of superoxide dismutase, catalase and glutathione peroxidase in cultured cells and tissue. *Nature protocols*. 2010;5(1):51-66.
 37. Gillespie KM, Ainsworth EA. Measurement of reduced, oxidized and total ascorbate content in plants. *Nature Protocols*. 2007;2(4):871-87.
 38. Rajasubramaniyam K, Hanumappa KR, Harish R, Narendra K. Estimation of amino acid, protein, proline and sugars in *Vigna mungo* L. exposed to cement dust pollution. *International Journal of Advanced Research in Biological Sciences*. 2018;5(7):173-186.
 39. Parihar P, Singh S, Singh R, Singh VP, Prasad SM. Effect of salinity stress on plants and its tolerance strategies: a review. *Environmental Science and Pollution Research*. 2015;22(6):4056-4075.
 40. Zhen-hua ZH, Qiang LI, Hai-xing SO, Xiang-min RO, Ismail AM. Responses of different rice (*Oryza sativa* L.) genotypes to salt stress and relation to carbohydrate metabolism and chlorophyll content. *African Journal of Agricultural Research*. 2012;7:19-27.
 41. Ozturk L, Demir Y, Unlukara A, Karatas I, Kurunc A, Duzdemir O. Effects of long-term salt stress on antioxidant system, chlorophyll and proline contents in pea leaves. *Romanian Biotechnological Letters*. 2012;17(3):7227-7236.
 42. Silva EN, Ribeiro RV, Ferreira-Silva SL, Vie'gas RA, Silveira JA. Salt stress induced damages on the photosynthesis of physic nut young plants. *Science Agriculture*. 2011;68(1):62-68.
 43. Oprica L, Sandu L. Impact of inorganic salt solution on antioxidative enzyme activity and photosynthetic pigments content in *Trogonella foenum-graecum* seedlings. *Analele Stiintifice ale logie Moleculara*. 2014;15(2):31-40.
 44. Wang G, Bi A, Amombo E, Li H, Zhang L, Cheng C, Hu T, Fu J. Exogenous calcium enhances the photosystem II photochemistry response in salt stressed tall fescue. *Frontier in Plant Science*. 2017;8:20-32.
 45. Tahjib-Ul-Arif M, Siddiqui MN, Sohag AAM, Sakil MA, Rahman MM, Polash MA S, Mostofa MG, Tran LSP. Salicylic acid-mediated enhancement of photosynthesis attributes and antioxidant capacity contributes to yield improvement of maize plants under salt stress. *Journal of Plant Growth Regulation*. 2018c;37:1318-1330.
 46. Agurla S, Gahir S, Munemasa S, Murata Y, Raghavendra AS. Mechanism of stomatal closure in plants exposed to drought and cold stress. In: Iwaya-Inoue M (ed) *Survival strategies in extreme cold and desiccation*. Springer, Singapore. 2018;215-232.
 47. Sibole JV, Cabot C, Poschenrieder C, Barcelo' J. Efficient leaf ion partitioning, an overriding condition for abscisic acid controlled stomatal and leaf growth responses to NaCl salinization in two legumes. *Journal of Experimental Botany*. 2003;54(390):2111-2119.
 48. Guérin A, Gravelle S, Dumais J. Forces behind plant cell division. *PNAS*. 2016;113(32):8891-8893.
 49. Fahad S, Bajwa AA, Nazir U, Anjum SA, Farooq A, Zohaib A, Sadia S, Nasim W, Adkins S, Saud S, Ihsan MZ, Alharby H, Wu C, Wang D, Huang J. Crop Production under Drought and Heat Stress: Plant

- Responses and Management Options. *Frontiers in plant science*. 2017;8:1147.
50. Mbarki S, Cerda` A, Zivcak M, Brestic M, Rabhi M, Mezni M, Jedidi N, Abdelly C, Pascual JA. Alfalfa crops amended with MSW compost can compensate the effect of salty water irrigation depending on the soil texture. *Process Safety and Environment Protection*. 2017;115: 8–16.
 51. Mbarki S, Sytar O, Cerda A, Zivcak M, Rastogi A, He X, Zoghlami A, Abdelly C, Brestic M. Strategies to mitigate the salt stress effects on photosynthetic apparatus and productivity of crop plants. In: Kumar V (ed) *Salinity responses and tolerance in plants*, vol 1. Springer, Cham. 2018;85–136.
 52. Yen WJ, Chyau CC, Lee C P, Chu HL, Chang LW, Duh PD. Cytoprotective effect of white tea against H₂O₂-induced oxidative stress in vitro. *Food Chemistry*. 2013;141:4107–4114.
 53. Demidchik V, Maathuis F J. Physiological roles of nonselective cation channels in plants: from salt stress to signaling and development. *New Phytology*. 2007;175(3):387–404.
 54. Asada K, Takahashi M. Production and Scavenging of Active Oxygen in Chloroplasts. *Photoinhibition*, Elsevier, Amsterdam. 1987;227-287.
 55. Fridovich I. Biological effects of the superoxide radical. *Archives of Biochemistry and Biophysics*. 1986;247(1):1-11.
 56. Sharma P, Jha AB, Dubey RS, Pessarakli M. Reactive oxygen species, oxidative damage, and antioxidative defense mechanism in plants under stressful conditions. *Journal of Botany*. 2012;1-26.
 57. Chao YY, Chen CY, Huang WD, Kao CH. Salicylic acid-mediated hydrogen peroxide accumulation and protection against Cd toxicity in rice leaves. *Plant and Soil*. 2010;329(1-2):327-337.
 58. Andrade FR, da Silva GN, Guimaraes KC, Barreto HBF, de Souza KRD, Guilherme LRG , Faquin V, Reis AR. Selenium protects rice plants from water deficit stress. *Ecotoxicology and Environmental Safety*. 2018;164:562–570.
 59. Verma S, Dubey RS. Lead toxicity induces lipid peroxidation and alters the activities of antioxidant enzymes in growing rice plants. *Plant Sci*. 2003;164: 645–655.
 60. De Vos CHR, Schat H. Free radicals and heavy metal tolerance, *Ecological Responses to Environmental Stresses*. Springer Netherlands UK. 1991;22–31.
 61. Hendry GAF, Baker AJM , Ewart C F. Cadmium tolerance and toxicity, oxygen radical processes and molecular damage in cadmium tolerant and cadmium sensitive clones of *Holcus lanatus* L. *Plant Biolog*. 1992;41(3):271–281.
 62. Balestrasse KB, Gardey L, Gallego SM, Tomaro ML. Response of antioxidant defence system in soybean nodules and roots subjected to cadmium stress. *Australian Journal of Plant Physiology*. 2001; 28:497–504.
 63. Hanin M, Ebel C, Ngom M, Laplaze L, Masmoudi K. New insights on plant salt tolerance mechanisms and their potential use for breeding. *Frontier in Plant Science*. 2016;7:1-17.
 64. Weisany W, Sohrabi Y, Heidari G, Siosemardeh A, Ghassemi- Golezani K. Changes in antioxidant enzymes activity and plant performance by salinity stress and zinc application in soybean (*Glycine max* L.). *Plant Omics*. 2012;5(2):60.
 65. Hossain MA, Fujita M. Hydrogen peroxide priming stimulates drought tolerance in mustard (*Brassica juncea* L.) seedlings. *Plant Gene and Trait*. 2013;4:109–123.
 66. Nounjan N, Nghia PT, Theerakulpisut P. Exogenous proline and trehalose promote recovery of rice seedlings from salt-stress and differentially modulate antioxidant enzymes and expression of related genes. *Journal of Plant Physiology*. 2012;169(6):596–604.
 67. Manivannan P, Jaleel CA, Kishorekumar A, Sankar B, Somasundaram R, Sridharan R, Panneerselvam R. Changes in antioxidant metabolism of *Vigna unguiculata* (L.) Walp. By propiconazole under water deficit stress. *Colloids Surf B Biointerfaces*. 2007;57(1): 69–74.
 68. Hayat S, Ali B, Hasan SA, Ahmad A. Brassinosteroid enhanced the level of antioxidants under cadmium stress in *Brassica juncea*. *Environmental Experimental Botany*. 2007;60(1):33–41.
 69. Qian HF, Peng XF, Han X, Ren J, Zhan KY, Zhu M. The stress factor, exogenous ascorbic acid, affects plant growth and the antioxidant system in *Arabidopsis thaliana*. *Russian Journal of Plant Physiology*. 2014;61:467–475.
 70. Veljović-Jovanović S, Vidović M, Morina F. Ascorbate as a key player in plant abiotic stress response and tolerance. In: Hossain

- MA, MunneBosch S, Burritt DJ, Dian-Vivancos P, Fujita M, Lorence A (eds) Ascorbic acid in plant growth, development and stress tolerance. Springer, Chemistry. 2017;47:109.
71. Ashraf M, Foolad MR. Roles of glycine betaine and proline in improving plant abiotic stress resistance. *Environmental and Experimental Botany*. 2007;59(2):206-216.
72. Hasanuzzaman M, Borhannuddin B, Zulfiqar F, Raza A, Mohsin SM, Mahmud JA, Fujita M, Fotopoulos V. Reactive oxygen species and antioxidant defense in plants under abiotic stress: revisiting the crucial role of a universal defense regulator. *Antioxidants*. 2020;9(8):11-52.
73. Ghosh B, Md NA, Gantait S. Response of rice under salinity stress: a review update. *Rice research: Open Access*. 2016;4(2):1-8.
74. Acosta-Motos JR, Diaz-Vivancos P, Acosta M, Hernandez JA. Effect of biostimulants on plant responses to salt stress. In: Hasanuzzaman M et al (eds) *Plant tolerance to environmental stress: role of phytoprotectants*. CRC Press, Boca Raton. 2019;363-380.

© 2023 Sahu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/100891>