



# Influence of Tank Water Depths on Reproductive Performance of Catfish (*Heterobranchus longifilis*) Valenciennes, 1840

Otoh A. J. <sup>a\*</sup>

<sup>a</sup> Department of Fisheries and Aquaculture, Faculty of Agriculture, Akwa Ibom State University, Obio, Akpa Campus, Oruk Anam, Nigeria.

## Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ajfar/2024/v26i6774>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/117628>

Original Research Article

Received: 16/03/2024

Accepted: 21/05/2024

Published: 10/06/2024

## ABSTRACT

**Aims:** To determine the optimum water depth required for catfish spawning in hatcheries.

**Study Design:** Treatments were assigned using complete randomized design.

**Place and Duration of Study:** Fish Farm Complex of the Akwa Ibom State University (AKSU), Nigeria.

**Methodology:** Eighteen (18) similarly-sized *H. longifilis* broodstock (2.51-2.53 kg, 64 – 66 cm): comprising six broodstock of three males and three females, were held for six months at 0.50 m, 0.75 m and 1.0 m pond water depth. All males were sacrificed for milt extraction without hormonal inducement; with milt from each treatment diluted with normal saline solution. Female broodstock from each treatment were separately induced with ovaprim hormones at a single dosage of 0.5ml/kg

\*Corresponding author: Email: [drotahaj@gmail.com](mailto:drotahaj@gmail.com);

**Cite as:** A. J., Otoh. 2024. "Influence of Tank Water Depths on Reproductive Performance of Catfish (*Heterobranchus Longifilis*) Valenciennes, 1840". *Asian Journal of Fisheries and Aquatic Research* 26 (6):36-42. <https://doi.org/10.9734/ajfar/2024/v26i6774>.

body weight of fish and allowed for 16 hours before manual stripping; 3g of egg from each broodstock and mixed with the diluted milt and activated with 100 ml of normal saline. The fertilized eggs were incubated and thereafter assessed for percentage fertilization, hatchability, survival and fry production success.

**Results:** broodstock reproductive parameters significantly increased ( $P = 0.00$ ) with water depth of broodstock culture tank. Percentage fertilizations were:  $49.50 \pm 0.78$ ,  $68.17 \pm 0.93$ , and  $82.50 \pm 1.44$ ; respectively. Percentage hatchability of broodstock eggs were:  $91.37 \pm 1.65$  significantly higher for 1.0 m water depth group than  $67.79 \pm 4.58$  from 0.75 m water depth treatment while the least value of hatchability  $50.31 \pm 0.78$  was recorded for broodstock raised at 0.50 m water depth. The fish group raised at 1.0 m water depth also exhibited highest percentage survival value of  $97.02 \pm 1.09$  and fry production success value of  $73.08 \pm 0.53$  while broodstock raised at 0.50 m water depth indicated the least values: % survival,  $8.38 \pm 0.48$  and fry production success of  $2.07 \pm 0.07$ .

**Conclusion:** In times or places of water scarcity, *H. longifilis* broodstock may be raised at pond water of 0.5 m to 1.0 m. But the water depth that would afford the best breeding and reproductive performance is at 1.0 m depth.

**Keywords:** Artificial fertilization; breeding; fish stress; hatchery production; fish culture; African catfish.

## 1. INTRODUCTION

Insufficient supply of water at different locality of the world, initiated the idea of water management for various agricultural purposes; this consequently influences the level or depth of pond water for fish culture [1]. Water is a natural component indispensable for all living organisms and existing as colourless, tasteless and odourless liquid. It is the natural habitat for aquatic organisms including fish – highly exploited for food by man. Unfortunately, the huge demand of animal protein for human sustenance re-enforces overfishing of wild catches [2] which drastically reduces the supply of fish from capture fisheries [3]; with aquaculture emerging as the only alternative to attaining world demand for fish [4,5].

In Nigeria, African catfish belonging to the family Clariidae (*Heterobranchus* and *Clarias* species) are the most cultivable species of significance [6,7,8]. This is due to the unique characteristics of the species such as fast growth rate, good taste, high stocking density, high market price and high resistance to disease and ability to reproduce in captivity, which makes it economical to culture [2,9,10,11,12,13]. However, the growth of fish depends on availability of good feed of which a single feed stuff component cannot achieve [4,5,14]. *Heterobranchus* and *Clarias*, readily accept supplementary feed and grow faster within a short period of culture compared to other species [1,7,15]. These species dominate fresh water

environments such as lakes, rivers and streams [16].

Sexual maturity of African catfish varies. *Clarias* species mature between 5 to 9 months while *Heterobranchus* species take a longer period of 18 months to attend sexual maturity in the case of females while males attend maturity as from 12 months [17,18,19]. Shallow water tolerance and broodstock maintenance is very crucial mostly in terms of good quality feeds, water parameters and culture environment to achieve the best gametes and success at the hatchery levels. The hopes of the fish farmers lie in adequate care of broodstock to achieve production of fish seeds. Hence, the need to determine the water depth or level for maintenance of broodstock for the best reproductive performances of African catfish is critical where water is in inadequate supply.

Several investigations had been carried out on the reproductive performances of African catfish based on hormonal induction [3,8,15], age of broodstock [6,7,19,20,21], egg batches [22,23], sperm diluents [2,24], influence of feed types [23,25], F1 offspring of [2,21,25], and growth of hybrids [26,27].

This study models the conditioning of *H. longifilis* broodstock at shallow and deeper ponds in hatchery. The aim being to determine the most suitable pond water depth for conditioning *H. longifilis* broodstock cultured at different water depths by examining their reproductive performances.

## 2. MATERIALS AND METHODS

### 2.1 Location of Study

The research was conducted at the Fish Farm Complex of Akwa Ibom State University, Obio Akpa campus in southeast Nigeria. The farm is between latitude 5°17'N and 7°27'N, Longitude 7°27'E and 7°58'E with an annual rainfall ranging from 3500mm-5000mm and average monthly temperature of 25° C. Akwa Ibom State is a coastal state lying between latitude 4°28'N and 5°3'N and between longitude 7°27'E and 8°20'E with a relative humidity between 60-70%. It is in the tropical rain forest zone of Nigeria [6,20].

### 2.2 Acquisition and Care of Broodstock

Eighteen (18) similarly-sized *H. longifilis* broodstock (2.51-2.53 kg, 64 – 66 cm) were carefully selected from the farm stock for this study. Groups of six broodstocks (three males and three females) were stocked in three separate concrete ponds, 1 x 3 x 1m<sup>3</sup> at water depth: 0.5 m, 0.75 m and 1.0 m; at the rate of 2 fishes m<sup>-2</sup>. Broodstock were fed twice daily at 5% body weight with Coppens commercial feed for six months.

### 2.3 Egg Stripping and Fertilization

The ratio of 1 male:1 female is taken for this experiment because both are of equal weights [1,2,4]. Approximately 3g eggs (2000 oocytes) from each broodstock was stripped and mixed with the diluted milt and activated with 100 ml of normal saline in nine separate containers: comprising three replicates per treatment [1,2,4].

### 2.4 Incubation

The fertilized eggs were spread uniformly in a monolayer on kakaban (shredded nylon sack) and incubated in aerated indoor concrete tanks (2 m x 1 m x 0.5 m at water depth of 10 cm) in three replicates until hatching. The incubation system was maintained within the range of 26 - 27°C, 6.5 - 7.0 pH and 4.5 - 5.0 mg L<sup>-1</sup> dissolved oxygen.

### 2.5 Hatching

The incubated eggs hatched between 26 - 27 hours after incubation at 26°C. Dead eggs were

siphoned from the breeding tanks 35 hours after incubation. Egg fertilization hatching and survival rates were determined as follows:

Fertilization rate (%) of eggs = (No. of eggs fertilized / total number of eggs) x 100

Hatchability (%) = (number of hatchlings / total number of fertilized eggs) x 100.

Percentage hatchability was obtained by direct counting of unhatched eggs as well as the numbers of eggs hatched in each incubating tank.

Hatching rate = (No of healthy fertilized eggs / No of fertilized eggs used) x 10 [24]

Survival rate (Ks) = (number of live hatchling / total number of hatchlings) x 100

during initial feeding [24]

Efficiency of hatching was evaluated as Fs (%) = Kf.Kh.Ks/10,000 where

Fs = Success rate (%) of fry production at 10-day post hatching.

Kf = Fertilization rate (%) of eggs

Kh = Hatching rate (%) of fry

Ks= survival rate (%) of 10-day-old swim-up fry [24]

### 2.6 Determination of Water Quality Parameters

pH (Vivosun pH meter) and dissolve oxygen (Extech 407510) and temperature (mercury in glass thermometer) readings were taken during the study.

### 2.7 Statistical Analysis

Statistical analyses of fertilization, hatching, survival rates at 10-day post hatching and fry production success were carried out using one-way ANOVA; the significant differences in mean were set at 0.05 significant level.




Fig. 1. Map showing the location of the Akwa Ibom State University fish farm complex

### 3. RESULTS AND DISCUSSION

#### 3.1 Physicochemical Parameters of Culture Media

The water parameters considered in this study (Table 1), showed no significant difference ( $P = .999$ ) among the treatments. Dissolved oxygen, temperature and pH measurements ranged between  $5.21 \pm 0.20 - 5.66 \pm 0.20$  mg l<sup>-1</sup>,  $26.25 \pm 0.05 - 26.82 \pm 0.04$  (°C) and  $6.90 \pm 0.02 - 6.95 \pm 0.01$ , respectively.

#### 3.2 Reproductive Performances of *H. longifilis* Broodstock with Pond Water Depth

Table 2 shows the fertility assessment of *H. longifilis* broodstock reared in concrete

aquaculture ponds at different water depths. All the measured reproductive parameters increased significantly with pond water depth.

The percentage fertilization and hatching of eggs were highest,  $82.50 \pm 1.44\%$  (80.00-85.00%) and  $91.37 \pm 1.63\%$  (88.24-93.75%), respectively, for broodstock raised in deeper waters at 1.0 m pond water depth. The corresponding values were least in shallower waters at pond water depth of 0.5 m ( $49.50 \pm 0.76\%$ ; 48.50-51.00% and  $50.31 \pm 4.69\%$ , 41.18-56.70%, respectively). Similarly fry survival and fry production success in this study increased significantly ( $P = 0.00$ ) with increase in the broodstock water depth (1.0 m > 0.75 m > 0.5 m) in that order in this study increased significantly ( $P = 0.00$ ) with increase in the broodstock water depth (1.0 m > 0.75 m > 0.5 m) in that order.

Table 1. Mean water quality parameters in egg incubation tanks

Water Parameters	Water Depth		
	0.50 m	0.75 m	1.0 m
Temperature (° C)	$26.24 \pm 0.05$	$26.40 \pm 0.01$	$26.80 \pm 0.02$
pH	$6.90 \pm 0.01$	$6.92 \pm 0.01$	$6.94 \pm 0.25$
Dissolved oxygen (mg l <sup>-1</sup> )	$5.21 \pm 0.20$	$5.62 \pm 0.40$	$5.64 \pm 0.150$

**Table 2. Fertility assessment of *H. longifilis* broodstock reared in concrete aquaculture ponds at different water depths for six months; Mean  $\pm$  SE (min – max)**

Reproductive Performance	Water depth of broodstock culture tank			F-Test	P
	0.50 m	0.75 m	1.0 m		
% Fertilization	49.50 $\pm$ 0.76* (48.50-51.00)	68.17 $\pm$ 0.93* (67.00-70.00)	82.50 $\pm$ 1.44* (80.00-85.00)	232.9	0.000
% Hatching	50.31 $\pm$ 4.69* (41.18-56.70)	67.79 $\pm$ 4.58* (62.22-76.87)	91.37 $\pm$ 1.63* (88.24-93.75)	27.94	0.001
% Survival	8.38 $\pm$ 0.48* (7.45-9.05)	45.66 $\pm$ 2.30* (41.26-49.05)	97.02 $\pm$ 1.01* (95.39-98.87)	905.2	0.000
% Fry Production	2.07 $\pm$ 0.10* (1.90-2.25)	20.95 $\pm$ 0.19(20.60-21.25)*	73.08 $\pm$ 0.53* (72.50-74.15)	12250	0.000

### 3.3 Discussion

The experimental water quality parameters were in line with the recommended standard for aquaculture operation for catfish breeding [20,22] as significant differences in water quality parameters were not observed in all the batches throughout the duration of the study. The idea of managing water depths in a culture enclosure arises in event of insufficient supply or scarce availability of suitable water. Consequently, fish may be cultured in ponds with low water depth for human sustainability.

*Clarias* and *Heterobranchus* actually tolerate low and shallow water pond culture; [1] but the standard pond water depth for catfish culture is 1.0 m which served as control in this study. The study was designed to cover the range at which catfish are generally cultured: 0.5 m to 1.0 m water depth [1], under different water supply scenarios. Successes of the aquaculture venture also require availability of quality fish seeds; apart from suitable pond environment.

This study investigated the optimum water depth required for catfish spawning in hatcheries; with particular reference to reproductive performance of *H. longifilis* broodstock in catfish hatchery operations. The study reveals reproductive parameters such as fertilization, hatchability and survival rates and fry production successes, significantly ( $P = 0.00$ ) increased with water depth of broodstock rearing pond.

The broodstock raised at the standard water level (of 1.0 m) achieved the best reproductive parameters than values for broodstock raised at lower water level of 0.75 m and 0.5 m. The broodstock raised at 0.75 m pond water depth

also outperformed ( $P = 0.00$ ) those from shallow ponds at 0.5 m. The differences might be attributed to the less crowding achieved at (standard) higher water depth of 1.0 m which afford better welfare for fish; compared to much crowding and competition, forced by halving or further reducing available space at 0.7 m and 0.5 m water depth. At lower water depth, fish are concentrated and packed into a limited space with high possibilities of bruising, injuries, cannibalism and reduced feeding or feeding inefficiencies. On the reverse, swim up fry grow better at lower pond water depth while higher depth impose greater energy requirement to swim up for feeding; which equally affect production and viability of gametes [1,23].

Other factors that could contribute to observed variations in the reproductive parameters include: level of egg maturity, hormonal development and overall physiological readiness for reproduction and viability of eggs [2,12]. Larger- sized eggs tend to perform better because of the residual potential energy the eggs possess at less strenuous environment. At normal or ideal level of water (1.0 m in this case), broodstock reserve and utilize energy for egg and immune system development; while they tend to expend their energy for maintenance in strenuous environments at lower water levels (0.5 m and 0.75 m). It could therefore be summarized that hatchery management system is a critical factor for successful fry production [4,21,25,26].

### 4. CONCLUSION

The study reveals that the best reproductive performance is achieved with rearing broodstock at 1.0 m pond water depth. Hence, matured catfish broodstock should be maintained or

conditioned at 1.0 m pond water depth before spawning in hatcheries. Catfish cultured at lower pond water depths of 0.75 m and 0.5 m may be applied to fattening and table-size fish production for processing and consumption.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

#### ACKNOWLEDGEMENTS

The author is grateful to the Akwa Ibom State University for providing enabling environments, ponds and ancillary facilities in the Department of Fisheries and Aquaculture for the experiments.

#### COMPETING INTERESTS

Author has declared that no competing interests exist.

#### REFERENCES

1. Nlewadim AA, Udoh JP, Otoh AJ. Growth response and survival of *Heterobranchus longifilis* cultured at different water levels in outdoor concrete tanks. *Aquaculture, Aquarium, Conservation & Legislation*. 2011;10(1):113-122.
2. Otoh AJ, Udoh JP, Nya E, Asuquo IE. Effect of incremental dilution of catfish sperm with normal saline solution on reproductive performance of *Clarias gariepinus*. *Journal of Wetlands and Waste Management*. 2023;5(1):74-78.
3. Otoh AJ, Udo MT, George UU. Comparative effect of inducing broodstock with natural and artificial hormones on reproductive performances of *Heterobranchus longifilis*. *Tropical Freshwater Biology*. 2022;31:95-102.
4. Otoh AJ, Udo MT, George UU. Comparative growth performance and sex ratio of *Heterobranchus longifilis* and its offspring induced with synthetic hormone and pituitary gland of *Heterobranchus longifilis*. *Journal of Wetlands and Waste Management*. 2023;5(1):106-111.
5. Otoh AJ, Okoko AC, Asangusung PS, Ekanem IE, George UU, Idiong TE. Effects of different ages of African catfish (*Clarias gariepinus*) broodstock on reproductive performance and fries production. *Asian Journal of Fisheries and Aquatic Research*. 2024;26(3):39-47.
6. Otoh AJ, Udoh JP. Semen quality of adult male offspring of *Heterobranchus longifilis* of different parental age groups fed two different iso-nitrogenous feeds. *Tropical Freshwater Biology*. 2018;27:1-22.
7. Otoh AJ, Udoh JP. Age related sperm quality of male *Heterobranchus longifilis* broodstock fed different isonitrogenous feeds. *Tropical Freshwater Biology*. 2018; 27:31-42.
8. Oyeleye OO, Ola SI, Omitogun OG. Ovulation induced in African catfish (*Clarias gariepinus*, Burchell 1822) by hormones produced in the primary culture of pituitary cells. *International Journal of Fisheries and Aquaculture*. 2016;8:67-73.
9. Udoh JP, Otoh AJ. Growth performance of silver catfish, *Chrysichthys nigrodigitatus* fingerlings fed salt-rich diet in fresh water system. *Aquaculture, Aquarium, Conservation and Legislation*. 2017;10(1): 113–122.
10. Udoh JP, Otoh AJ. Distribution and Size of Barnacle *Chelonibia patula* Fouling Blue Crab *Callinectes amnicola* in Southeast Nigeria. *Croatian Journal of Fisheries*. 2016;73:93-102.
11. Udoh JP, Otoh AJ. Aquaculture species biodiversity: relevance of the introduction of nonnative species for aquaculture development in Africa. In: Ayandele IA, Udom GN, Effiong EO, Etuk UR, Ekpo IE, Inyang UG, Edet GE, Moffat I. (editors). *Contemporary discourse on Nigeria's economic profile, a festschrift in honour of Prof. Nyaudoh, U. Ndaeyo. A. Publication of University of Uyo, Uyo, Akwa Ibom State*. 2023;168-197.
12. Otoh AJ, Umanah SI, Udo MT. Influence of feed type and age of parent broodstock on sex ratio of the African catfish, *Heterobranchus longifilis* offspring. *Nigerian Journal of Agriculture, Food and Environment*. 2020;16(4):113-118.
13. Nya E, Udosen I, Otoh A. Effect of Herbal based immunostimulant diets for disease control in African catfish *Clarias gariepinus* against *Aeromonas hydrophila* infections. *Journal of Biology, Agriculture and Healthcare*. 2017;7(16):49-54.
14. Ekanem SB, Otoh AJ, Enyehihi UK, Taege M. The response of juvenile *Chrysichthys*

- nigrodigitatus* (Lacepede) to different components. African Journal of Fisheries and Aquaculture. 2000;2:59-67.
15. Asangusung PS, Uka A, Otoh AJ. Economic evaluation of three hormonal preparations in artificial propagation of *Heterobranchus longifilis* (Valenciennes, 1840). Journal of Aquatic Sciences. 2020;35(2):173-179.
  16. Adewunmi AA, Olaleye VF. Catfish culture in Nigeria: Progress, prospects and problems. African Journal of Agricultural Research. 2011;6(6):1281–1285.
  17. Anibeze CIP, Eze A. Crawfish forage and feeding systems. Reviews in Aquatic Science. 2000;3(1):1-10.
  18. Ofor CO. A comparison of the yield economics of three types of semi-intensive grow out systems in the production of *Heterobranchus longifilis* (Teleostei; clariidae) (Val.1840), in southeast Nigeria. Aquaculture. 2007;269:402-413.
  19. Otoh AJ, Nlewadim AA. Influence of feed type and age of broodstocks on egg quality of *Heterobranchus longifilis*, Journal of Agriculture, Environmental Resource and Management. 2019;4(2):488-500.
  20. Otoh AJ, Ekanem IE, Okoko AC, Asangusung, PS, George UU. Comparative study of inducing broodstock with natural and artificial hormones on reproductive performances of *Clarias gariepinus*. Asian Journal of Fisheries and Aquatic Research. 2024.26(3):31-38.
  21. Otoh AJ, Okoko AC, Ekanem IE, George UU, Asangusung PS. Effects of multiple batches of stripped eggs on the reproductive performances of *Heterobranchus longifilis*. Asian Journal of Research in Zoology. 2024;7(2):9-16.
  22. Otoh AJ, Ekanem IE, Okoko AC, Asangusung, PS, George UU, Idiong TE. Effect of different ages of African catfish (*Clarias gariepinus*) broodstock on reproductive performance and fry production. Asian Journal of Fisheries and Aquatic Research. 2024;26(3): 39-47.
  23. Udoh JP. Survival, development and hatching of *Clarias gariepinus*: (Burchell) eggs in responses to fish extender composition. African Journal of Fisheries and Aquaculture. 2000;2:49-58.
  24. Otoh AJ, George UU, Asangusung PS, Okoko AC, Ekanem IE, Umaha MN. Investigation into the impacts of varied batches of stripped eggs on the reproductive success of *Clarias gariepinus*. Asian Journal of Research in Zoology. 2024;7(2):28-35.
  25. Otoh AJ, Udoh JP. Intergenerational consequences of maternal feed type and age on the egg quality of F1 offspring of *Heterobranchus longifilis*. Journal of Wetlands and Waste Management. 2019; 3(1):53-58.
  26. Otoh AJ, Udoh JP. Comparative study of the growth performances of African catfishes and their hybrids from fry to post fingerlings in outdoor concrete ponds. AKSU Journal of Agriculture and Food Science. 2018;1:81-88.
  27. Otoh AJ, Umanah SI, Udo MT. Comparative study of the effect of feed types and ages of broodstock on reproductive performances of *H. longifilis* in concrete pond. Nigerian Journal of Agriculture, Food and Environment. 2020;16(4):43-50.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/117628>