



Impact of Seed Priming and Seed Rate on Productivity and Economic Profitability of Wheat (*Triticum aestivum* L.)

Devashish Kumar ^a, Suborna Roy Choudhury ^{a*},
Anupam Das ^b, Sushil Kumar Pathak ^a, Sushant ^a,
Chandan Kumar Panda ^c, Pravesh Kumar ^a and Seema ^a

^a Department of Agronomy, Bihar Agricultural University, Bhagalpur, Bihar -813210, India.

^b Department of Soil Science and Agricultural Chemistry, Bihar Agricultural University, Bhagalpur, Bihar - 813210, India.

^c Department of Extension Education, Bihar Agricultural University, Bhagalpur, Bihar -813210, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jeai/2024/v46i82675>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119769>

Original Research Article

Received: 02/05/2024

Accepted: 04/07/2024

Published: 12/07/2024

ABSTRACT

In order to achieve optimal plant growth, the lifecycle of plants encounters various critical stage such as uneven seed germination, induce early flowering, poor and early seedling growth, poor root establishment which ultimately results in low crop yield. Keeping in view of the hypothesis of seed

*Corresponding author: E-mail: subornabau@gmail.com;

Cite as: Kumar, Devashish, Suborna Roy Choudhury, Anupam Das, Sushil Kumar Pathak, Sushant, Chandan Kumar Panda, Pravesh Kumar, and Seema. 2024. "Impact of Seed Priming and Seed Rate on Productivity and Economic Profitability of Wheat (*Triticum Aestivum* L.)". *Journal of Experimental Agriculture International* 46 (8):15-25. <https://doi.org/10.9734/jeai/2024/v46i82675>.

priming improves seedling emergence, reduces seedling emergence time, reduces seed dormancy, expands root growth, enhances nutrient uptake, resulted in better yield and yield contributing characters of plants, a field experiment was planned during Rabi season to figure out the effect of seed priming and seed rate on the yield performance of wheat. The experiment was laid out in factorial randomized block design which replicated thrice. The treatments included four seed priming methods (P1: no seed priming, P2: water seed priming, P3: 1% KNO₃ seed priming, P4: 1% CaCl₂ seed priming) and three seed rates (S1: 100 kg ha⁻¹, S2: 125 kg ha⁻¹, S3: 150 kg ha⁻¹) using the wheat variety HD-2967. The result revealed that the yield and yield attributes of wheat were significantly influenced by the priming techniques. Seed priming with 1% CaCl₂ solution increased grain yield (3977 kg ha⁻¹), number of ear head (275.61 m⁻²), length of ear head (11.29 cm), number of grains per ear head (46.17) and test weight (41.11 g) and was found to be statistically at par with seed priming with 1% KNO₃, but significantly superior over seed priming with water and dry seeding. Different seed rate significantly influenced grain and straw yield and yield contributing characters of wheat. The higher grain yield (3959 kg ha⁻¹), straw yield (5095 kg ha⁻¹) and no. of ear head (287.83) was obtained with the seed rate @150 kg/ha followed by 125 and 100 kg ha⁻¹ of seed rate.

Keywords: Productivity; profitability; seed priming; seed rate; wheat.

1. INTRODUCTION

Wheat (*Triticum aestivum* L.) is the one of the most staple food crops for rabi season grown in India as well as in the state Bihar. It is an annual, hexaploidy (2n = 6X= 42) crop, belongs to the family Poaceae (Gramineae). "Being the most predominant cultivated cereal crop of the world, wheat ranks first (30%) followed by rice (27%) and maize (25%) among the cereal crops. Wheat production of our country is 109.59 million tons from 29.65-million-hectare area with productivity of 3.47 t ha⁻¹ (Directorate of Economics and Statistics of India, 2021-22). Globally, India is the second largest producer of wheat contributing 12% in area and 11.9% in production, next to China. In India, U.P ranks first in area and production while Punjab ranks first in productivity. In Bihar, area under wheat cultivation is 2.22 m ha with production 6.34 m ton having average yield of 2855 kg ha⁻¹" (USDA 2021-22). Agronomic practices, particularly seed rate, play an indispensable role in maintaining optimum plant population and for realizing higher yield. The blanket recommendation of seed rates is not suitable for all varieties and in all agroecosystems. Hence, proper seed rate is a must to provide optimum space and proper availability of moisture and nutrients which certainly will help for better growth and development of a crop [1]. If plant density is above the optimum, the plant development might be poor because of competition for nutrients, light and space. On the other hand, if it is below optimum, at that point the nutrients, space and light won't be used to their full degree, subsequently coming about into poor yield. Yield increase to some extent

increases linearly with the increase in plant population until the light penetration in crop canopy becomes a limiting factor. Hence, seed rate must be standardized when different seed priming's has been used.

"By overcoming the germination limits under both favourable and unfavourable conditions, pre-sowing seed treatments i.e. seed priming, offer a potential solution to poor seedling establishment and boost vigour. But in adverse circumstances, such as the current low temperature in wheat, the effectiveness of seed priming is more noticeable [2,3]. The germination, seedling emergence, growth, and yield characteristics of the crop are all enhanced by the seed priming approach" [4,5,6]. In direct seeded rice, and in pulses like chickpea, soybean, the seed priming with KCl or CaCl₂ enhanced seedling development, stand establishment, and yield performance [7,8]. In wheat grown in late seeding conditions, CaCl₂ priming has also likely resulted in vigorous seedling growth, emergence, and yield performance [2]. The radicle length, plumule length, and root vigour of seedlings were enhanced by priming seeds of normal or low vigour: shoot to ratio and fresh weight [9]. According to Moosavi et al. [10], primed seeds considerably increased the percentage of germination, germination speed, root length, and seed vigour. Seeds with low vigour that were primed performed better during germination [11]. The best methods for increasing seed vigour and improving stand establishment were hydro priming, priming with growth regulators, and priming with different calcium and potassium

salts [12]. Osmo-primed wheat in late conditions had the highest number of productive tillers per unit area, biological yield, grain yield, and harvest index [13,14]. Prior research has demonstrated that, in comparison to non-primed crops, primed crops demonstrated quicker and more consistent emergence, strong stand establishment, and improved performance even in stressed environments [15,16]. Thus, the current study was carried out to access the impact of seed priming and seed rate for enhancing nutrient uptake, growth and performance of wheat.

2. MATERIALS AND METHODS

The experiment was carried during *rabi* seasons of 2020 and 2021, to evaluate the effect of seed priming and seed rate on growth and nutrient uptake of wheat (var. HD-2967) crop. The experimental plot had uniform topography. The experimental site is situated in the Middle Gangetic plain locale of Agro-climatic Zone III (A) in Bihar (latitude-25°23'N, longitude- 87°07'E at an elevation of 37.19 meters above mean ocean level). The climatic condition of this place is tropical to subtropical and somewhat semi-arid in nature and is characterized by very dry summer, moderate rainfall and very cold winter. The soil of the trial plot was Gangetic alluvial in its origin. The field experiment was laid out in factorial randomized block design replicated thrice. Altogether, there were 12 treatment combinations consisting of four seed priming; P₁- no seed priming (dry seed), P₂- seed priming with water, P₃- seed priming with 1% KNO₃, P₄- seed priming with 1% CaCl₂ and three seed rates S₁- seed rate @100 kg ha⁻¹, S₂- seed rate @ 125 kg ha⁻¹ and S₃- seed rate @ 150 kg ha⁻¹. Seed of wheat was primed with 1% KNO₃ and 1% CaCl₂ solutions and also in water for hours. Non-primed seed was included as a control treatment.

2.1 Yields and Yield attributes

The numbers of ear head bearing tillers, length of ear head, Number of grains per ear head were counted at maturity from randomly selected 0.25 m² areas at five locations in net plot area. It was recorded in the same way as the tiller numbers were recorded during progressive growth period. Hence, after threshing and weighing, a random sample of grains was taken from each plot and one-thousand grain were counted at random with the help of mechanical seed

counter and the weight (g) was taken as test weight.

Harvested bundles of wheat plants from each net plot were threshed and winnowed separately . After cleaning, the grain as well as the straw were dried plot wise and consequently the samples were taken from each plot to determine the moisture content with the help of moisture meter. Then the weight of both grain as well as straw were recorded separately at 12 per cent moisture before being subjected to its statistical analysis. The net plot yield was then finally converted into kg ha⁻¹.

Harvest index was calculated by dividing economic (grain) yield by the total biological (grain + straw) yield and expressed as percentage:

$$H. I. = \frac{\text{Economic yield (grain yield) kg ha}^{-1}}{\text{Biological yield (grain + straw yield) kg ha}^{-1}} \times 100$$

Economic Analysis: The common cost of cultivation was worked out by considering all the expenses based on the prevailing market prices incurred during the cultivation process. Grain and straw yield were separately compared to its current market price, and the values were summarised to achieve gross returns. Net returns were calculated by subtracting total cost of cultivation from gross return. The return per rupee investment or benefit: cost ratio was calculated by dividing the net return by the total cost of cultivation.

$$\text{Benefit: Cost ratio} = \frac{\text{Net return}}{\text{Total cost of cultivation}}$$

The experimental data for various growth, yield attributing characters, yield and economics were statistically analyzed by the methods of analysis of variance (ANOVA) [17]. The significance of treatment effects was computed with the help of 'F' (variance ratio) test and to judge the significance of differences between means of two treatments, critical differences (CD) was worked out [18,19,20].

3. RESULTS AND DISCUSSION

After two years of experimentation, it was observed that Seed priming with 1% CaCl₂ (P₄) had the highest number of ear head m⁻² (275.61), the longest ear heads (11.29 cm), the highest number of grains per ear heads (46.17) and the highest 1000 grain weight of (41.11 g)

Table 1. Effect of seed priming and seed rate on yield attributes of wheat

Treatments		Yield attributing characters			
		No. of ear head/m ²	Length of ear head (cm)	No. of grains/ ear head	1000 grain weight (g)
P ₁	Unprimed	261.94	10.11	42.11	37.22
P ₂	Seed priming with water	270.06	10.22	42.56	38.11
P ₃	Seed priming with 1% KNO ₃	274.61	10.94	45.22	40.89
P ₄	Seed priming with 1% CaCl ₂	275.61	11.29	46.17	41.11
SEm±		3.09	0.30	1.05	1.08
CD (P = 0.05)		9.08	0.89	3.08	3.17
Treatments					
S ₁	Seed rate@100 kg/ha	248.25	10.30	43.33	40.08
S ₂	Seed rate@125 kg/ha	275.58	10.96	44.75	39.25
S ₃	Seed rate@150 kg/ha	287.83	10.67	43.96	38.67
SEm±		2.68	0.26	0.91	0.93
CD (P=0.05)		7.86	NS	NS	NS
P X V		NS	NS	NS	NS

Table 2. Effect of seed priming and seed rate on grain yield, straw yield, biological yield and harvest index of wheat

Treatments		Grain yield (kg/ha)	Straw yield (kg/ha)	Biological yield (kg/ha)	Harvest index (%)
P ₁	Unprimed	3627	4897	8525	42.54
P ₂	Seed priming with water	3688	4980	8668	42.55
P ₃	Seed priming with 1% KNO ₃	3935	5262	9198	42.80
P ₄	Seed priming with 1% CaCl ₂	3977	5334	9311	42.72
SEm±		79	108	187	0.07
CD (P = 0.05)		233	316	548	NS
Treatments					
S ₁	Seed rate@100 kg/ha	3577	4808	8385	42.66
S ₂	Seed rate@125 kg/ha	3885	5229	9115	42.63
S ₃	Seed rate@150 kg/ha	3959	5317	9276	42.68
SEm±		68	93	162	0.06
CD (P=0.05)		201	274	475	NS
P X V		NS	NS	NS	NS

Table 3. Effect of seed priming and seed rate on economics of wheat

Treatments		Cost of cultivation (Rs. ha⁻¹)	Gross return (Rs. ha⁻¹)	Net return (Rs. ha⁻¹)	B:C ratio
P ₁	Unprimed	32405	91836	59431	1.83
P ₂	Seed priming with water	32680	92636	59956	1.84
P ₃	Seed priming with 1% KNO ₃	36645	98212	61567	1.68
P ₄	Seed priming with 1% CaCl ₂	34655	99390	64735	1.87
SEm±		-	1631	1631	0.03
CD (P = 0.05)		-	4784	4784	0.11
Treatments					
S ₁	Seed rate@100 kg/ha	33096	89830	56733	1.72
S ₂	Seed rate@125 kg/ha	34096	97386	63289	1.86
S ₃	Seed rate@150 kg/ha	35096	99340	64244	1.84
SEm±		-	1412	1412	0.03
CD (P=0.05)		-	4143	4143	0.09
P X V		NS	NS	NS	NS





Fig. 1. Meteorological data during Experimental Years 2020-2021




Fig. 2. Layout plan of the experimental design of single plots

however, had no any statistical differentiation with other two seed priming i.e. seed priming with water (P2) and seed priming with 1% KNO₃ (P3). Although the lowest values were obtained under the unprimed treatment (Table 1). Seed rate did not influence the various yield attributes significantly. Seed rate@150 kg/ha had the highest number of ear heads m⁻² (287.33), however, maximum length of ear heads (10.96 cm) and maximum number of grains per ear heads (44.75) were found under the seed rate@125 kg/ha.

Significantly the highest grain yield (3977 kg/ha) was recorded in P₄ treatment i.e., seed priming

with 1% CaCl₂ which was statistically at par with value (3935 kg/ha) realized under P₃ i.e., seed priming with 1% KNO₃. However, it was statistically superior with P₁ and P₂ i.e., unprimed (3627 kg/ha) and seed priming with water (3688 kg/ha), respectively. Likewise, the highest straw yield (5334 kg/ha) was recorded in P₄ treatment i.e., seed priming with 1% CaCl₂ which was statistically at par with value (5262 kg/ha) realized under P₃ i.e., seed priming with 1% KNO₃. However, it was statistically superior with P₁ and P₂ i.e., unprimed (4897 kg/ha) and seed priming with water (4980 kg/ha), respectively. So far as seed rate is concerned, significantly the highest grain yield of 3959 kg/ha as well as

straw yield of 5317 kg/ha was obtained from seed rate@150 kg/ha which was statistically at par with value (grain yield 3885 kg/ha and straw yield 5229 kg/ha) realized under S₂ i.e., seed rate@125 kg/ha (Table 2). However, it was statistically superior with S₁ i.e., seed rate@100 kg/ha. However, their interaction (P X S) did not reach the level of significance. Significantly the highest biological yield (9311 kg/ha) was recorded in P₄ treatment i.e., seed priming with 1% CaCl₂ which was statistically at par with value (9198 kg/ha) realized under P₃ i.e., seed priming with 1% KNO₃. However, it was statistically superior with P₁ and P₂ i.e., unprimed (8525 kg/ha) and seed priming with water (8668 kg/ha), respectively. Similarly, the highest biological yield of 9276 kg/ha was obtained from seed rate@150 kg/ha which was statistically at par with value (9115 kg/ha) realized under S₂ i.e., seed rate@125 kg/ha. However, it was statistically superior with S₁ i.e., seed rate@100 kg/ha. A perusal of ANOVA table revealed that the differences due to seed priming and seed rate did not significantly influence the harvest index. Even their interaction effects did not create any significant difference in this regard. However maximum numerical value of harvest index 42.80 was recorded in P₃ i.e., seed priming with 1% KNO₃ and lowest value of harvest index 42.54 was obtained at P₁ i.e., unprimed. Moreover, numerically the highest harvest index of 42.68 was recorded in case of seed rate@150 kg/ha. It was followed by seed rate@100 kg/ha (42.66) and the lowest numerical value of harvest index (42.63) was noted in seed rate@125 kg/ha. This is because of the fact that in primed seeds, the nutrients can easily be accessed by germinating seeds for its growth and vigour [21,22,23]. Studies proved that seed priming had a positive impact on various plant species specially to endure the environmental stress [24,25,26]. It aids the plant to elevate antioxidant enzyme activity and osmolytes and to build up the resistance to abiotic and non-abiotic stresses [27,28]. Moreover, the seed priming with CaCl₂ or KNO₃ improved the root growth to enhance the nutrient uptake and hastened the overall plant growth during the early stage of the crop [21,29,30].

Significantly the highest net return of 64735 Rs./ha was recorded under application of seed priming with 1% CaCl₂, which was statistically at par with the value realized under P₂ (59956 Rs./ha) and P₃ (61567 Rs./ha) i.e., seed priming with water and seed priming with 1% KNO₃, respectively. However, it was statistically

superior with the value obtained under P₁ (59431 Rs./ha) i.e., unprimed (Table 3). Similar trend was observed under benefit cost ratio, where, the highest benefit cost ratio (1.87) was recorded under P₄ i.e., seed priming with 1% CaCl₂ which was statistically at par with P₁ (1.83) and P₂ (1.84) i.e., unprimed and seed priming with water, respectively. However, it was statistically superior with the value realized under P₃ (1.28) i.e., seed priming with 1% KNO₃. Among all the seed rate, maximum net return of 64244 Rs./ha was recorded from seed rate@150 kg/ha which was statistically at par with the seed rate@125 kg/ha 63289 (Rs./ha) The lowest net returns of 56733 Rs/ha was recorded from seed rate@100 kg/ha. The cost-effective seed priming techniques strengthen the plants and enhance drought resistance through increased photosynthesis pigments, antioxidant defense, osmotic adjustments as well as membrane integrity [30,31].

4. CONCLUSIONS

We experienced from the experimental results that the priming method as pre-germinative enhancement technique significantly improved the productivity as well as profitability of wheat. Seed priming with 1% CaCl₂ resulted in better seedling emergence, uniform stand establishment, earlier flowering and, ultimately, better crop yields with good economic returns. Besides, the indiscriminate use of seeding rates not only increases production costs but usually decrease wheat grain yield. Therefore, the priming methods along with proper seed rate have the large potential for further enhancement in wheat production. The priming methods adopted should be simple and affordable for its easy spread and adaptability to the farmers.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. El-Wakeel, Sally E., Mohamed Mansour, Ashgan M. Abd El-Azeem, Sahar A.

- Ebrahim, and Tahany Noreldin. Field and economic evaluation of barley productivity As affected by seed rates and slow-release nitrogen fertilizer levels under rainfed conditions. Asian Journal of Advances in Agricultural Research. 2024; 24(4):16-26.
Available:<https://doi.org/10.9734/ajaar/2024/v24i44499>
2. Farooq M, Basra SMA, Rehman H, Saleem BA. Seed priming enhances the performance of late sown wheat (*Triticum aestivum* L.) by improving chilling tolerance. J. Agron. Crop Sci. 2008; 194(1):55-60.
 3. Riaz-ud-Din MS, Ghulam A, Naeem H, Makhdoom, Aziz Ur Rehman, Effect of temperature on development and grain formation in spring wheat. Pak. J. Bot. 2010;42:899–906.
 4. Devashish Kumar, Sushil Kumar Pathak, Shweta Kumari, Aditya Shri, Sanju Kumari, Priya Kumari, Chandini, Rakesh Kumar, Vibha Kumari and Prabhat Ranjan, Effect of seed priming and seed rate on LAI and uptake of nutrients in wheat (*Triticum aestivum* L.) The Pharma Innovation Journal. 2021;10(10): 779-782
 5. Agawane RB, Parhe SD. Effect of seed priming on crop growth and seed yield of soybean [*Glycine max* (L.) merill]. The Bioscan. 2015;10:265-70.
 6. Chavan NG, Bhujbal GB, Manjare MR. Effect of seed priming on field performance and seed yield of soybean [*Glycine max* (L.) Merrill] Varieties. The Bioscan. 2014;9(1):111-114.
 7. Farooq M, Basra SMA, Hafeez K, Rice seed invigoration by osmohardening. Seed Sci. Technol. 2006b;34:181–186.
 8. Farooq M, Hussain M, Imran M, Ahmad I, Atif M, Alghamdi SS. Improving the productivity and profitability of late sown chickpea by seed priming. International Journal of Plant Production. 2019 Jun 1;13:129-39.
 9. Kausar M, Mahmood T, Basra SMA, Arshad M, Invigoration of low vigor sunflower hybrids by seed priming. Int. J. Agric. Biol. 2009;11:521-528.
 10. Moosavi A, Afshari RT, Sharif-Zadeh F, Ayneband A. Effect of seed priming on germination characteristics, polyphenoloxidase, and peroxidase activities of four amaranth cultivars. J. Food, Agric. Env. 2009;7:353–358.
 11. Bray CM. Biochemical processes during the osmopriming of seeds. In: Seed Development and Germination, pp: 767–789. Kigel, J. and G. Galili. (eds.). New York: Marcel Dekker; 1995.
 12. Basra SMA, Farooq M, Khaliq A. Comparative study of pre-sowing seed enhancement treatments in fine rice (*Oryza sativa* L.). Pak. J. Life Soc. Sci. 2003;1(1):21-25.
 13. Sattar A, Cheema MA, Farooq M, Wahid MA, Wahid A, Babar BH, Evaluating the performance of wheat cultivars under late sown conditions. Int. J. Agric. Biol. 2010;12:561–565.
 14. Amin R, Khan AZ, Khalil SK, Khalil IH. Effect of osmopriming sources and moisture stress on wheat. Pak. J. Bot. 2012;44(3):867-871.
 15. Eivazi A. Induction of drought tolerance with seed priming in wheat cultivars (*Triticum aestivum* L.). Acta Agric. Slov. 2012;99(1):21-29.
 16. Gupta, Ajay. Effect of seed rate on yield of wheat (*Triticum aestivum*) under front line demonstrations in poonch. Asian Journal of Agricultural Extension, Economics & Sociology. 2022;40(10): 795-97.
Available:<https://doi.org/10.9734/ajaees/2022/v40i1031144>
 17. Panse WG, Sukhatme PV. Statistical methods for agriculture workers, 3rd Rev. Ed., ICAR, New Delhi; 1978.
 18. Cochran GW. Cox MG. Experimental Design. John Willey and Sons, Inc. U.S.A; 1967.
 19. Fischer RA, Yates F. Statistical tables for biological agricultural and medical research (6th Ed.). oliver and boyd, Edinburgh, Tweed dale Cowot, London; 1963
 20. Gomez KA, Gomez AA. Statistical procedures for agricultural research. John Willey and Sons, Inc., New York, 1984;180.
 21. Farooq M, Almamari SAD, Rehman A, Al-Busaidi WM, Wahid A, Al-Ghamdi SS. Morphological, physiological and biochemical aspects of zinc seed priming-induced drought tolerance in faba bean. Sci. Hortic2021;281:109894. DOI: 10.1016/j.scienta.2021.109894
 22. Babu R, Kakraliya SK, Prakash L, Kumar P, Yadav RA. Effect of plant geometry and seed rates on growth, yield attributes, productivity as well as weed dynamics of

- wheat (*Triticum aestivum*). Int J Curr Microbiol App Sci. 2017;6:81-88.
23. Giri GS, Schillinger WF. Seed priming in winter wheat for germination, emergence and yield. Crop Sci. 2003;43: 2135-2141.
 24. Bukhari MA, Ahmad Z, Ashraf MY, Afzal M, Nawaz F, Nafees M, et al. Silicon mitigates drought stress in wheat (*Triticum aestivum* L.) through improving photosynthetic pigments, biochemical and yield characters. Silicon. 2021;13: 4757–4772.
DOI: 10.1007/s12633-020-00797-4
 25. Farooq M, Basra SMA, Rehman H. Seed priming enhance the performance of late sown wheat (*Triticum aestivum* L.) by improving chilling Tolerance. Journal Agronomy & Crop Science; 2007. ISSN 0931-2250
 26. Khan MB, Ghurchani M, Hussain M, Mahmood K, Wheat seed invigoration by pre-sowing chilling treatments. Pak. J. Bot. 2010;42:1561–1566.
 27. Sattar A, Wang X, Abbas T, Sher A, Ijaz M, Ul-Allah S, et al. Combined application of zinc and silicon alleviates terminal drought stress in wheat by triggering morpho-physiological and antioxidants defense mechanisms. PloS One. 2021; 16(10):e0256984.
DOI: 10.1371/journal.pone.0256984
 28. Ali H, Iqbal N, Shahzad AN, Sarwar N, Ahmad S, Mehmood A. Seed priming improves irrigation water use efficiency, yield and yield components of late-sown wheat under limited water conditions. Turk. J. agric. 2013;37:534-544.
 29. Malko MM, Khanzada A, Xiao WANG, Samo A, Li Q, Dong JIANG, et al. Chemical treatment refines drought tolerance in wheat and its implications in changing climate: A review. Plant Stress. 2022;100118.
DOI: 10.1016/j.stress.2022.100118
 30. Saha D, Choyal P, Mishra UN, Dey P, Bose B, Prathibha MD, et al. Drought stress responses and inducing tolerance by seed priming approach in plants. Plant Stress. 2022;4:100066.
DOI: 10.1016/j.stress.2022.100066
 31. Sime G, Aune JB. On-farm seed priming and fertilizer micro-dosing: Agronomic and economic responses of maize in semi-arid Ethiopia. Food and Energy Security. 2020 Feb;9(1):e190.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/119769>