



Influence of IBA and NAA on Rooting of Terminal Cuttings of Chrysanthemum (*Dendranthema grandiflora* L.)

**Sivakumar. B ^a, Senthilkumar. P ^{b++*}, Deivamani. M ^c,
Sasikumar. K ^b, Ayyadurai. P ^d, K. Govindan ^b,
Senthilkumar. T ^c and Mangammal. P ^e**

^a Forest College and Research Institute, Tamil Nadu Agricultural University, Mettupalayam, Tamil Nadu, India.

^b Regional Research Station, Tamil Nadu Agricultural University, Paiyur, Krishnagiri Dt, Tamil Nadu, India.

^c ICAR-Krishi Vigyan Kendra, Dharmapuri, Tamil Nadu Agricultural University, Tamil Nadu, India.

^d Centre of Excellence for Millets, Tamil Nadu Agricultural University, Athiyandal, Tamil Nadu, India.

^e Horticultural College and Research Institute, Tamil Nadu Agricultural University, Paiyur, Krishnagiri Dt, Tamil Nadu, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jsrr/2024/v30i102493>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/122067>

Original Research Article

Received: 01/08/2024

Accepted: 04/10/2024

Published: 09/10/2024

⁺⁺Associate Professor (Agrl. Nematology);

*Corresponding author: E-mail: senthikumar@tnau.ac.in;

ABSTRACT

The present investigation was conducted at the Agricultural College and Research Institute, Vazhavachanur, Tiruvannamalai to study the effect of Indole-3-Butyric Acid (IBA) and Naphthalene Acetic Acid (NAA) on the rooting of terminal cuttings of chrysanthemum. An application of growth regulators (IBA and NAA) had a positive effect on the rooting of chrysanthemum terminal cuttings. The result of the experiment revealed that the treatment of IBA @ 400ppm significantly increased rooting percentage (89%), length of roots (5.67 cm), root spread (39.18 cm), number of roots (22.67), fresh weight of roots (4.94 g), plantlet weight (1.17 g), and plant height (8.87 cm) followed by NAA @ 400 ppm (79.33%), while minimum rooting percentage (47) was observed in control. It can be concluded that the treatment of IBA @ 400ppm significantly increased rooting percentage, roots length, number of roots, plantlet weight and plant height has greater survival percentage in the terminal cuttings of chrysanthemum (*Dendranthema grandiflora*).

Keywords: *Dendranthema grandiflora*; terminal cuttings; indole-3-butyric acid (IBA); naphthalene acetic acid (NAA).

1. INTRODUCTION

India has diverse agro-climatic conditions, and flower crops are cultivated throughout the year in conventional and non-conventional areas. Floriculture is a fast growing agro-industry and has been associated with social culture and heritage in India. Since ancient times, it has been a tradition of growing flowers largely for religious purposes, the perfume industry, and landscaping. Chrysanthemum (*Dendranthema grandiflora* L.) belongs to the family Asteraceae, a popular perennial flowering plant, second only to roses in the international flower trade [1]. It is known for its long vase life and comes in a wide range of types, sizes, and colors. Chrysanthemum is used both as cut and loose flowers. Loose flowers are used for making garlands and for social and religious functions. Standard chrysanthemums are grown for cut flowers in bouquets and vase decorations [2,3]. The rooting percentage low is a serious problem for the potted chrysanthemum because the cuttings are rooted in the definitive pot and the rooting unevenness. Consequently it will reflect on the unevenness of the final product. The commercial propagation of chrysanthemum is done by rooting of herbaceous cuttings previously treated with synthetic auxin, with indol butyric acid (IBA) being the most used, at a concentration of 1000 mg·L⁻¹ [4]. In India, it is commonly propagated through terminal cuttings and suckers from June to August. The propagation by stem cuttings is more feasible with the application of auxins. However, the root development percentage is low in some varieties, which can overcome by the use of growth regulators. The effect of IBA and NAA on rooting of chrysanthemum terminal cuttings indicated

that cuttings with IBA at 400 ppm performed well concerning the root development percentage [5]. Sidhu and Singh [6] studied the effect of auxins on the rooting of terminal cuttings of *Chrysanthemum morifolium* cv. Flirt and recorded the highest number of roots per cutting with IBA at 250 ppm. The treatment of cuttings with auxins increases the number of adventitious roots that can naturally regenerate into roots. [7] reported that treating chrysanthemum cuttings with IAA and IBA at concentrations 0, 100, 200, 300 or 400 ppm significantly increased the number of roots compared with non- treated cuttings. However, their effectiveness varies among different species and varieties. Rooting in chrysanthemum cuttings is facilitated by auxins, such as Indole Butyric Acid (IBA), Naphthalene Acetic Acid (NAA), and Indole Acetic Acid (IAA), which are commonly applied in liquid or powder form to promote rooting [8]. The present study investigates the effect of Indole-3-Butyric Acid (IBA) and Naphthalene Acetic Acid (NAA) on the rooting of terminal cuttings of chrysanthemum.

2. MATERIALS AND METHODS

The present investigation was conducted in a shade net house at Agricultural College and Research Institute, Vazhavachanur, Tiruvannamalai, Tamil Nadu. The experiment was laid out in Completely Randomized Design (CRD) with twelve treatments and three replications. The treatment details are furnished in Table 1. The required quantity of growth regulators IBA and NAA were carefully weighed using an electronic balance and dissolved in 10 ml of 95% ethyl alcohol with thorough shaking. The same was transferred to a volumetric flask to make the required volume with water. The

growth regulators were prepared at 100, 200, 400, and 800 ppm concentrations. The chrysanthemum terminal cuttings were soaked with the growth regulators at different concentrations for 24 hours. River sand was treated with 0.2% carbendazim fungicide to prevent soil diseases. Same size pots were selected and filled with the media. Cocopeat was used as the rooting medium, filled into portraits. All pots were kept in partial shade. The rooting percentage, length of roots (cm), root spread (cm), number of roots plantlet⁻¹, fresh weight of roots (g), plantlet weight (g), and plant height (cm), were recorded in each treatment. The data were statistically analysed following the procedures outlined [9].

Table 1. Treatment details

Treatments
T ₀ - Control (water spray)
T ₁ - IBA @ 100 ppm
T ₂ - IBA @ 200 ppm
T ₃ - IBA @ 400 ppm
T ₄ - IBA @ 800 ppm
T ₅ - NAA @ 100 ppm
T ₆ - NAA @ 200 ppm
T ₇ - NAA @ 400 ppm
T ₈ - NAA @ 800 ppm
T ₉ - IBA 100 ppm + NAA @ 100 ppm
T ₁₀ - IBA 200 ppm + NAA @ 200 ppm
T ₁₁ - IBA 400 ppm + NAA @ 400 ppm

3. RESULTS AND DISCUSSION

The effects of different concentrations of auxins varied significantly among treatments in terms of cutting length, rooting percentage, number of roots per cutting, root length, days to root initiation, and plant survival in pots, as shown in Table 2. The observation on rooting revealed that all the growth regulators had significant effect on rooting percentage (Table 2). The maximum rooting percentage was recorded in IBA @ 400 ppm (89%), followed by NAA @ 400 ppm (79.33%), while minimum rooting percentage (47) was observed in control (Fig. 1). The results are also in conformity with the findings of [10]. It was also found that all the growth regulator treatments showed significant effect on length of roots. The maximum root length (5.67 cm) was observed in IBA @ 400 ppm followed by IBA @ 100 ppm + NAA @ 100 ppm combination (5.17cm), whereas minimum length of roots (3.60 cm) was recorded in control. Similar result was obtained by [5,11] in *Juniperus procera* and [12] in jasmine.

All the growth regulator treatments showed a significant effect on root spread as compared to the control. The maximum root spread (39.18cm²) was observed in IBA @ 400 ppm followed by NAA @ 800 ppm (38.89cm²) and it was minimum in control (25.27cm²). Indole butyric acid treatment recording maximum root spread in chrysanthemum cuttings was also observed by [5]. The growth regulators showed a significant effect on the number of roots plantlet⁻¹. The maximum number of roots per cutting (22.67) was recorded in IBA @ 400 ppm, followed by IBA @ 100 ppm + NAA @ 100 ppm combination (20.67) as compared to control (15.67). [13] studied the effect of different levels of IBA on rooting, growth and flower yield of chrysanthemum cv. Sonali Tara and reported maximum survival percentage with IBA at 2000ppm followed by IBA at 1000ppm. [14] Panahi and Morteza16, studied the effect of auxins on rooting and flowering of carnation cultivars (*Dianthus caryophyllus* L.) and recorded highest number of roots per cutting with IBA at 100ppm. Moreover, Chrysanthemum cuttings treated with IBA at 250 ppm exhibited the longest root lengths at 10 and 20 DAP (6.53 cm and 11.98 cm, respectively), matching those treated with NAA at 250 ppm (6.09 cm and 11.08 cm, respectively), while the control group showed the shortest shoot lengths (3.49 cm and 5.34 cm, respectively), The maximum number of roots (53.07) was observed in cuttings treated with IBA at 250 ppm, similar to those treated with NAA at 200 ppm (52.98), while the control group had the minimum (26.00), likely due to increased cell division and enlargement [15]. The IBA at 250 ppm treatment also resulted in the shortest rooting time (5.33 days), comparable to NAA at 200 ppm (5.63 days), with the control group requiring the longest time (11.13 days), possibly due to endogenous auxin translocation and the impact of exogenous IBA on cell division and elongation [16]. Auxin treatment significantly reduces the time-to-rooting and early rooting was recorded with the 500 ppm IBA (20.60 days) over 200 ppm (22.53 days). This might be due to the internal auxin amount is not enough for root induction of cuttings. IBA significantly increased the number of roots per cutting and found maximum (13.84) in 500 ppm IBA. The more number of roots obtained with the application of growth chemicals clearly reflects that they not only initiate rooting but also help in subsequent rapid growth of roots in numerical strength [17]. Shin and Zee [18] also reported that IBA treated cuttings had increased the number of roots per cutting in chrysanthemum, [19] also observed

Table 2. Effect of IBA and NAA on rooting percentage, length of roots (cm), root spread (cm²), no. of roots plantlet⁻¹, fresh weight of roots (g) and height of plant in Chrysanthemum terminal cuttings

Treatments	Rooting Percentage	Length of roots (cm)	Root spread (cm ²)	No. of roots plantlet ⁻¹	Fresh weight of roots (g)	Weight of plantlet (g)	Height of plant (cm)
T ₁ IBA @100 ppm	53.00	3.83	30.57	17.67	3.29	0.68	7.70
T ₂ IBA @ 200 ppm	60.00	4.50	34.78	17.00	3.12	0.93	7.73
T ₃ IBA @ 400 ppm	89.00	5.67	39.18	22.67	4.94	1.17	8.87
T ₄ IBA @ 800 ppm	75.67	4.27	31.29	20.67	4.10	0.59	8.57
T ₅ NAA @ 100 ppm	59.00	3.80	30.09	17.00	3.08	0.58	7.93
T ₆ NAA @ 200 ppm	53.00	4.13	29.66	16.33	2.93	0.63	7.63
T ₇ NAA @ 400 ppm	79.33	4.47	34.44	20.00	3.71	0.75	8.03
T ₈ NAA @ 800 ppm	55.00	4.67	38.89	18.00	3.78	0.77	8.47
T ₉ IBA @ 100 ppm + NAA @ 100 ppm	71.00	5.17	29.17	20.67	3.43	0.77	7.67
T ₁₀ IBA @ 200 ppm + NAA @ 200 ppm	55.00	4.20	27.52	17.00	3.44	0.61	7.83
T ₁₁ IBA @ 400 ppm + NAA @ 400 ppm	52.33	4.10	26.61	16.33	3.44	0.60	7.33
T ₁₂ Control (water/spray)	47.00	3.60	25.27	15.67	2.88	0.52	7.17
F-test	S	S	S	S	S	S	S 0.12 0.26
S. Ed (±)	2.37	0.09	0.59	0.43	0.10	0.04	
C. D.(P= 0.05)	4.89	0.18	0.22	0.89	0.22	0.08	




Fig. 1. Effect of IBA and NAA in different concentration on rooting percentage

similar results in carnation stem cutting. Bose and Mandal [20] obtained not only high percentage of rooting but also a large number of roots in many root trees and climbers by treating with IBA and NAA in talc. The growth regulators had a significant effect on the fresh weight of roots. The maximum fresh weight of roots (4.94 g) was observed in IBA @ 400 ppm, followed by IBA at 400 ppm (4.10 g) while the minimum fresh weight of roots (2.88 g) was recorded in control. A similar result was obtained in the stem cuttings of chrysanthemum by [5,6]. All the treatments showed a significant effect on plantlet weight. The maximum plantlet weight (1.17 g) was also recorded in IBA at 400 ppm, followed by IBA at 200 ppm (0.93 g). A similar results were obtained in chrysanthemum cuttings by [5,21]. The plant height was also influenced by the growth regulator treatments. The maximum plant height (8.87cm) was recorded in IBA at 400 ppm, followed by IBA at 400 ppm (8.57cm), [19] and [21] also obtained increased in plant height with IBA treatment.

4. CONCLUSION

Based on the experiment, it can be concluded that the treatment of IBA @ 400ppm significantly increased rooting percentage, roots length, number of roots, plantlet weight and plant height has greater survival percentage in the terminal cuttings of chrysanthemum (*Dendranthema grandiflora*).

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during the writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Anonymous, Indian Horticulture Data Base. Area, production and productivity of flowers, National Horticulture Board, Gurguan. 2006;3-4.
2. Singh P, Chettri R. A new propagation method for rapid multiplication of chrysanthemum under *in vivo* conditions. Int. J Conserv. Sci. 2013;4(1): 95-100.
3. Manjunath DR, Patil RT, Patil BC, Mukund Shiragur, Sachin Kumar T, Nandimath, Srikantaprasad D. Effect of auxins on rooting of terminal cuttings of chrysanthemum (*Dendranthema grandiflora*). The Pharma Innovation Journal. 2022;11(3):1436-1439.
4. Nogueira MR, Ferraz MV, Bezerra AKD, Mazzini-Guedes RB, Costa CRX, de

- Almeida LCP, Pereira STS, Pivetta KFL. Indolbutyric acid and time of the year influence on rooting of chrysanthemum cuttings. American Journal of Plant Sciences. 2018;9:507-516. Available:<https://doi.org/10.4236/ajps.2018.93038>
5. Grewal HS, Kumar R, Chauhan, Rapinder. Effect of IBA and NAA on rooting in Chrysanthemum (*Dendranthema grandiflora*) terminal cuttings. J. Orna. Hort. 2005;8(3):230-232.
 6. Sidhu GS, Singh P. Effect of auxins on propagation in chrysanthemum morifolium. Ind. Flor. in the new millennium, Lal Bagh, Bangalore. 2002;25-27.
 7. Amin KA, Hashim HA. Effect of two types of plant auxin on rooting of cuttings of chrysanthemum (*Chrysanthemum morifolium*). Third Scientific Conference of Technical Education. Agricultural Research. 1992;477-487.
 8. Poteza GJ. Effect of rooting hormone and media on propagation of chrysanthemum (*Dendranthema grandiflorum* Ramat Kitam.), M.Sc. (Agri.) Thesis, Punjab Agri. Univ., Ludhiana (India); 2020.
 9. Panse VC, Sukhatme PV. Statistical methods for agricultural workers. 2nd edition, ICAR, New Delhi. 1967;328.
 10. Cuquel FL, Minami K. Rooting of chrysanthemum cuttings treated with IBA applied in talc. Scientia Agricola. 1994; 51:28-35.
 11. Beshe D, Nagesh L. A sexual propagation of *Juniperus procera* from Ethiopia; a contribution. Forest Ecology and Management. 1998;112: 179-190.
 12. Nagaraja GS, Raj BGM, Guru Prasad TR. Effect on intermittent mist and growth regulators on propagation of *Jasminum grandiflorum* by different types of cuttings. Haryana J. Hort. Sci. 1991;20:183-188.
 13. Ranpise SA, Bharmal VS, Darwade RT. Effect of different levels of Indole Butyric Acid (IBA) on rooting, growth and flower yield of chrysanthemum cv. Sonali Tara. J. Orn. Hort. 2004;7(3-4):331-337.
 14. Panahi, R, Khoushkhohi Morteza., Effects of auxins on rooting and flowering of two cultivars of carnation (*Dianthus caryophyllus* L.). Iranian J. Hort. Sci. Techn. 2000;1(3-4): 91-108.
 15. Ratna Kumari N, Bharathi N. The Influence of Auxins on the Rooting Process of Terminal Cuttings of Chrysanthemum (*Dendranthema grandiflora*). Biological Forum – An International Journal. 2024; 16(2):121-123.
 16. Yusnita Y, Jamaludin J, Agustiansyah A, Hapsoro D. A combination of IBA and NAA resulted in better rooting and shoot sprouting than single auxin on Malay apple [*Syzygium malaccense* (L.) Merr. and Perry] stem cuttings. J Agric. Sci. 2017;40(1):80-90.
 17. Prince, Arvind Mailk, Vivek Beniwal. Influence of Indole-3-Butyric Acid on Rooting Efficacy in Different Carnation (*Dianthus caryophyllus* L.) Genotypes under Protected Condition. Chemical Science Review and Letters. 2017;6(23): 1858-1862.
 18. Shin HJ, Zee JM. Effect of propagation methods and growth regulators on Chrysanthem cuttings. J. Korean Soc. Hort. Sci. 1997;20:111-116.
 19. Sharma RK, Singh VP, Pande DS, Misra HR. Effect of IAA and IBA on growth and flowering of chrysanthemum Cv. Margaeite, Plant Sci. 1978;10:179-181.
 20. Bose TK, Mandal DP. Rooting of cutting under mist. Indian J. Agric. Sci. 1973; 43(3):229-233.
 21. Samanda N, Ormvod DP, Adedipe NO. Rooting of chrysanthemum stem cuttings as affected by phosphonic acid and IBA. Ann. Bot. 1972;63:961-965.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/122067>