



The Role of Organic Amendments and Their Impact on Soil Restoration: A Review

**Godspower Oke Omokaro^{a,b*},
Itohanosa Omolara Osarhiemen^{b,c}, Vivian Idama^{b,c},
Edmond Osemwengie Airueghian^b,
Sokari Tomipapamieari West^b, Franklin Endurance Igbigbi^d,
David Chibuokem Nnake^e, Etyo Obolokor^f,
Abdulraheem Ahmed^g and Vessy Onyeamachi Omoshie^h**

^a Institute of Environmental Engineering, Peoples Friendship University of Russia, Moscow, Russia.

^b Faculty of Agriculture, University of Benin of Benin, Benin City, Nigeria.

^c School of Public Health, I.M. Sechenov First Moscow State Medical University, Moscow, Russia.

^d College of Science, Federal University of Petroleum Resources, Effurun, Delta State, Nigeria.

^e Applied Natural Sciences, Enugu State University of Science and Technology, Enugu, Nigeria.

^f College of Technology, Federal University of Petroleum Resources, Effurun, Delta State, Nigeria.

^g Faculty of Health Science, Environmental Health Science, Kwara State University, Nigeria.

^h Department of Science and Technology, Delta State Polytechnic, Ozoro, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajee/2024/v23i11620>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/125606>

Review Article

Received: 21/08/2024

Accepted: 23/10/2024

Published: 29/10/2024

*Corresponding author: E-mail: omokaro.kelly@gmail.com;

Cite as: Omokaro, Godspower Oke, Itohanosa Omolara Osarhiemen, Vivian Idama, Edmond Osemwengie Airueghian, Sokari Tomipapamieari West, Franklin Endurance Igbigbi, David Chibuokem Nnake, Etyo Obolokor, Abdulraheem Ahmed, and Vessy Onyeamachi Omoshie. 2024. "The Role of Organic Amendments and Their Impact on Soil Restoration: A Review". Asian Journal of Environment & Ecology 23 (11):41-52. <https://doi.org/10.9734/ajee/2024/v23i11620>.

ABSTRACT

Soil organic matter (SOM) plays a pivotal role in maintaining soil health by enhancing its physical, chemical, and biological properties. It improves soil structure, water retention, nutrient availability, and microbial activity, contributing significantly to sustainable agriculture. Organic amendments, including animal manure, compost, biosolids, crop residues, and various organic by-products, are widely used to restore degraded soils by increasing SOM content. These amendments not only supply essential nutrients but also promote biological aggregation, suppress soil-borne pathogens, and improve soil's resistance to environmental stressors such as drought and heavy metal toxicity. The positive impacts of organic amendments on soil properties are well-documented, with key benefits including increased soil organic carbon (SOC), improved nutrient cycling, enhanced microbial diversity, and greater water-holding capacity. Organic materials such as compost and animal manure have been shown to significantly improve soil fertility and restore degraded soils. The application of green manure and crop residues also boosts soil biodiversity, enhances rhizospheric processes, and supports overall ecosystem resilience. This paper emphasizes the importance of adopting holistic soil management practices that include organic amendments, conservation tillage, and sustainable cropping systems to address soil degradation. By focusing on replenishing SOM, improving nutrient availability, and increasing soil biodiversity, these practices can contribute to the long-term sustainability of agricultural systems and environmental health. The restoration of soil organic matter is essential for enhancing soil quality and mitigating degradation, ultimately ensuring food security and ecosystem stability for future generations.

Keywords: Amendments; soil restoration; soil properties; agriculture; soil fertility; organic manure.

1. INTRODUCTION

Human activities have profoundly altered Earth's ecosystems, leading to significant environmental degradation, particularly through changes to soil properties. Anthropogenic influences, such as agricultural practices, industrial development, and urban expansion, have resulted in the deterioration of soil's physical, chemical, and biological integrity. For instance, Ellis et al. (2010) estimated that in 1700, 95% of Earth's ice-free land was categorized as wild or semi-natural landscapes. However, by 2000, human activities had transformed 55% of this land into rangelands, croplands, and urbanized areas, leaving only 45% of natural habitats intact. This shift not only reduces biodiversity but also disrupts soil ecosystems. Zika and Erb (2009) reported that dryland degradation alone leads to the loss of 2% of global terrestrial net primary productivity annually, further diminishing soil quality. Without organic matter, the soil's ability to retain water, regulate temperature, and support microbial activity is significantly compromised. This creates a cycle of degradation that hinders future regeneration, making it increasingly difficult to restore degraded soils for productive use. Given the importance of soil organic matter for soil health, restoring degraded lands often hinges on

rebuilding a surface horizon rich in organic matter. While importing topsoil is an option, it is often economically and environmentally unsustainable, as the donor areas themselves suffer degradation (Larney and Pan 2006). As an alternative, the use of organic amendments—such as manure, biosolids, and other organic wastes—has been practiced for centuries to restore soil fertility (Montgomery, 2007). These materials help replenish soil organic matter and improve soil structure, nutrient content, and biological activity (Akala and Lal, 2000). However, despite the well-documented benefits of organic amendments (Stewart et al., 2000), they are often viewed with skepticism due to concerns about potential contaminants, such as heavy metals, pathogens, and excess nutrients, which can leach into water sources (Larney et al., 2011). This paper explores the critical role of organic matter in soil restoration, examining how organic amendments can enhance soil properties and contribute to the recovery of degraded lands.

2. SOIL ORGANIC AMENDMENTS

Organic matter plays a crucial role in maintaining the health and functionality of soil ecosystems. It provides the essential substrates for decomposing microbes, which in turn release

nutrients that plants require for growth. Furthermore, organic matter enhances soil structure, improves water-holding capacity, increases natural resistance to soil-borne pathogens, and mitigates the toxicity of heavy metals (Abiven et al., 2009; Park et al., 2011). Given these critical functions, restoring depleted organic matter levels in soil is a vital aspect of soil management. The application of organic amendments, such as compost, has proven to be an effective strategy for improving both the physical structure and the chemical and biological fertility of soils, as well as for suppressing soil-borne diseases (Scotti et al., 2013; Zaccardelli et al., 2013). The use of organic amendments to enhance soil fertility dates back thousands of years. Ancient civilizations, including the Greeks and Romans, recognized the benefits of applying animal manure and human sewage to soils, as well as the advantages of rotating crops with legumes to improve soil fertility (Goss et al., 2013). Modern organic amendments include a variety of materials such as compost, animal manure, peat moss, wood chips, straw, sewage sludge, and sawdust, which can be categorized into five main types (Goss et al., 2013).

2.1 Types of Organic Amendments

2.1.1 Animal manure

Animal manure, composed of a mixture of animal feces, urine, and bedding material, has long been used to improve soil fertility. Manure is derived from various livestock, including cattle, poultry, and pigs, and its nutrient content depends on the type of animal, the duration of storage, and whether the manure has been composted. Manure provides not only essential nutrients for plants but also organic matter that enhances soil structure and fertility. During aerobic decomposition, manure releases carbon dioxide and more stable organic compounds, while anaerobic conditions can produce methane and volatile organic acids (Goss et al., 2013). Animal manure is a primary source of organic fertilizer in agriculture, predominantly sourced from poultry, pig, cattle, and sheep farms (Kobayashi et al., 2009). With increasing food demand and shifts in human lifestyle and dietary habits, animal production has expanded significantly in recent years. This growth in the production of meat, eggs, dairy, and leather has introduced several new environmental challenges (Tahmasbian et al., 2017).



Fig. 1. Illustration of organic amendment types
Source: Author's Creation

2.1.2 Municipal biosolids

Changing lifestyles have led to a global increase in wastewater production. As a result, biosolids which are residual byproducts from the treatment of municipal wastewater are also commonly applied to soil as fertilizer (Melo et al., 2018; Eid et al., 2019). Municipal biosolids, or treated sewage sludge, are organic materials processed to reduce odors and pathogens, making them suitable for agricultural use. Biosolids are rich in nutrients and organic matter but are subject to strict regulations to control levels of heavy metals, pathogens, and other contaminants before being applied to soils (Goss et al., 2013).

2.1.3 Green manure and cover crops

Green manure involves growing specific crops and incorporating them into the soil while still green, which boosts soil fertility by adding organic matter and nutrients. Cover crops, such as legumes or non-leguminous species like sorghum and buckwheat, are used to protect soil from erosion and improve soil health during fallow periods or between crop rotations. Legumes are particularly beneficial as they fix nitrogen from the atmosphere, enriching the soil with both nitrogen and organic matter (Goss et al., 2013). Cover crops can also increase microbial activity, improve water retention, and reduce erosion during off-season periods (Sullivan, 2003).

2.2 Compost and Waste from Manufacturing Processes

In recent years, compost has gained attention as an alternative to synthetic fungicides for controlling phytopathogens. Research shows that the increase in microbial biomass following compost application can lead to general disease suppression (Siddiqui et al., 2008). Composting is widely regarded as the most common method for recovering materials from organic waste. Composts are composed of a stable mixture of decayed organo-mineral materials derived from sources such as food scraps, green waste, animal manure, and sewage sludge (Baldantoni et al., 2017). In Europe, nearly half of the compost produced is applied to agricultural land (Latifah et al., 2018). Beyond enhancing soil organic carbon content, compost application has been shown to increase the abundance, activity, and biodiversity of soil microbiota (Gandolfi et al., 2010). Additionally, compost made from crop residues and green waste—such as leaves,

stalks, and stubble—has been found to improve soil water retention, significantly alter microbial communities, and help suppress plant diseases (Feng et al., 2014). However, there are potential downsides to compost use. High electrical conductivity and a high carbon-to-nitrogen (C/N) ratio can lead to soil salinity and nitrogen immobilization. Compost derived from the organic fraction of municipal waste often contains elevated levels of heavy metals, which can accumulate in the soil with repeated use, posing environmental risks (Kodesova et al., 2012). The presence of both organic and inorganic contaminants in compost, depending on its source, can therefore threaten agricultural systems (Kobayashi et al., 2009). While decades of research on composting have yielded extensive knowledge about the process and its effects on soil (Woignier et al., 2016), concerns about contaminants remain.

Meta-analyses have also demonstrated varying degrees of effectiveness in disease suppression depending on the type of phytopathogen (Kavroulakis et al., 2010). However, the wide variability in compost quality, maturity, organic matter content, and microbial composition often results in inconsistent outcomes, making it difficult to identify specific indicators of pathogen suppression (Milinković et al., 2019). Compost is one of the most widely used organic amendments and is created through the controlled decomposition of organic waste materials. It provides multiple benefits, including the recycling of nutrients, improving soil organic carbon content, enhancing microbial activity, and supplying essential nutrients such as nitrogen, phosphorus, potassium, and magnesium (Scotti et al., 2015). Compost also improves soil porosity, water retention, and cation exchange capacity (CEC), while promoting the suppression of soil-borne pathogens such as *Pythium*, *Phytophthora*, and *Fusarium* spp. (Borrero et al., 2004). In recent years, on-farm composting has gained popularity as an environmentally friendly and cost-effective method for recycling agricultural waste into high-quality soil amendments (Pane et al., 2015). This practice not only reduces waste disposal costs but also provides a sustainable source of compost for farmers, contributing to soil health and fertility. On the other hand, organic by-products from various manufacturing industries, such as seed residues, animal feathers, biochar, and paper mill biosolids, are increasingly being used as soil amendments. However, these materials are used less frequently in agriculture due to limited

knowledge about their decomposition rates and nutrient availability (Goss et al., 2013).

3. SOURCES OF ORGANIC AMENDMENTS FOR SOIL RESTORATION

A wide range of organic amendments is available for soil restoration, originating from agriculture, forestry, and urban waste streams. These materials provide essential organic matter and nutrients that help rehabilitate degraded soils, improving both their fertility and structure. Agricultural sources of organic amendments are among the most prevalent and include livestock manure whether fresh, composted, or derived from solid fractions of anaerobic digesters produced by cattle, hogs, and poultry. Crop residues, such as straw and leguminous plants, and spent mushroom compost are also commonly used. These amendments supply organic matter and nutrients that are vital for restoring soil health. Forestry also contributes organic amendments, including wood chips, shavings, and deinking sludge. In the past, wood residuals from the lumber industry were often incinerated to produce wood ash, which was also used as a soil amendment (Charmley et al., 2006). However, these forestry by-products are particularly useful for improving soil structure and increasing organic matter content in degraded lands.

Urban waste streams provide additional sources of organic amendments, such as biosolids (sewage sludge and municipal sludge) produced by wastewater treatment plants. The biodegradable fraction of municipal solid waste, which includes food scraps, yard waste, and paper products, is increasingly used in land restoration as well. These materials, previously destined for landfills, are now being diverted for agricultural use due to concerns over diminishing landfill space (MacLeod et al., 2006). The food processing industry, which handles vegetables, grains, meat, and fish, generates significant amounts of organic by-products that could be used for soil restoration, although they are not yet widely utilized in this context (MacLeod et al., 2006). Historically, organic amendments from urban areas, such as biosolids and food processing waste, were sent to landfills, but this practice has decreased in favor of land application as environmental concerns over landfill space have grown (Edwards & Someshwar, 2000).

4. EFFECTS OF ORGANIC AMENDMENTS ON SOIL PROPERTIES

Organic amendments significantly impact soil properties, both directly through their inherent characteristics and indirectly by modifying the soil's physical, biological, and chemical attributes. These amendments play a crucial role in soil restoration, particularly in degraded environments, where they enhance soil health, nutrient availability, and structure. Organic matter is essential for soil health and plays a critical role in maintaining overall environmental quality (Mohan et al., 2018). Intensive agricultural practices, when not supplemented with organic amendments to restore soil organic carbon, along with industrial pollution, can adversely affect the soil's chemical properties. This depletion of soil carbon content has harmful effects on microbial activity, diversity, and enzymatic functions (Rahman et al., 2016). Additionally, these practices can lead to a significant increase in soil salinity (Nawab et al., 2016). It is well established that adding organic amendments to soil increases soil organic matter, which in turn enhances water-holding capacity, porosity, and aggregate stability—key factors in improving soil quality (Beesley et al., 2010). Additionally, applying other waste materials such as rice husk, wheat straw, reeds, and biochar, alongside composts and manures, has been shown to improve soil aggregate stability and reduce bulk density (Nawab et al., 2016). Since organic fertilizers have a much higher water-holding capacity compared to degraded soils, their application can significantly boost the soil's water retention. This mixed approach to soil supplementation can amplify the benefits of organic amendments (Puglisi et al., 2007). However, the overall impact and long-term effects depend heavily on the type and number of organic amendments used (Houben et al., 2012).

4.1 Soil Organic Matter Content

Soil organic matter is considered one of the most vital components influencing soil quality and functionality (Gregorich et al., 1994). The application of organic amendments typically results in an immediate increase in soil organic carbon, with the amount of carbon added often proportional to the quantity of organic matter applied (Chantigny et al., 1999). However, detecting measurable changes in soil organic carbon can be challenging, especially in soils

with high background organic carbon or variability (Viaud et al., 2011). The long-term retention of soil organic carbon depends on the rate of decomposition and the intrinsic quality of the organic amendment (Lashermes et al., 2009). Carbon introduced into the soil through organic amendments originates from plant material, with its stability influenced by factors such as plant type and post-harvest processing. For example, cereal straw incorporated into soil as part of manure decomposes differently than fresh straw, as composting stabilizes the carbon, leading to more durable improvements in soil organic matter (Lashermes et al., 2009).

4.2 Soil Microbial Communities

It is widely understood that after adding organic amendments to soil, the introduced microbial communities usually give way to the native soil microbes, except for some persistent pathogens (Anastasi et al., 2009). The changes in soil microbial communities that follow are mainly linked to the shifts in the soil's physical and chemical properties caused by the amendments (Tahmasbian et al., 2017). Organic amendments can alter the composition, biomass, and activity of native microbes by providing them with more nutrients for growth (Gandolfi et al., 2010). However, research on this topic often shows site-specific results. For instance, some studies have found no changes in microbial communities in soils contaminated with lead after adding organic materials like poultry litter, sheep manure, paper mill sludge, or cow slurry, possibly due to the severe contamination (Chen et al., 2018). On the other hand, other studies have shown beneficial effects of organic amendments on native microbiota in agricultural soils (Tahmasbian et al., 2017). In degraded soils, adding organic amendments has been found to boost soil life both in terms of quantity and biodiversity (Hashimoto et al., 2009). For example, animal manure has been shown to increase earthworm populations compared to soils only treated with synthetic fertilizers, likely due to the added carbon and improved water retention from organic fertilizers (Kang et al., 2016). Organic amendments positively influence soil biota, enhancing soil biochemical activities such as enzymatic function and overall biodiversity (Bastida et al., 2008). Studies have shown that organic amendments, particularly animal manure, increase the abundance of earthworms compared to inorganic fertilizers (Leroy et al., 2008). The composition of the amendment also matters; farmyard manure and cattle slurry, for

instance, tend to support higher earthworm populations than composts, due to the greater availability of carbon in manure. Similarly, microbial communities benefit from organic amendments, as they provide a significant carbon source that stimulates microbial growth and activity. This is often measured by the increase in soil microbial biomass carbon following the addition of organic amendments (Belyaeva & Haynes, 2009). The level of microbial activity, however, varies depending on the decomposability and rate of application of the amendments (Albiach et al., 2000).

4.3 Biological Aggregation and Soil Restoration

Organic amendments initiate soil biological activity, which plays a central role in soil reclamation. This process takes place in the detritosphere, a localized area around decomposing organic residues, where intense microbial activity occurs (Gaillard et al., 1999). Studies have shown that decomposing plant residues significantly alter the physical structure of soil, promoting the formation of stable soil aggregates. As organic residues are colonized by microbes, they bind with mineral particles, creating stable aggregates that protect organic matter from rapid decomposition (Golchin et al., 1994). The application of organic amendments such as wood-derived sludge, compost, and cattle manure has been shown to promote the formation of soil aggregates, stabilizing soil organic carbon and contributing to the early stages of soil formation in highly degraded areas (Sere et al., 2010).

4.4 Nitrogen and Phosphorus Availability

Organic amendments or fertilizers contain varying amounts of nitrogen and phosphorus in different forms (Purnomo et al., 2010). Organic amendments contain nitrogen in varying concentrations, with some forms more readily available than others (Lashermes et al., 2010). Nitrogen is typically found in an organic form, but in animal manures especially liquid ones most of the nitrogen is in a mineral form that is readily available for plants to absorb (Baldantoni et al., 2017). When nitrogen is present in its organic form, it must undergo a process called mineralization to be converted into a form that plants can use (Zhang et al., 2018). One of the main advantages of using organic amendments instead of synthetic fertilizers is the different ways in which phosphorus becomes available.

Studies have shown that applying animal manure is often more effective than using inorganic phosphorus fertilizers, leading to greater phosphorus availability. In contrast, the increase in phosphorus from synthetic fertilizers is often minimal, even with high application rates (Adam et al., 2015; Naser et al., 2018). In liquid animal manures, nitrogen is present in a mineral form (ammonia), which plants can absorb immediately (Chantigny et al., 2001). However, in most

organic amendments, nitrogen is in an organic form, requiring mineralization to convert it into a plant-available form. Low-nitrogen amendments may even immobilize nitrogen, competing with plants for available nitrogen. Predicting nitrogen availability from organic amendments depends on their chemical composition, including factors such as soluble cellulose and lignin-like fractions, which influence the rate of nitrogen mineralization (Lashermes et al., 2010).

Table 1. Positive impacts of soil organic amendments

Positive Impact	Details/Description	References
Increases Soil Organic Matter	Organic amendments increase soil organic carbon, which improves overall soil quality and functioning.	Gregorich et al., 1994; Chantigny et al., 1999; Lashermes et al., 2009
Enhances Soil Structure	Organic matter improves poor soil physical conditions, leading to better structure and reduced compaction.	Viaud et al., 2011; Golchin et al., 1994; Sere et al., 2010
Boosts Microbial Activity	Organic amendments provide carbon, which serves as an energy source for soil microbes, increasing microbial biomass.	Viaud et al., 2011; Belyaeva & Haynes, 2009
Increases Soil Biota	Organic amendments, particularly animal manure, increase earthworm abundance and soil biodiversity.	Bastida et al., 2008; Leroy et al., 2008
Promotes Biological Aggregation	Induces biological activity that forms stable soil aggregates, protecting organic matter and contributing to pedogenesis.	Gaillard et al., 1999; Golchin et al., 1994; Sere et al., 2010
Improves Nitrogen Availability	Organic amendments can provide slow-release nitrogen, reducing the risk of leaching and enhancing plant uptake.	Chantigny et al., 2001; Lashermes et al., 2010
Increases Water-Holding Capacity	Organic matter retains more water, enhancing water-holding capacity, particularly in sandy and degraded soils.	Camberato et al., 2006; Fierro et al., 1999; Larney et al., 2000
Suppresses Soil-Borne Pathogens	Compost amendments have been shown to suppress diseases caused by soil pathogens like <i>Pythium</i> and <i>Phytophthora</i> .	Borrero et al., 2004; Donn et al., 2014
Enhances Cation Exchange Capacity (CEC)	Compost improves soil CEC, increasing the soil's ability to retain and exchange nutrients.	Scotti et al., 2013
Promotes Long-Term Carbon Stability	Composted materials stabilize carbon, resulting in longer-lasting increases in soil organic matter.	Lashermes et al., 2009
Reduces Heavy Metal Toxicity	Organic matter can help mitigate the harmful effects of heavy metals in soil.	Park et al., 2011

Source: Author's Compilation

4.5 Water-Holding Capacity and Plant-Available Water

Organic amendments typically increase the water-holding capacity of soils, particularly in sandy, degraded soils that lack this essential property (Camberato et al., 2006). The organic matter in these amendments absorbs and retains water, providing more plant-available water. The extent of this benefit depends on the type and decomposition state of the organic amendment. While some studies have shown significant increases in water retention following the addition of organic matter, the effects can vary depending on the soil texture and other factors (Fierro et al., 1999). For instance, biosolids have been found to improve water retention in fine-textured soils such as silt loams but have less effect on sandy soils, where decreased bulk density may offset water retention gains (Gardner et al., 2010). Similar results have been observed with manure applications, which tend to increase water retention primarily due to the water absorbed by the organic matter itself (Larney et al., 2000).

In general, organic amendments play a critical role in improving soil properties, from increasing organic matter and microbial activity to enhancing water retention and nutrient availability. These changes support soil restoration, particularly in degraded environments, by creating conditions that promote long-term soil health and productivity. More so, organic amendments for soil restoration come from diverse sources, including agriculture, forestry, and urban waste. Each type of amendment offers unique benefits for improving soil quality, adding organic matter, and enhancing nutrient availability, all of which are critical for the successful rehabilitation of degraded soils.

5. POTENTIAL RISKS OF ORGANIC AMENDMENTS

1. Application of sewage sludge and animal manures can introduce pathogenic microorganisms (e.g., *Escherichia coli*, *Salmonella*, *Campylobacter*, *Yersinia*), posing risks to soil health (Watteau et al., 2018).
2. Organic amendments may contain heavy metals, which can accumulate in the soil.
3. The application of sewage sludge contributes significantly to microplastic contamination in soils (120 to 860 tons annually in Europe) (Scheurer et al., 2018).

Microplastics and nanoplastics have toxic effects on various organisms, including microbial communities, earthworms, birds, and mammals, raising concerns about ecosystem health (Watteau et al., 2018; Xu et al., 2019).

4. Organic amendments decompose over time, releasing bound contaminants such as metals and micro/nanoplastics, potentially affecting plant growth (Imran et al., 2019).
5. Contaminants from organic amendments can transfer through the food chain, impacting animal and human health (Imran et al., 2019).

6. CONCLUSION

Soil organic matter (SOM) is vital for soil health, affecting its physical, chemical, and biological properties. Effective SOM management supports soil fertility, sustainable agriculture, and ecosystem health. However, practices like continuous cropping and poor soil management deplete soil organic carbon (SOC), impacting soil structure, water retention, and productivity. Organic amendments such as manure, compost, and biosolids help restore SOC, improve soil structure, boost microbial activity, and enhance nutrient cycling. They also mitigate contamination risks but must be used cautiously to avoid releasing harmful substances. Integrating organic amendments with sustainable practices like conservation tillage and crop rotations is key to restoring soil health and ensuring long-term agricultural sustainability.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Abiven, S., Menassero, S., and Chenu, C. 2009. The effect of organic inputs over time on soil aggregate stability - a literature analysis. *Soil Biological Biochem.* 41, 1-12.

- Adam, I.K.U., Miltner, A., & Kästner, M. (2015). Degradation of 13C-labeled pyrene in soil-compost mixtures and fertilized soil. *Appl Microbiol Biotechnol*, 99, 9813–9824 <https://doi.org/10.1007/s00253-015-6822-8>
- Akala, V. A. and Lal, R. 2000. Potential of mine land reclamation for soil organic carbon sequestration in Ohio. *Land Degradation Development* 11: 289_297.
- Albiach, R., Canet, R., Pomares, F. and Ingelmo, F. 2000. Microbial biomass content and enzymatic activities after the application of organic amendments to a horticultural soil. *Bioresour. Technol.* 75: 43_48.
- Anastasi, A., Coppola, T., Prigione, V., Varese, GC. (2009). Pyrene degradation and detoxification in soil by a consortium of basidiomycetes isolated from compost: Role of laccases and peroxidases, *Journal of Hazardous Materials*, 165, 1–3, 1229-1233.
- Baldantoni, D., Morelli, R., Bellino, A., Prati, M. V., Alfani, A., & Nicola, F. D. (2017). Anthracene and benzo(a)pyrene degradation in soil is favoured by compost amendment: Perspectives for a bioremediation approach. *Journal of Hazardous Materials*, 339, 395–400. doi: 10.1016/j.jhazmat.2017.06.043
- Bastida, F., Kandeler, E., Hernandez, T. and Garcia, C. 2008. Long-term effect of municipal solid waste amendment on microbial abundance and humus-associated enzyme activities under semiarid conditions. *Microbial Ecology* 55: 651_661.
- Beesley, L., Moreno-Jiménez, E., & Gomez-Eyles, J. L. (2010). Effects of biochar and greenwaste compost amendments on mobility, bioavailability and toxicity of inorganic and organic contaminants in a multi-element polluted soil. *Environmental Pollution*, 158(6), 2282–2287. doi: 10.1016/j.envpol.2010.02.003
- Belyaeva, O. N. and Haynes, R. J. 2009. Chemical, microbial and physical properties of manufactured soils produced by composting municipal green waste with coal fly ash. *Bioresour. Technol.* 100: 5203_5209.
- Borrero, C., Trillas, M. I., Ordovás, J., Tello, J. C., and Avilés, M. 2004. Predictive factors for the suppression of *Fusarium* wilt of tomato in plant growth media. *Phytopathology*. 94, 1094-1101.
- Camberato, J. J., Gagnon, B., Angers, D. A., Chantigny, M. H. and Pan, W. L. 2006. Pulp and papermill by-products as soil amendments and plant nutrient sources. *Can. J. Soil Sci.* 86: 641_653.
- Chantigny, M. H., Angers, D. A. and Beauchamp, C. J. 1999. Aggregation and organic matter decomposition in soils amended with de-inking paper sludge. *Soil Science Society American Journal* 63: 1214_1221.
- Chantigny, M. H., Rochette, P. and Angers, D. A. 2001. Shortterm C and N dynamics in a soil amended with pig slurry and barley straw: A field experiment. *Canadian Journal Soil Sci.* 81: 131_137.
- Charmley, E., Nelson, D. and Zvomuya, F. 2006. Nutrient cycling in the vegetable processing industry: Utilization of potato by-products. *Can. J. Soil Sci.* 86: 621_629.
- Chen, C., Guron, G. K., Pruden, A., Ponder, M., Du, P., & Xia, K. (2018). Antibiotics and Antibiotic Resistance Genes in Bulk and Rhizosphere Soils Subject to Manure Amendment and Vegetable Cultivation. *Journal of Environmental Quality*, 47(6), 1318–1326. doi: 10.2134/jeq2018.02.0078
- Donn, S., Wheatley, R. E., McKenzie, B. M., Loades, K. W., and Hallett, P. D. 2014. Improved soil fertility from compost amendment increases root growth and reinforcement of surface soil on slope. *Ecological Engineering* 71, 458-465.
- Edwards, J. H. and Someshwar, A. V. 2000. Chemical, physical, and biological characteristics of agricultural and forest by-products for land application. Pages 1_62 in J. F. Power and W. A. Dick, eds. *Land application of agricultural, industrial and municipal by-products*. SSSA Book Series No. 6, SSSA, Madison, WI.
- Eid, E. M., Hussain, A. A., Taher, M. A., Galal, T. M., Shaltout, K. H., & Sewelam, N. (2019). Sewage Sludge Application Enhances the Growth of *Corchorus olitorius* Plants and Provides a Sustainable Practice for Nutrient Recirculation in Agricultural Soils. *Journal of Soil Science and Plant Nutrition*, 20(1), 149–159. doi: 10.1007/s42729-019-00113-z
- Ellis, E. C., Klein Goldewijk, K., Siebert, S., Lightman, D. and Ramankutty, N. 2010. Anthropogenic transformation of the

- biomes, 1700 to 2000. *Global Ecol. Biogeogr.* 19: 589_606.
- Feng, L., Zhang, L., Feng, L. (2014). Dissipation of polycyclic aromatic hydrocarbons in soil amended with sewage sludge compost, *International Biodeterioration & Biodegradation*, 95, Part A, 200-207.
- Fierro, A., Angers, D. A. and Beauchamp, C. J. 1999. Restoration of ecosystem function in an abandoned sandpit: plant and soil responses to paper de-inking sludge. *J. Applied Ecology* 36: 244_253.
- Gaillard, V., Chenu, C. and Recous, S. 2003. Carbon mineralization in soil adjacent to plant residues of contrasting biochemical quality. *Soil Biol. Biochem.* 35: 93_99.
- Gaillard, V., Chenu, C., Recous, S. and Richard, G. 1999. Carbon, nitrogen and microbial gradients induced by plant residues decomposing in soil. *Eur. J. Soil Sci.* 50: 567_578.
- Gandolfi, I., Sicolo, M., Franzetti, A., Fontanarosa, E., Santagostino, A., Bestetti, G. (2010). Influence of compost amendment on microbial community and ecotoxicity of hydrocarbon-contaminated soils, *Bioresource Technology*, 101, 2, 568-575.
- Gardner, W. C., Broersma, K., Naeth, A., Chanasyk, D. and Jobson, A. 2010. Influence of biosolids and fertilizer amendments on physical, chemical and microbiological properties of copper mine tailings. *Can. J. Soil Sci.* 90: 571_583.
- Golchin, A., Oades, J. M., Skjemstad, J. O. and Clarke, P. 1994. Soil structure and carbon cycling. *Aust. J. Soil Res.* 32: 1043_1068.
- Goss, M. J., Tubeileh, A., and Goorahoo, D. 2013. A review of the use of organic amendments and the risk to human health. *Adv. Agron.* 120, 275-379.
- Gregorich, E. G., Carter, M. R., Angers, D. A., Monreal, C. M. and Ellert, B. H. 1994. Towards a minimum data set to assess soil organic matter quality in agricultural soils. *Can. J. Soil Sci.* 74: 367_385.
- Gupta, S. C., Dowdy, R. H. and Larson, W. E. 1977. Hydraulic and thermal properties of a sandy soil as influenced by incorporation of sewage sludge. *Soil Sci. Soc. Am. J.* 41: 601_605.
- Hashimoto, Y., Taki, T., & Sato, T. (2009). Extractability and leachability of Pb in a shooting range soil amended with poultry litter ash: Investigations for immobilization potentials. *Journal of Environmental Science and Health, Part A*, 44(6), 583–590. doi: 10.1080/10934520902784617
- Houben, D., Pircar, J., & Sonnet, P. (2012). Heavy metal immobilization by cost-effective amendments in a contaminated soil: Effects on metal leaching and phytoavailability. *Journal of Geochemical Exploration*, 123, 87–94. doi: 10.1016/j.gexplo.2011.10.004
- Imran, M., Das, K. R., & Naik, M. M. (2019). Co-selection of multi-antibiotic resistance in bacterial pathogens in metal and microplastic contaminated environments: An emerging health threat. *Chemosphere*, 215, 846–857. doi: 10.1016/j.chemosphere.2018.10.114
- Kang, Y., Hao, Y., Shen, M., Zhao, Q., Li, Q., & Hu, J. (2016). Impacts of supplementing chemical fertilizers with organic fertilizers manufactured using pig manure as a substrate on the spread of tetracycline resistance genes in soil. *Ecotoxicology and Environmental Safety*, 130, 279–288. doi: 10.1016/j.ecoenv.2016.04.028
- Kavroulakis, N., Ntougias, S., Besi, M. I., Katsou, P., Damaskinou, A., Ehaliotis, C., ... Papadopoulou, K. K. (2010). Antagonistic bacteria of composted agro-industrial residues exhibit antibiosis against soil-borne fungal plant pathogens and protection of tomato plants from *Fusarium oxysporum* f.sp. *radicis-lycopersici*. *Plant and Soil*, 333(1-2), 233–247. doi: 10.1007/s11104-010-0338-x
- Kobayashi, T., Murai, Y., Tatsumi, K., & Imura, Y. (2009). Biodegradation of polycyclic aromatic hydrocarbons by *Sphingomonas* sp. enhanced by water-extractable organic matter from manure compost. *Science of The Total Environment*, 407(22), 5805–5810. doi: 10.1016/j.scitotenv.2009.06.041
- Kodešová, R., Kočárek, M., Hajková, T., Hýbler, M., Drábek, O., & Kodeš, V. (2012). Chlorotoluron mobility in compost amended soil. *Soil and Tillage Research*, 118, 88–96. doi: 10.1016/j.still.2011.10.014
- Larney, F. J. and Pan, W. L. 2006. Organic waste to resource: recycling nutrients. *Can. J. Soil Sci.* 86: 585_586.
- Larney, F. J., Hao, X. and Topp, E. 2011. Manure management. Pages 247_263 in J. L. Hatfield and T. J. Sauer, eds. *Soil*

- management: building a stable base for agriculture. ASA, SSSA, Madison, WI.
- Larney, F. J., Janzen, H. H., Olson, B. M. and Lindwall, C. W. 2000. Soil quality and productivity responses to simulated erosion and restorative amendments. *Can. J. Soil Sci.* 80: 515_522.
- Lashermes, G., Nicolardot, B., Parnaudeau, V., Thuriès, L., Chaussod, R., Guillotin, M. L., Line`res, M., Mary, B., Metzger, L., Morvan, T., Tricaud, A., Vilette, C. and Houot, S. 2009. Indicator of potential residual carbon in soils after exogenous organic matter application. *Eur. J. Soil Sci.* 60: 297_310.
- Latifah, O., Ahmed, O.H., Majid, N.M.A. (2018). Soil pH buffering capacity and nitrogen availability following compost application in a tropical acid soil. *Compost Sci Util.* 26(1), 1–15.
- Leroy, B. L. M., Herath, H. M. S. K., Sleutel, S., De Neve, S., Gabriels, D., Reheul, D. and Moens, M. 2008. The quality of exogenous organic matter: Short-term effects on soil physical properties and soil organic matter fractions. *Soil Use Manage.* 24: 139_147.
- MacLeod, J. A., Kuo, S., Gallant, T. L. and Grimmett, M. 2006. Seafood processing wastes as nutrient sources for crop production. *Can. J. Soil Sci.* 86: 631_640.
- Melo, W., Delarica, D., Guedes, A., Lavezzo, L., Donha, R., Araújo, A. D., ... Macedo, F. (2018). Ten years of application of sewage sludge on tropical soil. A balance sheet on agricultural crops and environmental quality. *Science of The Total Environment*, 643, 1493–1501. doi: 10.1016/j.scitotenv.2018.06.254
- Milinković, M., Lalević, B., Jovičić-Petrović, J., Golubović-Čurguz, V., Kljujev, I., & Raičević, V. (2019). Biopotential of compost and compost products derived from horticultural waste—Effect on plant growth and plant pathogens suppression. *Process Safety and Environmental Protection*, 121, 299–306. doi: 10.1016/j.psep.2018.09.024
- Mohan, D., Abhishek, K., Sarswat, A., Patel, M., Singh, P., Pittman, C.U. (2018). Biochar production and applications in soil fertility and carbon sequestration—a sustainable solution to crop-residue burning in India. *RSC Adv*, 8(1), 508–520
- Montgomery, D. R. 2007. *Dirt—the erosion of civilizations*. University of California Press, Berkeley, CA. 60: 297_310
- Naser, H., Rahman, M., Sultana, S., Quddus, M., & Haoque, M. (2018). Remediation of heavy metal polluted soil through organic amendments. *Bangladesh Journal of Agricultural Research*, 42(4), 589. doi:10.3329/bjar.v42i4.35786
- Nawab, J., Khan, S., Aamir, M., Shamshad, I., Qamar, Z., Din, I., Huang, Q. (2016). Organic amendments impact the availability of heavy metal (loid) s in mine-impacted soil and their phytoremediation by *Penisitum americanum* and *Sorghum bicolor*. *Environ. Sci. Pollut. Res.* 23 (3), 2381–2390.
- Pane, C., Celano, G., Piccolo, A., Vilecco, D., Spaccini, R., Palese, A.M., Zaccardelli, M. 2015. Effects of on-farm composted tomato residues on soil biological activity and yields in a tomato cropping system. *Chemical Biological Technol. Agric.* 2, 4.
- Park, J. H., Lamb, D., Paneerselvam, P., Choppala, G., Bolan, N., and Chung, J. W. 2011. Role of organic amendments on enhanced bioremediation of heavy metal(loid) contaminated soils. *J. Hazard. Mater.* 185, 549-574.
- Puglisi, E., Cappa, F., Fragoulis, G., Trevisan, M., & Re, A. A. D. (2007). Bioavailability and degradation of phenanthrene in compost amended soils. *Chemosphere*, 67(3), 548–556. doi: 10.1016/j.chemosphere.2006.09.058
- Purnomo, A. S., Koyama, F., Mori, T., & Kondo, R. (2010). DDT degradation potential of cattle manure compost. *Chemosphere*, 80(6), 619–624. doi: 10.1016/j.chemosphere.2010.04.059
- Rahman, F., Rahman, M.M., Rahman, G.K.M.M., Saleque, M.A., Hossain, A.S., Miah, M.G. (2016). Effect of organic and inorganic fertilizers and rice straw on carbon sequestration and soil fertility under a rice–rice cropping pattern. *Carbon Manage*, 7(1–2), 41–53.
- Scheurer, M., & Bigalke, M. (2018). Microplastics in Swiss Floodplain Soils. *Environmental Science & Technology*, 52(6), 3591–3598. doi: 10.1021/acs.est.7b06003
- Scotti, R., Conte, P., Berns, A. E., Alonzo, G., and Rao, M. A. 2013. Effect of organic amendments on the evolution of soil organic matter in soils stressed by

- intensive agricultural practices. *Curr. Org. Chem.* 17, 2998-3005.
- Scotti, R., D'Ascoli, R., Bonanomi, G., Caceres, M. G., Sultana, S., Cozzolino, L., Scelza, R., Zoina, A., Rao, and M. A. 2015. Combined use of compost and wood scraps to increase carbon stock and improve soil quality in intensive farming systems. *Eur. J. Soil Sci.* doi: 10.1111/ejss.12248
- Sere, G., Schwartz, C., Ouvrard, S., Renat, J.-C., Watteau, F., Villemin, G. and Morel, J. L. 2010. Early pedogenic evolution of constructed Technosols. *J. Soils Sediments* 10: 1246_1254.
- Siddiqui, Y., Meon, S., Ismail, R., Rahmani, M., & Ali, A. (2008). Bio-efficiency of compost extracts on the wet rot incidence, morphological and physiological growth of okra (*Abelmoschus esculentus* [(L.) Moench]). *Scientia Horticulturae*, 117(1), 9–14. doi: 10.1016/j.scienta.2008.03.008
- Stewart, B. A., Robinson, C. A. and Parker, D. B. 2000. Examples and case studies of beneficial reuse of beef cattle byproducts. Pages 387_407 in J. F. Power and W. A. Dick, eds. *Land application of agricultural, industrial and municipal byproducts*. SSSA Book Series No. 6, SSSA, Madison, WI.
- Sullivan, P. 2003. Overview of cover crops and green manures: fundamentals of sustainable agriculture.
- Tahmasbian, I., Safari Sinangani, A.A., Nguyen, T.T.N. (2017). Application of manures to mitigate the harmful effects of electrokinetic remediation of heavy metals on soil microbial properties in polluted soils. *Environ Sci Pollut Res* 24, 26485–26496. <https://doi.org/10.1007/s11356-017-0281-y>
- Viaud, V., Angers, D. A., Parnaudeau, V., Morvan, T. and MenasseriAubry, S. 2011. Response of organic matter to reduced tillage and animal manure in a temperate loamy soil. *Soil Use Manage.* 27: 84_93.
- Watteau, F., Dignac, M.-F., Bouchard, A., Revallier, A., & Houot, S. (2018). Microplastic Detection in Soil Amended With Municipal Solid Waste Composts as Revealed by Transmission Electronic Microscopy and Pyrolysis/GC/MS. *Frontiers in Sustainable Food Systems*, 2. doi: 10.3389/fsufs.2018.00081
- Woignier, T., Clostre, F., Fernandes, P. (2016). Compost addition reduces porosity and chlordecone transfer in soil microstructure. *Environ Sci Pollut Res* 23, 98–108.
- Xu, B., Liu, F., Cryder, Z., Huang, D., Lu, Z., He, Y., ... Xu, J. (2019). Microplastics in the soil environment: Occurrence, risks, interactions and fate – A review. *Critical Reviews in Environmental Science and Technology*, 1–48. doi: 10.1080/10643389.2019.1694822
- Zaccardelli, M., De Nicola, F., Villecco, D., and Scotti, R. 2013. The development and suppressive activity of soil microbial communities under compost amendment. *J. Soil Sci. Plant Nutr.* 13, 730-742.
- Zhang, J., Sui, Q., Tong, J., Zhong, H., Wang, Y., Chen, M., & Wei, Y. (2018). Soil types influence the fate of antibiotic-resistant bacteria and antibiotic resistance genes following the land application of sludge composts. *Environment International*, 118, 34–43. doi: 10.1016/j.envint.2018.05.029
- Zika, M. and Erb, K. H. 2009. The global loss of net primary production resulting from human-induced soil degradation in drylands. *Ecol. Econ.* 69: 310_318.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/125606>