



# **Analgesic Effects of Methanol Leaf Extracts from *Datura alba ness* in Male Wistar Rats through Reduction of C-reactive Protein Levels**

**Blessing Zeinab Ovili-Odili <sup>a</sup>,  
Olaghaguo Macstephen Adienbo <sup>b</sup>,  
Tarela Melish Elias Daubry <sup>a</sup>  
and Bruno Chukwuemeka Chinko <sup>b\*</sup>**

<sup>a</sup> Department of Physiology, Faculty of Basic Medical Sciences, Delta State University Abraka, Delta State, Nigeria.

<sup>b</sup> Department of Human Physiology, Faculty of Basic Medical Sciences, University of Port Harcourt, Port Harcourt, Rivers State, Nigeria.

## **Authors' contributions**

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## **Article Information**

DOI: <https://doi.org/10.9734/jocamr/2024/v25i11583>

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/125537>

**Original Research Article**

**Received: 20/08/2024  
Accepted: 22/10/2024  
Published: 30/10/2024**

## **ABSTRACT**

**Introduction:** The increasing interest in utilizing natural products from medicinal plants for pain management stems from their wide availability and minimal side effects. This study aims to evaluate the toxicity profile of *Datura alba ness* (DAN) extracts and assess their potential analgesic

\*Corresponding author: E-mail: [bruno.chinko@uniport.edu.ng](mailto:bruno.chinko@uniport.edu.ng);

**Cite as:** Ovili-Odili, Blessing Zeinab, Olaghaguo Macstephen Adienbo, Tarela Melish Elias Daubry, and Bruno Chukwuemeka Chinko. 2024. "Analgesic Effects of Methanol Leaf Extracts from *Datura Alba Ness* in Male Wistar Rats through Reduction of C-Reactive Protein Levels". *Journal of Complementary and Alternative Medical Research* 25 (11):57-66. <https://doi.org/10.9734/jocamr/2024/v25i11583>.

activity through preventive and therapeutic analgesia models in Wistar rats, as well as to investigate the possible mechanisms of action using molecular docking.

**Materials and Methods:** Fresh leaves of DAN were collected from a garden in the Abua/Odua Local Government Area of Rivers State, Nigeria. The leaves were extracted using cold maceration to obtain ethanolic extracts of *Datura alba ness* (MEDAN). Twelve mice (19-30g) were used for toxicity study, while sixty Wistar rats (140-180g) were used for analgesic studies. Serum C-reactive protein levels were determined using ELISA methods, and molecular docking simulations were conducted using AutoDock and Discovery Studio Visualizer.

**Results:** The results indicated that MEDAN was very safe, with an LD<sub>50</sub> of 5000 mg/kg. The findings demonstrate that MEDAN significantly increased the pain threshold in male Wistar rats after three weeks of administration in a dose-dependent manner during the preemptive analgesia study, while pain thresholds increased within one hour but decreased over time in the therapeutic analgesia study. The combination of MEDAN with a standard drug (Flexicam) resulted in improved analgesic effects at the 500 mg/kg dose. Additionally, the study revealed that MEDAN significantly reduced serum C-reactive protein levels after three weeks of oral administration. Molecular docking studies indicated that stigmasterol, gamma-sitosterol, and cholest-5-en-3-ol, 24-propylidene-(3 $\beta$ ) from MEDAN demonstrated binding affinity to C-reactive protein receptors, mimicking the effects of the reference drug, celecoxib.

**Conclusion:** This study concludes that ethanolic extracts of *Datura alba ness* exhibited significant efficacy in pain reduction by lowering C-reactive protein levels. This effect was especially enhanced when the 500 mg/kg dose of MEDAN was combined with the standard drug, Flexicam. The findings provide a scientific basis supporting the traditional medicinal use of *Datura alba ness* for pain management.

*Keywords: Datura alba ness; analgesia; c-reactive protein; molecular docking.*

## 1. INTRODUCTION

Pain is defined as an "unpleasant sensory or emotional experience associated with actual or potential tissue damage or described in terms of such damage." The pain perception threshold refers to the point at which an individual first perceives pain, and this threshold exhibits minimal variability among individuals [1,2]. Pain signals are transmitted through a complex network of pathways from peripheral receptors to the central nervous system. Primary afferent pain fibres synapse with second-order neurons in the dorsal horn of the spinal cord, with ascending spinothalamic and spinoreticular tracts carrying pain signals to the brain. These signals are processed in the thalamus and relayed to the cortex for interpretation. Pain modulation occurs through descending tracts involving the periaqueductal grey matter and nucleus raphe magnus. Neuropathic pain arises from nerve damage, with proposed mechanisms involving both peripheral and central sensitization pathways [1,3]. Pain is classified based on various criteria, including the affected body region (somatic or visceral pain), duration (acute or chronic), causative factors (neuropathic or inflammatory), and the mechanism of transmission (central, peripheral, or nociceptive) [2-4]. Analgesics are drugs used in pain

management. They exert their effects through various mechanisms which highlight the diverse approaches to pain management. For example, acetaminophen (paracetamol) and non-steroidal anti-inflammatory drugs (NSAIDs) relieve pain by inhibiting cyclooxygenase enzymes (COX) in the brain, reducing pain and fever while opioid receptors act by inhibiting the pain pathway and general modulation and dampening of pain perception [5-7].

Traditional African medicine primarily relies on botanical preparations for the treatment of illnesses in both animals and humans. In Traditional African Herbal Medicine (TAHM), remedies are created by extracting components from various parts of plants, including roots, barks, leaves, flowers, seeds, and aerial parts, either as single entities or through combinations of different parts from various plant species [8,9]. Research aimed at validating traditional knowledge has demonstrated the efficacy of TAHM preparations in alleviating pain. Bioactive compounds have been identified in certain plant species, and further research could lead to the acceptance and integration of some traditional pain treatments into modern medical practices. This exploration may pave the way for discovering new bioactive compounds that could be developed into evidence-based

pharmaceutical medications [10, 11]. Some of the African plants screened for their potential analgesic activity include *Acacia modesta* [12], *Alstonia boonei* [13], *Annona muricata* [14], *Bryophyllum pinnatum* [15], *Chromolaena odorata* [16], *Hymenocardia acida* [17], *Ipomoea involucre* [18], *Myristica Fragrans Houtt* [19], *Tamarindus indica* [20], *Zea mays* and many others [21].

The *Datura* genus is generally characterized as comprising of annual or perennial herbs and shrubs that are part of the Solanaceae family. The leaves have petioles, and the tubular calyx splits around near the base, while the funnel-shaped, elongated corolla features cuspidate lobes [22,23]. The seeds are dark, flat, and kidney-shaped, while the spiny fruits resemble walnuts, giving rise to the name "thorn apple." The plant is also commonly known as Jimsonweed or Devil's trumpet [23,24]. *Datura alba ness* (DAN) is one of the most prevalent species in Nigeria. It is commonly known as *Zakami*, while the Ogoni and Ndokwa tribes refer to it as *Jegemi* and *Gegemu*, respectively [25]. Despite its toxic properties, *Datura* has long been used in traditional medicine as an anaesthetic and for treating bruises, wounds, and skin ulcers [25,26]. Our earlier report on the proximate, phytochemical and bioactive analysis of *Datura alba ness* leaf extract showed it contains various bioactive compounds with potential medicinal properties [25]. *Datura* species, *Datura fastuosa* and *Datura metel*, have been shown to exhibit strong analgesic properties [27,28]. However, no research has yet explored the potential analgesic effects of *Datura alba ness* (DAN), despite anecdotal evidence suggesting that brewing its leaves in local gin may help manage pain. This study aims to evaluate the potential analgesic activity of DAN extracts using preventive and therapeutic analgesia models in Wistar rats and the possible mechanism of action using molecular docking.

## 2. MATERIALS AND METHODS

### 2.1 Collection, Identification, Preparation and Extraction of Plant Material

Fresh leaves of *Datura alba ness* were collected from a garden in the Abua/Odua Local Government Area of Rivers State, Nigeria. The plant was identified and authenticated by a taxonomist from the Department of Plant Science and Biotechnology at the University of Port Harcourt and assigned the voucher number

UPH/295. The leaves were washed and air-dried at room temperature for two weeks until they became crisp. They were then ground into coarse particles. The extraction process followed a cold maceration technique as previously described [29,30]. Approximately 400 g of the coarsely ground *Datura alba ness* leaves were soaked in 2000 ml of methanol and left to stand for 72 hours with occasional stirring. The macerated mixture was then filtered using filter paper, and the filtrate was evaporated in a water bath at 50°C to yield the methanol extract of *Datura alba ness* (MEDAN).

### 2.2 Research Animals

Twelve (12) mice weighing between 19 and 30 grams, along with sixty (60) Wistar rats weighing between 140 and 180 grams, were obtained from the animal house at the Faculty of Basic Medical Sciences, University of Port Harcourt, for the study. All animal experiments adhered to the principles of laboratory animal care [31]. The animals were given a 14-day acclimatization period at the experimental site before the study began. They were housed in wooden cages under standard conditions, with a temperature of 22±3°C, 12 hours of light, and 12 hours of darkness. Throughout the study, they had access to standard laboratory rat chow *ad libitum*. At the commencement of the experiment, all MEDAN administration was done orally.

### 2.3 Acute Toxicity Test

Twelve (12) mice were used to assess acute toxicity profile following Lorke's method [32]. The test was conducted in two phases. In the first phase, nine (9) mice were divided into three (3) groups of three (3) mice each. Groups 1, 2, and 3 were administered 10, 100, and 1000 mg/kg of MEDAN, respectively, to determine the dose range that might cause toxic effects. In the second phase, the remaining three (3) mice received 1600, 2900, and 5000 mg/kg of MEDAN. They were monitored for 24 hours for signs of weakness or mortality. The LD<sub>50</sub> was calculated to be 5000 mg/kg of MEDAN.

### 2.4 Preemptive and Therapeutic Analgesia Tests

For the preemptive analgesia test, Wistar rats were divided into six groups, with five animals in each group. Group I, the negative control, received distilled water, while Group II, the positive control, was given the standard pain

medication Flexicam (1.5 mg/kg) (Hovid, Malaysia). Groups III and IV were administered 500 mg/kg and 1000 mg/kg of MEDAN, respectively. Groups V and VI received 500 mg/kg and 1000 mg/kg of MEDAN in combination with the standard drug. Both MEDAN and the standard drug were administered orally for two weeks before the experiment. The tests were carried out on day 15, 17 and 19<sup>th</sup>. The therapeutic analgesia test followed a similar design to the preemptive test involving sixty Wistar and grouped six (6) groups of five (5) animals treated with MEDAN and standard drug in the same manner as the preemptive analgesia test, however, the pain was first induced using an analgesimeter before treatment with extracts and standard drugs.

## 2.5 Analgesimeter Test

This experiment followed the previously described method [33,34]. The rat was suspended vertically, and its left hind paw was placed between the plinth and the finger of the Ugo Basile analgesimeter (Ugo Basile 7200, Italy). The machine measured pain thresholds by applying increasing pressure to the hind paw. The force at which each rat withdrew its paw was recorded. This was considered the pain threshold and expressed as force per gram (f/g). The recording of pain threshold was done the 15, 17 and 19<sup>th</sup> days for the preemptive analgesia and 1, 2 and 4 hours post-treatment for the therapeutic analgesia test.

## 2.6 Sample Collection and Analysis

After the study, blood samples were obtained from the recto-orbital sinus and transferred to a plain sample bottle. The blood was allowed to sit for 30 minutes before being centrifuged at 3000 rpm for 5 minutes. The supernatant serum was carefully decanted using a micropipette and stored at 2°C until laboratory analysis. The concentration of C-reactive protein in the serum was measured using standard laboratory test kits (Elabscience, China) via the ELISA method.

## 2.7 Insilco Molecular Docking

This method combines computer science and computational methods with experimental biology by simulating and predicting molecular interactions, gene regulation and drug targets [35,36]. The crystal structure of C-reactive protein (CRP) was retrieved from the Protein Data Bank and prepared using AutoDock v4.2

(Scripps Research Institute, USA) by removing ligands and water, adding missing hydrogens, and saving the structure in PDBQT format. Ligands were docked to various protein targets, and binding affinities were evaluated using AutoDock Vina. The docking grid for CRP was established, and cluster analysis based on RMSD values identified the most reliable binding conformations. Compounds with binding affinities greater than the standard antimalarial (artesunate) were visualized with Discovery Studio Visualizer (BIOVIA, USA).

## 2.8 Statistical Analysis

Data analysis was performed using GraphPad Prism version 8.0.1 (244). One-way analysis of variance (ANOVA) and post hoc tests were conducted to assess mean differences. The results are presented as means and standard errors of the mean ( $M \pm SEM$ ), with statistically significant values indicated at  $p < 0.05$  within a 95% confidence interval.

## 3. RESULTS

Table 1 presents the pain threshold in male Wistar rats following three weeks of administration of methanolic leaf extract of *Datura alba ness* (MEDAN). The data indicate that MEDAN at doses of 500 and 1000 mg/kg significantly increased the pain threshold on day 15 in a dose-dependent manner compared to both negative and positive controls ( $p < 0.05$ ). However, when combined with Flexicam, the 500 mg/kg MEDAN dose resulted in a significantly reduced pain threshold, while the 1000 mg/kg MEDAN dose improved the pain threshold. On day 17, there was a significant decrease in pain threshold for rats treated with 500 and 1000 mg/kg MEDAN ( $p < 0.05$ ), which was reversed when MEDAN was combined with a standard drug, leading to significantly higher pain thresholds for both dosages. By day 19, the increased pain threshold persisted in rats treated with 1000 mg/kg MEDAN, and further improvement was observed when the 500 mg/kg dose was combined with the standard drug. These findings suggest that *Datura alba ness* exhibits potential analgesic effects, which may be enhanced when used in conjunction with standard pain relief medications.

Table 2 presents the changes in pain threshold in male Wistar rats following oral administration of leaf extracts of *Datura alba Ness* (MEDAN). The results show that doses of 500 and 1000 mg/kg of MEDAN, as well as 500 mg/kg MEDAN in

**Table 1. Preventive effects of methanolic leave extract of *Datura alba ness* on pain threshold in male *Wistar* rats using an analgesimeter after 3 weeks of administration**

Research Groups	Pain Threshold (f/g)		
	Day15	Day17	Day 19
Negative control (distilled water)	142.50±2.79	208.50±6.34	165.00±2.34
Positive control (Felxicom 1.5 mg/kg)	148.00±1.89	222.10±2.92	239.00±2.58
500mg/kg MEDAN	172.30±4.24 <sup>ab</sup>	155.10±4.55 <sup>ab</sup>	166.00±2.82 <sup>b</sup>
1000mg/kg MEDAN	227.50±4.85 <sup>ab</sup>	190.70±3.50 <sup>b</sup>	222.80±3.65 <sup>ab</sup>
500mg/kg MEDAN +Felxicom	114.30±1.71 <sup>ab</sup>	228.10±3.43 <sup>a</sup>	261.80±3.83 <sup>ab</sup>
1000mg/kg MEDAN +Felxicom	165.60±5.31 <sup>ab</sup>	169.20±2.62 <sup>ab</sup>	184.00±4.57 <sup>ab</sup>

Data are expressed as Mean ± SEM; p < 0.05<sup>a</sup>: significant compared to negative control (Group 1); p < 0.05<sup>b</sup>: significant compared to Standard drug (Felxicom) (Group 2)

**Table 2. Therapeutic effects of methanolic leaves extract of *Datura alba ness* on pain threshold in male *Wistar* rats**

Research Groups	Pain Threshold (f/g)		
	After 1 hr	After 2 hrs	After 4 hrs
Negative control (distilled water)	145.70±2.72	170.40±3.32	212.90±3.45
Positive control (Felxicom 1.5 mg/kg)	87.33±2.85	243.90±3.68	257.10±4.35
500mg/kg MEDAN	206.50±4.80 <sup>ab</sup>	162.90±4.84 <sup>b</sup>	230.10±3.87 <sup>ab</sup>
1000mg/kg MEDAN	163.30±4.13 <sup>ab</sup>	172.10±2.50 <sup>b</sup>	203.50±2.37 <sup>b</sup>
500mg/kg MEDAN +Felxicom	238.80±5.01 <sup>ab</sup>	169.90±3.56 <sup>b</sup>	208.00±3.88 <sup>b</sup>
1000mg/kg MEDAN +Felxicom	124.30±2.01 <sup>ab</sup>	160.00±3.13 <sup>b</sup>	250.30±4.06 <sup>a</sup>

Data are expressed as Mean ± SEM; p < 0.05<sup>a</sup>: significant compared to negative control (Group 1); p < 0.05<sup>b</sup>: significant compared to Standard drug (Felxicom) (Group 2)

combination with a standard drug, resulted in an increased pain threshold one hour post-administration. However, after two hours, this effect diminished, as all MEDAN doses and their combinations with the standard drug led to significantly lower pain thresholds compared to the standard drug (p < 0.05). After four hours, 500 mg/kg MEDAN caused an increase in pain threshold compared to both the negative control and the standard drug, while 1000 mg/kg MEDAN significantly decreased the pain threshold in comparison to the standard drug. When combined with the standard drug, both 500 and 1000 mg/kg MEDAN resulted in a significant reduction in pain threshold compared to the standard drug and the negative control, respectively.

Table 3 illustrates the effects of a three-week oral administration of methanolic leaf extract of *Datura alba ness* (MEDAN) on serum C-reactive protein levels. The data reveal that all MEDAN doses and their combinations with a standard drug significantly reduced the mean levels of C-reactive protein compared to the standard drug (p < 0.05). However, no changes were observed when compared to the negative control.

Table 4 shows the binding Affinity of compounds to CRP using celecoxib as the reference compound, the binding of the C=reactive protein to the reference drug is -9.0 out of the 29 compounds from the extract [25]. The following compound showed a binding affinity with the


**Table 3. Effects of methanolic leaf extract of *Datura alba ness* on serum level C Reactive Protein of male *Wistar* rats after three weeks of administration.**

Research Groups	C-Reactive Protein(mg/l)
Negative control (distilled water)	266 ± 3.81
Positive control (Felxicom 1.5 mg/kg)	380± 6.52
500mg/kg MEDAN	298 ± 7.08 <sup>b</sup>
1000mg/kg MEDAN	300± 17.8 <sup>b</sup>
500mg/kg MEDAN +Felxicom	278 ± 3.53 <sup>b</sup>
1000mg/kg MEDAN +Felxicom	279 ± 2.20 <sup>b</sup>


Data are expressed as Mean ± SEM; p < 0.05<sup>a</sup>: significant compared to negative control (Group 1); p < 0.05<sup>b</sup>: significant compared to Standard drug (Felxicom) (Group 2)

**Table 4. *In silico* molecular docking chart showing binding affinity of compounds to CRP using celecoxib as the reference compound**

S/N	Compound Name	GCMC Area%	GCMC Height%	GCMC H/A mark	Bat Ref Drug C-pr	Bat Cpd C-pr	Lipinski Violation
R	Celecoxib				-9.0		
1	Dodecanoic acid, 1,2,3-propanetriyl ester	4.32	2.21	5.07			2
2	Stigmasterol	3,11	2.74	3.01		-9.2	1
3	Gamma sitosterol	10.77	7.98	3.57		-9.2	1
4	Cholest-5-en-3-ol, 24-propylidene-, (3β)	2.18	1.40	4.12		-9.7	1



**Fig. 1. 2D and 3D view of the interaction between celecoxib and amino acid in the binding site of c-reactive protein**



**Fig. 2. 2D and 3D view of the interaction between stigmasterol and amino acid in the binding site of c-reactive protein respectively**

CPR receptor. Dodecanoic acid, 1,2,3-propanediol ester, Stigmasterol, gamma sitosterol and Cholest-5-en-3-ol, 24-propylidene-, (3β).

#### 4. DISCUSSION

Traditional medicine in Africa and Asia relies on the use of leaves, roots, seeds, and stems from various plants to treat different ailments. Numerous studies have sought to validate the effectiveness of some of these plants for pain

management, leading to the identification of bioactive compounds that could be developed into standardized pharmaceuticals. Additionally, the occurrence of side effects linked to synthetic drugs has sparked a considerable interest in natural products, resulting in an increase in research focused on their potential benefits [37-39]. While our previous research focused on the proximate composition, phytochemicals, and bioactive components of *Datura alba nesa* (DAN), the current study aims to evaluate the potential analgesic activity of DAN extracts using

preemptive and therapeutic analgesia models in Wistar rats, along with investigating the possible mechanism of action through molecular docking.

Our findings show that the methanolic leaf extract of *Datura alba nesa* (MEDAN) significantly increased the pain threshold in male Wistar rats after three weeks of administration in a dose-dependent manner, with the 1000 mg/kg dose showing sustained effects. Combining MEDAN with Flexicam enhanced pain relief at higher doses (Table 1). However, for the therapeutic analgesic test, we observed that MEDAN increases pain thresholds within one hour but diminishes over time, with effects varying by dose and combination with standard drugs (Table 2). These observed analgesic effects of MEDAN could be attributed to a variety of bioactive compounds present in MEDAN. Our earlier finding showed the presence of some fatty acids (n-Hexadecanoic acid, Octadecanoic acid and Ricinoleic acid) plant sterols (Stigmasterol and Gamma-Sitosterol), Vitamins (Delta-Tocopherol (Vitamin E and [40]) and Triterpene (Squalene) [25] which have been shown to possess potent modulators of pain and inflammatory response [41-45]. These compounds have been reported to suppress pain and inflammation by inhibiting inflammatory cytokines which can contribute to pain relief or by inhibiting prostaglandin synthesis, which is involved in pain signaling. They have also been shown to be potent antioxidants which help in pain management by reducing oxidative stress and inflammation [41-45]. These findings indicate that *Datura alba nesa* may possess analgesic properties, which could be amplified when combined with standard pain relief medications.

The present study also reveals that *Datura alba nesa* significantly lowered serum C-reactive protein levels after three weeks of oral administration, both at all doses and in combination with a standard drug, compared to the standard drug, but showed no difference from the negative control. C-reactive protein (CRP) is an acute-phase protein synthesized by the liver in response to inflammation and stimulated by inflammatory cytokines, particularly interleukin-6 (IL-6) [46,47]. They are often elevated in conditions associated with pain, such as arthritis, fibromyalgia, and other inflammatory diseases. The presence of CRP reflects ongoing inflammation, which is a key driver of pain [48,49]. Measuring CRP levels can help assess the severity of inflammation and, indirectly, the intensity of pain. Higher CRP levels often

correlate with increased pain levels in various conditions [50]. Also, the molecular docking results from this study indicated that certain compounds in the extract, including stigmasterol, gamma-sitosterol, and cholest-5-en-3-ol, 24-propylidene-(3 $\beta$ ), showed binding affinity to C-reactive protein receptors, mimicking the effects of the reference drug, celecoxib. Celecoxib is prescribed for symptomatic relief of osteoarthritis and rheumatoid arthritis in adults, as well as for acute pain from various causes. This medication acts as a selective noncompetitive inhibitor of the cyclooxygenase-2 (COX-2) enzyme, which is predominantly expressed in inflamed tissues and is induced by inflammatory mediators [51]. Inhibiting COX-2 reduces the production of metabolites such as prostaglandin E2 (PGE2), prostacyclin (PGI2), thromboxane (TXA2), prostaglandin D2 (PGD2), and prostaglandin F2 (PGF2) [52]. The reduced level of serum CRP observed suggests a potent ability of MEDAN to reduce pain sensitivity and this may be my potent than standard pain relief medication as observed in the study. These findings are in tandem with analgesic properties observed in other *Datura* species, *Datura fastuosa* and *Datura metel* [27,28].

## 5. CONCLUSION

The use of natural products derived from medicinal plants as therapeutic remedies for pain management is gaining increasing attention due to their broad acceptability, availability, affordability, and minimal or nonexistent side effects. According to the data from the current study, methanol extracts from *Datura alba nesa* leaves demonstrate significant efficacy in reducing pain by lowering C-reactive protein levels. This effect was particularly pronounced when the 500 mg/kg dose of MEDAN was combined with the standard drug, Flexicam. This study offers a scientific foundation that supports the traditional medicinal use of *Datura alba nesa* in treating pain and inflammation.

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative Artificial intelligence (AI) technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

## CONSENT

It is not applicable.

## ETHICAL APPROVAL

The research design and protocol adhered to all guidelines regarding the use of laboratory animals. Ethical approval was obtained from the University of Port Harcourt Research Ethics Committee [REC/MM89/221].

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Steeds CE. The anatomy and physiology of pain. Surgery (Oxford). 2009;27(12):507-11.
2. Schaible H-G, Richter F. Pathophysiology of pain. Langenbeck's Archives of Surgery. 2004;389:237-43.
3. Møller AR. Anatomy and physiology of pain. Textbook of Tinnitus. 2011;121-32.
4. Hakim AJ, Keer RJ, Grahame R. Hypermobility, fibromyalgia and chronic pain: Elsevier Health Sciences; 2010.
5. Helms JE, Barone CP. Physiology and treatment of pain. Crit. Care Nurs. 2008;28(6):38-49.
6. Berde CB, Sethna NF. Analgesics for the treatment of pain in children. NEJM. 2002;347(14):1094-103.
7. Farquhar-Smith WP. Anatomy, physiology and pharmacology of pain. Anaesthesia & Intensive Care Medicine. 2008;9(1):3-7.
8. Stark TD, Mtui DJ, Balemba OB. Ethnopharmacological survey of plants used in the traditional treatment of gastrointestinal pain, inflammation and diarrhea in Africa: Future perspectives for integration into modern medicine. Animals. 2013;3(1):158-227.
9. Khumalo GP, Van Wyk BE, Feng Y, Cock IE. A review of the traditional use of southern African medicinal plants for the treatment of inflammation and inflammatory pain. J. Ethnopharmacol. 2022;283:114436.
10. Nasim N, Sandeep IS, Mohanty S. Plant-derived natural products for drug discovery: current approaches and prospects. The Nucleus. 2022;65(3):399-411.
11. Patwardhan B, Vaidya AD, Chorghade M, Joshi SP. Reverse pharmacology and systems approaches for drug discovery and development. Curr. Bioact. Compd. 2008;4(4):201-12.
12. Bukhari IA, Khan RA, Gilani AH, Ahmed S, Saeed SA. Analgesic, anti-inflammatory and anti-platelet activities of the methanolic extract of *Acacia modesta* leaves. Inflammopharmacology. 2010;18:187-96.
13. Olajide OA, Awe SO, Makinde JM, Ekhelar AI, Olusola A, Morebise O, Okpako DT. Studies on the anti-inflammatory, antipyretic and analgesic properties of *Alstonia boonei* stem bark. J. Ethnopharmacol. 2000;71(1-2):179-86.
14. Ishola IO, Awodele O, Olusayero AM, Ochieng CO. Mechanisms of analgesic and anti-inflammatory properties of *Annona muricata* Linn. (Annonaceae) fruit extract in rodents. J. Med. Food. 2014;17(12):1375-82.
15. Igwe SA, Akunyili DN. Analgesic Effects of Aqueous Extracts of the leaves of *bryophyllum pinnatum*. Pharm. Biol. 2005;43(8):658-61.
16. Owoyele BV, Oguntoye SO, Dare K, Ogunbiyi BA, Aruboula EA, Soladoye AO. Analgesic, anti-inflammatory and antipyretic activities from flavonoid fractions of *Chromolaena odorata*. J. Med. Plant Res. 2008;2(9):219-25.
17. Sofidiya MO, Odukoya OA, Adedapo AA, Mbagwu HOC, Afolayan AJ, Familoni OB. Investigation of the anti-inflammatory and antinociceptive activities of *Hymenocardia acida* Tul.(Hymenocardiaceae). AJB. 2010;9(49):8454-9.
18. Ijeoma UF, Aderonke SO, Ogbonna O, Augustina MA, Ifeyinwa C-N. Antinociceptive and anti-inflammatory activities of crude extracts of *Ipomoea involucrata* leaves in mice and rats. Asian Pac. J. Trop. Med. 2011;4(2):121-4.
19. Olorunfemi OJ, Nworah DC, Chinko BC, Joffa PPK, Pughikumo DT. The analgesic effect of ethanolic extract of *Myristica fragrans* Houtt (nutmeg) on mice. Am J PharmTech. Res. 2012;2:265-70.
20. Komakech R, Kim Y-g, Matsabisa GM, Kang Y. Anti-inflammatory and analgesic potential of *Tamarindus indica* Linn.(Fabaceae): A narrative review. Integr. Med. Res. 2019;8(3):181-6.
21. Okokon JE, Davies K, Antia BS. Analgesic and anti-inflammatory activities of *Zea mays* leaves. J. Med Herb. 2016;7(2):73-82.
22. Ibrahim M, Siddique S, Rehman K, Husnain M, Hussain A, Akash MSH, Azam F. Comprehensive analysis of phytochemical constituents and



- ethnopharmacological investigation of genus *Datura*. *Critical Reviews™ in Eukaryotic Gene Expression*. 2018;28(3).
23. Sharma M, Dhaliwal I, Rana K, Delta AK, Kaushik P. Phytochemistry, pharmacology, and toxicology of *Datura* species—A review. *Antioxidants*. 2021;10(8):1291.
  24. Priya KS, Gnanamani A, Radhakrishnan N, Babu M. Healing potential of *Datura alba* on burn wounds in albino rats. *J. Ethnopharmacol*. 2002;83(3):193-9.
  25. Ovili-Odili BZ, Adienbo OM, Daubry TME, Chinko BC. Phytochemical profiling and asian plant res. *J. Nutraceutical benefits of methanol leaf extracts from *Datura alba* ness*. 2024;12(6): 1-11.
  26. Maheshwari NO, Khan A, Chopade BA. Rediscovering the medicinal properties of *Datura* sp.: A review. *J. Med. Plant Res*. 2013;7(39):2885-97.
  27. Abena AA, Miguel LM, Mouanga A, Assah TH, Diatwa M. Evaluation of analgesic effect of *Datura fastuosa* leaves and seed extracts. *Fitoterapia*. 2003;74(5):486-8.
  28. Adeola BS. *Datura Metel* L: analgesic or hallucinogen?"Sharo" Perspective. *Middle East J. Sci. Res*. 2014;21:993-7.
  29. Nerlekar N, Patil P, Khot S, Kulkarni A, Dandge P, Berde A. Cold maceration extraction of wild fruit *Terminalia bellirica* (Gaertn.) Roxb.: exploring its bioactives for biomedical applications. *Prep Biochem Biotech*. 2024;1-19.
  30. Lezoul NEH, Belkadi M, Habibi F, Guillén F. Extraction processes with several solvents on total bioactive compounds in different organs of three medicinal plants. *Molecules*. 2020; 25(20):4672.
  31. Albus U. Guide for the care and use of laboratory animals (8th edn). SAGE Publications Sage UK: London, England; 2012.
  32. Lorke D. A new approach to practical acute toxicity testing. *Arch. Toxicol*. 1983;54(4):275-87.
  33. Mbiantcha M, Kamanyi A, Teponno RB, Tapondjou AL, Watcho P, Nguelefack TB. Analgesic And Anti-Inflammatory Properties Of Extracts From The Bulbils Of *Dioscorea Bulbifera* L. *Var Sativa* (Dioscoreaceae) In Mice And Rats. *Evid. - Based Complementary Altern. Med*. 2010;2011:1-9.
  34. Nguelefack TB, Fotio AL, Watcho P, Wansi SL, Dimo T, Kamanyi A. Analgesic properties of the aqueous and ethanol extracts of the leaves of *Kalanchoe crenata* (Crassulaceae). *Phytother Res*. 2004;18(5):385-8.
  35. Wang W, Ye Z, Gao H, Ouyang D. Computational pharmaceutics-A new paradigm of drug delivery. *J Control Release*. 2021;338:119-36.
  36. Jakhar R, Dangi M, Khichi A, Chhillar AK. Relevance of molecular docking studies in drug designing. *Current Bioinformatics*. 2020;15(4):270-8.
  37. Ugoeze KC, Aja PC, Nwachukwu N, Chinko BC, Egwurugwu JN, Oluigbo KE. Evaluation of the wound healing potentials of aqueous topical creams containing aqueous extract of *Azadirachta indica* leaves as bioactive ingredient. *J Pharm Pharmacol Res*. 2021;5(1):176-87.
  38. Hart VO, Chinko BC. Effect of *Terminalia catappa* Seed Extract on Liver Enzymes and Lipid Profile in Male Wistar Rat. *Asian Journal of Research in Botany*. 2022;8(4):25-32.
  39. Chinko BC, Orlu CN. Antipyretic and Anti-Inflammatory Effects of *Ocimum gratissimum* in Male Wistar Rats. *South Asian Research Journal of Natural Products*. 2023;6(1):28-38.
  40. Jiang Q, Moreland M, Ames BN, Yin X. A combination of aspirin and  $\gamma$ -tocopherol is superior to that of aspirin and  $\alpha$ -tocopherol in anti-inflammatory action and attenuation of aspirin-induced adverse effects. *J Nutr Biochem*. 2009;20(11):894-900.
  41. Mohammad H, Prabhu, K, Rao, M R K, Sundram, L., Dinakar, S., Sathish Kumar, M & Vijayalakshmi N. The gas chromatography-mass spectrometry study of one ayurvedic pain relieving oil "Mahamasha Thailam". *Drug Invention Today*, 2019; 12(7), 1524-1527.
  42. Zilani MNH, Islam MA, Biswas P, Anisuzzman M, Hossain H, Shilpi JA. Metabolite profiling, anti-inflammatory, analgesic potentials of edible herb *Colocasia gigantea* and molecular docking study against COX-II enzyme. *J. Ethnopharmacol*. 2021;281:114577.
  43. Vieira C, Evangelista S, Cirillo R, Lippi A, Maggi CA, Manzini S. Effect of ricinoleic acid in acute and subchronic experimental models of inflammation. *Mediators Inflamm*. 2000;9(5):223-8.
  44. Walker CIB, Oliveira SM, Tonello R, Rossato MF, da Silva Brum E, Ferreira J, Trevisan G. Anti-nociceptive effect of stigmasterol in mouse models of acute and

- chronic pain. *Naunyn Schmiedebergs Arch Pharmacol.* 2017;390:1163-72.
45. Morgan LV, Petry F, Scatolin M, de Oliveira PV, Alves BO, Zilli GAL. Investigation of the anti-inflammatory effects of stigmaterol in mice: insight into its mechanism of action. *Behav. Pharmacol.* 2021;32(8):640-51.
  46. Eklund CM. Proinflammatory cytokines in CRP baseline regulation. *Adv. Clin. Chem.* 2009;48: 111-36.
  47. Sproston NR, Ashworth JJ. Role of C-reactive protein at sites of inflammation and infection. *Front Immunol.* 2018;9:754.
  48. Thudium CS, Löfvall H, Karsdal MA, Bay-Jensen A-C, Bihlet AR. Protein biomarkers associated with pain mechanisms in osteoarthritis. *J. Proteom.* 2019;190:55-66.
  49. Graham-Engeland J, DeMeo NN, Jones DR, Mathur A, Smyth JM, Sliwinski MJ, et al. Individuals with both higher recent negative affect and physical pain have higher levels of C-reactive protein. *BBI-Health.* 2022;21:100431.
  50. Ansar W, Ghosh S, Ansar W, Ghosh S. Inflammation and inflammatory diseases, markers, and mediators: Role of CRP in some inflammatory diseases In *Biology of C Reactive Protein in Health and Disease.* Springer, New Delhi. 2016:67-107.
  51. Isakson PC. Pharmacology of Cox-2 inhibitors. *Prog. Exp. Tumor Res.* 2003;37:25-51.
  52. Iezzi A, Ferri C, Mezzetti A, Cipollone F. COX-2: friend or foe? *Current Pharmaceutical Design.* 2007;13(16):1715-21.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/125537>