



A Case Report: Post-Operative Physiotherapy Rehabilitation Program for a Patient with Tibial Osteosarcoma

Sapna Dhote ^a, Dushyant Bawiskar ^{a#} and Pratik Phansopkar ^{a##}

^a Department of Musculoskeletal Physiotherapy, Ravi Nair Physiotherapy College, Datta Meghe Institute of Medical Sciences, Sawangi Meghe, Wardha, Maharashtra, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i58A34143

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/71126>

Case Report

Received 10 October 2021

Accepted 14 December 2021

Published 15 December 2021

ABSTRACT

Tumour or bone tumour can occur in any bone and called it as a Osteosarcoma. It is more prone in the long bones of the body especially at metaphyseal growth plate of the bone. Femur, Tibia and Humours are the most common bone for the osteosarcoma and in some research pelvis bone and skull or jaw bone also involved. There are many factors which cause the sarcoma might be due lack of physical activity. The Chondroblastoma as Giant Cell Tumour of the Humours of Epiphysial Chondromatous in nature. Proximal humerus and Knee are the most common and most affected bone by the Chondroblastomas at the level of epiphyses of the longer bone. 62% cases show the thinning of the Subchondral structure and 33% cases shows the collapse and chondral breach Mostly (90%) it is found in the age group of 5 to 25 years of age with the male and female ratio is of 2:1. Psychological and physical improvement is seen in the patient of sarcoma and enhances participation in the exercise with great enthusiasm and help in the reduction the impact of symptoms of diseases. Physical therapy exercise protocol starts from low to moderate intensity which involved resistance exercise, aerobic exercise and combination of both exercises.

Conclusion: Physiotherapy has been demonstrated to be useful in post-operative osteosarcoma

⁼ Intern;

[#] Associate Professor & HOD;

^{*}Corresponding author: E-mail: drpratik77@gmail.com;

patients in our case study. Physiotherapy assists the patient in de-stressing from both mental and physical stress. The goal of improving function through health-care management and goal-setting was met on schedule. Taking excellent care of yourself after surgery can help you avoid issues.

Keywords: Osteosarcoma; oncology rehabilitation; blastosarcoma; chondroblastoma; giant cell tumour rehabilitation.

1. INTRODUCTION

In 1928, Codman said about the chondroblastoma as giant cell tumour of the Humours of Epiphysial Chondromatous in nature. Proximal humerus and Knee are the most common and most affected bone by the Chondroblastomas at the level of epiphyses of the longer bone. 62% cases show the thinning of the Subchondral structure and 33% cases shows the collapse and chondral breach. Mostly (90%) it is found in the age group of 5 to 25 years of age with the male and female ratio is of 2:1. Pain during activity or at rest, tenderness at local place with effusion and decrease in the ROM of the affected limb are the common symptoms which present in it [1]. The exact aetiology to include the genetics and environmental factors and genetics are the two factors for the presenting the neoplasm in the patient and few abnormalities at different chromosomes are mentioned in few researchers that predispose patients to the formation of these neoplasms is not well understood [2]. Surgical Excision is the best procedure for the neoplasm because during the procedure complete neoplasm is removed and prevent relapse and dysfunction of joint [3]. An intensive procedure or curettage is suggested for lesions, which could compromise an open forum for the epiphysis and result in the growth and development of the limb being shortened and deformed [4]. Before the operative procedure, conventional radiology is used as a first line of investigation in the patient with tumours [5]. Telangiectatic, osteoblastic, fibroblastic, chondroblastic, parosteal, multifocal and periosteal types are the different conventional types for the osteosarcoma [6]. The treatment of osteosarcoma requires a multidisciplinary approach involving the family physician, orthopaedic oncologist, medical oncologist, radiologist and pathologist. In children and adolescents, it is 3rd most common

condition. Localized diseases are presents with patients. Sclerotic and lytic lesion are seen in the radiographic investigation in Metaphyseal region of the bone. Amputation used to carried out before 1970 for the treatment of osteosarcoma which caused more death rate (80%) [7].

2. PATIENT INFORMATION

Patient came to OPD with Complaint of pain and instability over right knee for 12 months. Patient gives alleged history of fall while riding bicycle 12 months ago. Pain was sudden in onset, gradually progressive in nature and dull aching type which gets aggravated by walking, movement and gets relieved by taking rest and medication. Pain was associated with swelling for 2 months. The swelling was firm in consistency, hard to touch with local temperature raised. Patient previously went to private hospital where he was managed conservatively and after which he was referred to this hospital. On local examination patient examined in lying down position with both ASIS at same level. On Inspection Right Knee overlying skin normal diffuse swelling present over knee. No limb length discrepancy seen. On Palpation Right Knee Local temperature raised over medial condyle of tibia. Tenderness presents over medial condyle of tibia, Quadriceps and Calf. On USG guided biopsy was done and sample was sent for Histopathological examination which was suggestive of Chondroma of Medial Condyle Proximal Tibia Right Side. Then patient was managed with Tumour Excision and reconstruction with Bone Grafting for Chondroma of Medial Condyle proximal tibia right side under Spinal Anaesthesia. Bone graft was harvested from ASIS by giving 5cm incision and bone wax was applied. After Operative Procedure, patient shifted to Physiotherapy Department for further management. On Assessment at Physiotherapy OPD.

Initial Examination Findings:

Table 1. Pre-range of motion

Pre-Physiotherapy Assessment Day 1:				
Joint	Right		Left	
	Active	Passive	Active	Passive
Hip Flexion	0-70(assisted)	0-72	0-112	0-115
Hip Extension	0-18(assisted)	0-22	0-30	0-32
Hip Abduction	0-16(assisted)	0-20	0-40	0-45
Hip Adduction	0-12(assisted)	0-14	0-27	0-29
Knee Flexion	0-90(assisted)	0-91	0-140	0-144
Knee Extension	80-0(assisted)	82-0	140-0	144-0
Ankle Planter Flexion	0-45	0-46	0-50	0-50
Dorsiflexion	0-8	0-10	0-10	0-10

Table 2. Pre-manual muscle testing

Pre-Physiotherapy Assessment Day 1:			
		Right	Left
Hip	Abductor	3/5	5/5
	Adductor	3/5	5/5
	Flexors	3/5	5/5
	Extensors	3/5	5/5
	Internal Rotators	3/5	5/5
	External Rotators	3/5	5/5
Knee	Extensors	3/5	5/5
	Flexors	3/5	5/5
Ankle	Plantar Flexors	4/5	5/5
	Dorsi Flexors	4/5	5/5

*Pain Assessment Day 1: - On NPRS at rest 6/10 and on slight movement 9/10.
Berg balance scale: -4/56
Dynamic Gait index: -0/24*

INTERVENTION

Week 1:- Maximum Protection

Ice Pack is used for the reduction of the inflammation at the site of incision and it was given for 10-15min. Exercises which are active assisted in nature are started with the right hip, right knee and Right Ankle Toe movements given to increase the range of motion and 5 repetition twice a day. Resistance exercise must be given to the left lower limb and both upper limbs with the weights. Statics or Isometric Contraction exercises are main to increase the strength of the muscle so it must be given to the Quadriceps, Hamstrings and Gluteus with the hold of 5 sec with 5 repetitions twice a day. Along with these exercise, the patient must be counsel for the Important of Exercise and advantages of exercise.

Week 2 to 6-Progression in strengthening and ROM

For the control of inflammation use of ice pack is continued. For the prevention of the knee deformity, flexion and extension was carried out and which helped from formation of contracture. Uninvolved limbs was continued with resistance exercise and intensity should be increase and repetitions also be increase. Dynamic quadriceps exercise help to increase the strength of the Right limb in pain free range of motion. With the help of walker, partial weight bearing exercise was continued and improvement in functional task must be observed. Every exercise must be done twice a day.

Week 6 to 8:- Strengthening and proprioceptive phase

Passive Manual training was given to increase the range of motion of knee joint. Heel Slides

with help of towel pulling and or with help of opposite lower limb by assisting. In functional Training, Surge Faradic Current is used for the stimulation to the Quadriceps and Hamstrings Muscles. Training for balance and training for equilibrium was given with 3 repetitions. Now the gait training must be start with the help of walker with touching the lower limb by taking the weight over the planter surface with 50-75 percent and progression must be after every 1 week. Parallel bars are used for the gait training.

Week 8 to 10:- Advanced strengthening and balance regime

For the improvement of the Proprioception, all the above exercise must be carried out in

proper manner with progression level. Right lower limb was continued with strengthen exercise so it will not get fatigue early. Progression of the weight on affected limb must be increased in proper training program. 100% weight bearing can be done after contacted with Orthopaedic surgeon and Radiographic Examination must be carried out for the complete recovery of the patient. After Gait training, treadmill should be start in slow manner. 30 minutes session per day for 3-4 days per week must be done. Heart rate and other vitals must be observed. Endurance protocol should be start at the time of Treadmill Exercise.

FOLLOW UP

After Exercise Examination Findings

Table 3. Post-range of motion

Post-Physiotherapy Assessment				
Joint	Right		Left	
	Active	Passive	Active	Passive
Hip Flexion	0-90(assisted)	0-94	0-112	0-115
Hip Extension	0-22(assisted)	0-24	0-30	0-32
Hip Abduction	0-18(assisted)	0-20	0-40	0-45
Hip Adduction	0-14(assisted)	0-16	0-27	0-29
Knee Flexion	0-110(assisted)	0-112	0-140	0-144
Knee Extension	110-0(assisted)	112-0	140-0	144-0
Ankle Planter Flexion	0-46	0-47	0-50	0-50
Dorsiflexion	0-10	0-11	0-10	0-10

Table 4. Post-manual muscle testing

Post-Physiotherapy Assessment			
		Right	Left
Hip	Abductor	4/5	5/5
	Adductor	4/5	5/5
	Flexors	4/5	5/5
	Extensors	4/5	5/5
	Internal Rotators	4/5	5/5
	External Rotators	4/5	5/5
Knee	Extensors	4/5	5/5
	Flexors	4/5	5/5
Ankle	Plantar Flexors	4/5	5/5
	Dorsi Flexors	4/5	5/5

PAIN ASSESSMENT

Pre-rehabilitation: NPRS 6/10 at rest, 9/10 on slight movement.

Post-rehabilitation: NPRS 2/10 at rest, 4/10 on activity.

SWELLING: MID PATELLAR GIRTH

Pre-rehabilitation: Left=27cm and Right=33cm
Post-rehabilitation: Left=27cm and Right=33cm.

BERG BALANCE SCORE

Pre-rehabilitation:4/56
Post-rehabilitation:48/56

DYNAMIC GAIT INDEX

Pre-rehabilitation:0/24
Post-rehabilitation:19/24

3. DISCUSSION

Profession which belong to health care and deal with function and movement is a Physical Therapy Profession. Physiotherapist diagnose and treat all individuals from Neonates to Old Age persons as the have medical issue or any medical problem or any other condition which limit or decrease the ability of the work or movement get hampered and cause difficulty in carry out daily activity. Musculoskeletal, Cardiopulmonary, Neuromuscular, and Integumentary are the various departments who have big role in the treatment of Oncology Setting [8]. Hawley C. Almstedt and his team explained in his project combined aerobic and resistance training improves bone health of female cancer survivors and concluded that osteoporosis risk can be reduce by doing exercise and maintaining the bone strength [9]. Exercise improves quality of life in patients with cancer is the study done by Jasper K W Gerritsen, Arnaud J P E Vincent and explained about different exercise which directly and indirectly help to improve the quality of life of the patient [10]. Compliance and satisfaction with intensive physiotherapy treatment during chemotherapy in patients with bone tumours and evaluation of related prognostic factors: An observational study done by MattiaMorri suggested it is a feasible treatment for bone tumour patients who have shown they can positively adhere to it [11,12]. Wiebke Jensen and team said in their research that physical exercise and therapy in terminally ill cancer patients will get help from physical exercises, relaxation therapy, and breathing training and these were the most commonly used modalities and it is more feasibility and acceptability [13,14]. Radiotherapy may cause malignant transformation so it is contraindicated in the management of the chondroblastoma [15,16].

Cryotherapy and analgesic help to reduce the pain and swelling in the osteochondroma patient [17,-20]. Dr. Paramanadam and his team concluded study that Physical rehabilitation is the most important and key for the good outcomes for the functional group and if the early rehabilitation is not done then there will be decrease in the quality of life of the patient [21-24].



Fig. 1. Pre-operative Xray



Fig. 2. Post-operative Xray



Fig. 3. Post-operative attitude of limb with scar mark



Fig. 4. Minimal weight bearing over the affected limb at the end of third week



Fig. 5. Bone graft taken from ASIS

4. CONCLUSION

Physiotherapy has been demonstrated to be useful in post-operative osteosarcoma patients in our case study. Physiotherapy assists the patient in de-stressing from both mental and physical stress. The goal of improving function through health-care management and goal-setting was met on schedule. Taking excellent care of yourself after surgery can help you avoid issues.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and

producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

CONSENT AND ETHICAL APPROVAL

As per international standard or university standard guideline Patient's consent and ethical approval has been collected and preserved by the authors.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Fitzgerald J, Broehm C, Chafey D, Treme G. Chondroblastoma of the Knee Treated with Resection and Osteochondral Allograft Reconstruction [Internet]. Vol. 2014, Case Reports in Orthopedics. Hindawi. 2014 [cited 2021 Feb 15];e543959.
2. Limaïem F, Tafti D, Rawla P. Chondroblastoma. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2020 [cited 2021 Feb 15]. Available: <http://www.ncbi.nlm.nih.gov/books/NBK536947/>
3. Yang Z, Tao H, Ye Z, Huang X, Lin N, Yang M, et al. The diagnosis and treatment of tibial intercondylar chondroblastoma. Clinics [Internet]. 2018 [cited 2021 Feb 15];73.
4. Daniel V, Daniel K. Exercises training program: It's effect on muscle strength and activity of daily living among elderly people. Nursing and Midwifery. 2020;1(01):19-23.
5. Xiong Y, Lang Y, Yu Z, Liu H, Fang X, Tu C, et al. The effects of surgical treatment with chondroblastoma in children and adolescents in open epiphyseal plate of long bones. World J Surg Oncol [Internet]. 2018 [cited 2021 Feb 15];16. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5781291/>
6. Andrade Neto F, Teixeira MJD, Araújo LH do C, Ponte CEB. Knee bone tumors:

- findings on conventional radiology. *Radiol Bras.* 2016;49(3):182–9.
7. Jr, BFP, Federico R. Tewes. What attorneys should understand about Medicare set-aside allocations: How Medicare Set-Aside Allocation Is Going to Be Used to Accelerate Settlement Claims in Catastrophic Personal Injury Cases. *Clinical Medicine and Medical Research.* 2021; 2(1):61-64. Available:<https://doi.org/10.52845/CMMR/2021v1i1a1>
 8. Jerome TJ, Varghese M, Sankaran B, Thomas S, Thirumagal SK. Tibial Chondroblastic Osteosarcoma—Case Report. *Foot Ankle Surg.* 2009;15(1):33–9.
 9. Wittig JC, Kellar-Graney K, Shmookler B. Osteosarcoma: A Multidisciplinary approach to diagnosis and treatment. 2002;65(6):10.
 10. Punzalan M, Hyden G. The role of physical therapy and occupational therapy in the rehabilitation of pediatric and adolescent patients with osteosarcoma. In: Jaffe N, Bruland OS, Bielack S, editors. *Pediatric and Adolescent Osteosarcoma* [Internet]. Boston, MA: Springer US; 2009 [cited 2021 Feb 17];367–84. (Cancer Treatment and Research; vol. 152). Available:http://link.springer.com/10.1007/978-1-4419-0284-9_20
 11. Daniel V, Daniel K. Perception of nurses' work in psychiatric clinic. *Clinical Medicine Insights,* 2020;1(1):27-33. Available: <https://doi.org/10.52845/CMI/2020v1i1a5>
 12. Almstedt HC, Grote S, Korte JR, Perez Beaudion S, Shoepe TC, Strand S, et al. Combined aerobic and resistance training improves bone health of female cancer survivors. *Bone Rep.* 2016;5:274–9.
 13. Gerritsen JKW, Vincent AJPE. Exercise improves quality of life in patients with cancer: a systematic review and meta-analysis of randomised controlled trials. *Br J Sports Med.* 2016;50(13):796–803.
 14. Morri M, Raffa D, Barbieri M, Ferrari S, Mariani E, Vigna D. Compliance and satisfaction with intensive physiotherapy treatment during chemotherapy in patients with bone tumours and evaluation of related prognostic factors: An observational study. *Eur J Cancer Care (Engl).* 2018;27(6):e12916.
 15. Daniel V, Daniel K. Diabetic neuropathy: New perspectives on early diagnosis and treatments. *Journal of Current Diabetes Reports.* 2020;1(1):12–14. Available: <https://doi.org/10.52845/JCDR/2020v1i1a3>
 16. Goyal C, Naqvi WM, Sahu A. An atypical case of febrile infection-related epilepsy syndrome following acute encephalitis: impact of physiotherapy in regaining locomotor abilities in a patient with neuroregression. *Pan Afr Med J* [Internet]. 2020 Jun 17 [cited 2021 Mar 14];36. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7392866/>
 17. Jensen W, Bialy L, Ketels G, Baumann FT, Bokemeyer C, Oechsle K. Physical exercise and therapy in terminally ill cancer patients: A retrospective feasibility analysis. *Support Care Cancer.* 2014;22(5):1261–8.
 18. Wane M, Naqvi WM, Vaidya L, Kumar K. Kinesiophobia in a Patient With Postoperative Midshaft Fracture: A Case Report of Its Impact on Rehabilitation in a 16-Year-Old Girl. *Cureus* [Internet]. 2020 Nov 5 [cited 2020 Dec 11];12(11). Available:<https://www.cureus.com/articles/42869-kinesiophobia-in-a-patient-with-postoperative-midshaft-fracture-a-case-report-of-its-impact-on-rehabilitation-in-a-16-year-old-girl>
 19. Strong DP, Grimer RJ, Carter SR, Tillman RM, Abudu A. Chondroblastoma of the femoral head: management and outcome. *Int Orthop.* 2010;34(3):413–7.
 20. Vaidya L, Kumar K, Naqvi W, Narang S, Pisulkar G, Dadlani M. Revision of total hip replacement surgery in elderly patient and its recovery based on periprosthetic fracture rehabilitation. 2020;11.
 21. Bawiskar D, Dhote S, Phansopkar P. Early physical rehabilitation post-surgery in a complex type 5 Schatzker Tibial plateau fracture improves functional outcomes: A case report. 2020;8.
 22. Goyal C, Naqvi W, Sahu A. Xia-Gibbs Syndrome: A Rare Case Report of a Male Child and Insight into Physiotherapy Management. *Cureus* [Internet]. [cited 2021 Mar 14];12(8). Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7478925/>
 23. Lane JM, Christ GH, Khan SN, Backus SI. Rehabilitation for limb salvage patients. *Cancer.* 2001;92(S4):1013–9.

24. Khatib MN, Shankar AH, Kirubakaran R, Gaidhane A, Gaidhane S, Simkhada P, Quazi SZ. Ghrelin for the management of cachexia associated with cancer. Cochrane Database of Systematic Reviews. 2018;2(2018). Available:<https://doi.org/10.1002/14651858.CD012229.pub2>.

© 2021 Dhote et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/71126>