



First Report of Common Bean (*Phaseolus vulgaris* L.) Root Rot Caused by *Rhizoctonia Solani* in Jammu and Kashmir, India

**Shafat Ahmad Ahanger^{a*}, Sabiya Bashir^b, Mohammad Najeeb Mughal^c,
Rouf Ahmad Rather^a, Reyaz ul Rouf Mir^d, Muntazir Mushtaq^e
and Umer Basu^{f*}**

^a Division of Plant Pathology, Faculty of Agriculture, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, Wadura, Sopore-193201, India.

^b Dryland Agriculture Research Station, Rangreth Srinagar, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, 191132, India.

^c Division of Plant Pathology, Faculty of Horticulture, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, Shalimar-190025, India.

^d Division of Genetics and Plant Breeding, Faculty of Agriculture, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, Wadura, Sopore -193201, India.

^e Division of Germplasm Evaluation, ICAR-National Bureau of Plant Genetic Resources, Pusa Campus, New Delhi - 110012, India.

^f Division of Plant Pathology, Sher-e-Kashmir University of Agricultural Sciences and Technology of Jammu, 180009, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2022/v12i1131168

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/91263>

Original Research Article

Received 14 June 2022

Accepted 25 August 2022

Published 26 August 2022

ABSTRACT

Root rot of common bean (*Phaseolus vulgaris*) is one of the most serious disease, causes unadorned damage to crop health and leads to the heavy yield loss. In the present investigation, *Rhizoctonia solani*, a new root rot pathogen was found pathogenic and cause of beans root rot in the region of Jammu and Kashmir, India. The pathogen was characterized by morpho-cultural as well as through molecular sequencing using ITS primer for establishing the proper identification of

isolated pathogen. Based on the pathogenicity test carried on common bean cv. Shalimar Bean-1 and available literature, it was recorded as the first report of root rot in common beans caused by *Rhizoctonia solani* in Jammu and Kashmir.

Keywords: ITS primers; pathogenicity; *Phaseolus vulgaris*; *Rhizoctonia solani*; root rot.

1. INTRODUCTION

Common bean (*Phaseolus vulgaris* L.) is a regularly consumed crop, thereby occupying a prime position among the legumes in the world including India. Diseases are becoming more frequent encountered constraints in the successful production of common bean. Among important diseases, soil borne pathogenic fungi causing root rot/damping off are widely distributed and causing serious damage to the crop that led to heavy yield losses [1]. The main soil borne pathogens viz., *Fusarium solani*, *Pythium ultimum*, *Rhizoctonia solani* [2-4] and *Macrophomina phaseolina* [5], which are responsible for fusarium root rot (FRR), pythium seed rot/damping off, Rhizoctonia root rot and Macrophomina charcoal rot respectively, and have been reported world-wide. In Kashmir, root rot of beans has been reported to be caused mainly by *Fusarium solani* [6,7]. However, in present study *Rhizoctonia solani* was found pathogenic and cause beans root rot.

2. MATERIALS AND METHODS

2.1 Symptomatology

The root rot samples of common bean were collected from various districts viz., Anantnag, Baramulla and Pulwama of Kashmir valley of UT of Jammu and Kashmir. The symptoms of the infected plants showed dwarfing and yellowing on above ground plant and irregular to circular sunken scaled lesions with reddish brown to black borders on roots and hypocotyl region. The infected root system was devoid of lateral roots. The pathogen destroyed the pith which resulted in a brick-red discoloration of interior part of stem near the soil line (Fig. 1A and 1B).

2.2 Isolation and Purification

The pathogen was isolated from the symptomatic roots samples using tissue bit isolation technique of Sicard et al. [8]. The suspected tissue bits were inoculated on potato dextrose agar plates and were incubated at 25±1°C for seven days.

Hypal tip technique was used to obtain the axenic culture of pathogen [9].

2.3 Pathogenicity Test

To ascertain ability of the isolate to cause root rot in bean plants, pathogenicity tests was carried on common bean cv. Shalimar Bean-1. Sick-soil method was employed for performing the pathogenicity test [10]. Plastic pots (15 cm height) were filled with sterilized soil (1.5 kg pot⁻¹) and inoculated with mass culture of the pathogen @ 10g kg⁻¹ soil, five days prior to sowing. One un-inoculated pot filled with sterilized soil of same quantity was kept as a control. Three apparently healthy bean seeds, first surface sterilized by 1 per cent sodium hypochlorite for two minutes followed by 3 to 4 washings with sterile distilled water, were sown in each pot. Both the inoculated and un-inoculated pots were kept under poly-house for observations on disease development.

2.4 Genomic DNA Extraction

Genomic DNA was extracted by using modified CTAB (*Cetyltrimethyl ammonium bromide*) method (Murray and Thompson, 1980). Two ITS primers viz., ITS1F (5'TCCGTAGGTGAACCTGCG'3) and ITS4R (3'TCCTCCGCTTATTTGATATGC'5) were used for sequence analysis of the root rot pathogens. The PCR cycle for ITS primers with initial denaturation at 94°C for 5 minute, followed by 35 cycles of denaturation at 94°C for 30 seconds, annealing at 58°C for 1 minute, extension at 72°C for 2 minutes with final elongation of 72°C for 10 minutes was performed. After PCR amplification, 40 to 45 µl PCR product of root rot pathogen was sent for custom sequencing (Bio Kart India Pvt. Ltd. Bengaluru, Karnataka, India). ITS sequence of pathogen was retrieved from chromatograms using BioEdit version 7.0 (Hall, 1999) and the sequenced product was subjected to BLASTn in NCBI (National Centre of Biotechnology Information) database and identification of pathogen was established.



Fig. 1. Symptoms of root rot observed under field condition; A: Rotting symptoms developed on roots, B: Above ground plant symptoms due to root rot

3. RESULTS AND DISCUSSION

The pathogen was identified by morpho-cultural as well as through molecular sequencing. The pure culture of *R. solani* produced a profuse white aerial mycelium which turned to brown to dark brown in colour after seven days of incubation (Fig. 2A and 2B). Hyphal branches of pathogen were formed at right angle from the main hyphae with prominent constriction and a septum at point of origin (Fig. 2C and 2D). The diameter of hyphae varied from 6.60 to 7.78 μ m with an average of 7.01 μ m. Sclerotia were small to large in size and developed on upper surface of media in circular manner (target board) after five days of incubation. These were 95 to 160 x 62 to 145 μ m with an average of 81.5 to 107 μ m in size, spherical/oval, dirty white and later turning to dark brown. Monilliod cells were formed either singly or in chains, barrel shaped, thick walled, initially hyaline and then turned to brown in

colour and measuring 18.01 to 23.45 x 8.2 to 13.7 μ m in size. On the basis of mycelial characters, sclerotia and monilliod cells, the fungus was identified as *Rhizoctonia solani* [11].

To confirm the precise identity of the pathogen, the PCR amplification was carried out for Internal Transcribed Spacer (ITS) region of the ribosomal DNA with primers ITS1/IST4 [12]. The sequence was retrieved from the chromatogram and was analyzed through Basic Local Alignment Search Tool-Nucleotide (BLASTn) programme in NCBI (National Centre for Biotechnology Information) database (ncbi.nlm.nih.gov/genbank) and identification of pure cultures was established. The sequence of the isolated pathogen showed 98 per cent similarity with reference *Rhizoctonia solani* sequences in Gen Bank (Fig. 3). The sequences were submitted in the Gen Bank under accession no: MZ708012.




Fig. 2. Morphological and cultural characterization of *Rhizoctonia solani* causing root rot of common bean. (A) Front view of *Rhizoctonia solani* culture, (B) Back view of *Rhizoctonia solani* culture, (C) Branches from the main hyphae formed at right angle with prominent constriction at point of origin (D) Monilliod cells formed in chains




Fig. 3. Phylogenetic analysis of the original isolate by using the Maximum Likelihood method and Kimura 2-parameter mode on Internal Transcribed Spacer (ITS sequence).The position of bean root rot pathogen *Rhizoctonia solani* (SKU-W-BRR2) is depicted with red (♦) colour



Un-inoculated Inoculated with *Rhizoctonia solani*



Control Symptoms of *Rhizoctonia solani*

Fig. 4. Pathogenicity test of isolated fungal pathogen *Rhizoctonia solani* on common bean cv. Shalimar Bean-1 under pot culture

Pathogenicity of isolated pathogen was carried on common bean cv. Shalimar Bean-1 under pot culture experiment by adopting the sick soil method [10]. These pots were maintained under greenhouse with 12 hours of photoperiod at 25±1°C. The pathogen inoculated plants showed the development of characteristic symptoms viz., yellowing and drying of leaves after 16 days of inoculation and the roots showed irregular to circular sunken lesions with reddish brown-black margins on roots, rotted similar to those originally detected under field conditions. No symptoms development was observed in controlled plants (Fig. 4). The pathogen was also reported to cause disease in other crops [13-16]. But to our knowledge and on the basis of available literature, this is the first report of common beans root rot caused by *Rhizoctonia solani* in Kashmir. Thus it becomes important to study the impacts of this new disease on common bean production for the potential infestation of the pathogen in Kashmir region.

4. CONCLUSION

Based on the morpho-cultural characteristics of the pathogen and pathogenicity test carried on common bean cv. Shalimar Bean-1 and available literature, it was recorded as the first report of root rot in common beans caused by *Rhizoctonia solani* in Jammu and Kashmir, India.

ACKNOWLEDGEMENT

The author acknowledges the support and help of all the members of Division of Plant Pathology, SKUAST-K.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Panth M, Hassler SC, Baysal-Gurel F. Methods for management of soilborne diseases in crop production. *Agriculture*. 2020;10(1):16.
- Singh SP, Schwartz HF. Breeding common bean for resistance to diseases: a review. *Crop Science*. 2010;50(6):2199-223.
- Scandiani MM, Ruberti DS, Giorda LM, Pioli RN, Luque AG, Bottai H, Ivancovich JJ, Aoki T, O'Donnell K. Comparison of inoculation methods for characterizing relative aggressiveness of two soybean sudden-death syndrome pathogens, *Fusarium virguliforme* and *F. tucumaniae*. *Tropical Plant Pathology*. 2011;36:133-40.
- Vural C, Soylu S. Prevalence and incidence of fungal disease agents affecting bean (*Phaseolus vulgaris* L.) plants. *Research on Crops*. 2012;13(2): 634-40.
- Clare MM, Melis R, Dereta J, Laing M, Buruchara RA. Identification of sources of resistance to *Fusarium* root rot among selected common bean lines in Uganda. *Journal of Animal & Plant Sciences*. 2010;7(3):876-91.
- Hassan Dar GH, Zargar MY, Beigh GM. Biocontrol of *Fusarium* root rot in the common bean (*Phaseolus vulgaris* L.) by using symbiotic *Glomus mosseae* and *Rhizobium leguminosarum*. *Microbial Ecology*. 1997;34(1):74-80.
- Badri ZA, Bhat NA, Raina FA, Wani TA, Yousuf V. Status of late blight disease (*Phytophthora infestans*) of potato at gurez in Jammu and Kashmir. *Indian Phytopathology*. 2014;67(2):164-7.
- Sicard D, Michalakis Y, Dron M, Neema C. Genetic diversity and pathogenic variation of *Colletotrichum lindemuthianum* in the three centers of diversity of its host, *Phaseolus vulgaris*. *Phytopathology*. 1997; 87(8):807-13.
- Rangaswami G. Diseases of Crop Plants in India. Prentice Hall of India. New Delhi. 1972:520.
- Dubey SC, Aradhika T, Birendra S. Combination of soil application and seed treatment formulations of *Trichoderma* species for integrated management of wet root rot caused by *Rhizoctonia solani* in chickpea (*Cicer arietinum*). *Indian Journal of Agricultural Sciences*. 2012;82(4): 356-62.
- Parmeter JR, Whitney HS. Taxonomy and nomenclature of the imperfect stage In: *Rhizoctonia solani* Biology and pathology. *Journal Research of University of California*. 1970;7-19.
- White TJ, Bruns T, Lee SJ, Taylor J. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR protocols: a Guide to Methods and Applications*. 1990;18(1): 315-322.
- Venegas P. Identification of rust resistance and a molecular marker in a cross within tertiary gene pool of common bean and characterization of *Rhizoctonia* Spp.

- Isolates from Western Nebraska (MS thesis) University of Nebraska-Lincoln: 141.
14. Aliferis, Konstantinos A, Suha Jabaji. "FT-ICR/MS and GC-EI/MS metabolomics networking unravels global potato sprout's responses to *Rhizoctonia solani* Infection. 2012:e42576.
 15. Paparu P, Acur A, Kato F, Acam C, Nakibuule J, Musoke S, Nkalubo S, Mukankusi C. Prevalence and incidence of four common bean root rots in Uganda. *Experimental agriculture*. 2018;54(6): 888-900.
 16. Tolêdo-Souza ED, Júnior ML, da Silveira PM, Café Filho AC. Interactions between *Fusarium solani* f. sp. *phaseoli* and *Rhizoctonia solani* on the severity of dry bean root rot. *Pesquisa Agropecuária Tropical*. 2009;39(1):13.

© 2022 Ahanger et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/91263>