



Inadvertent Gallbladder Stenting During Endoscopic Management of Choledocholithiasis Detected During Laparoscopic Cholecystectomy: A Case Report

Jayant Kumar Biswal^{a++}, Sujit Kumar Mohanty^{a#},
Suraj Ethiraj^{a*†} and Satyanarayan Behera^{a#}

^a Department of Surgery, SCB Medical College and Hospital, Cuttack, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/95737>

Received: 20/11/2022

Accepted: 28/01/2023

Published: 03/02/2023

Case Study

ABSTRACT

Endoscopic retrograde cholangiopancreatography (ERCP) is the standard management protocol for choledocholithiasis, which more often than not involves a placement of stent in the common bile duct. Here we present a case of inadvertent stenting of the cystic duct and gallbladder during ERCP for choledocholithiasis which was detected during laparoscopic cholecystectomy. A 38 year old lady presented with jaundice and right upper quadrant pain, diagnosed with choledocholithiasis, underwent ERCP with stenting and stone extraction. After 6 weeks, she underwent laparoscopic

⁺⁺ Professor;

[#] Assistant Professor;

[†] Postgraduate resident;

*Corresponding author: E-mail: surajethiraj@gmail.com;

cholecystectomy, when it was noticed that the biliary stent was inadvertently placed into the cystic duct and gallbladder. The stent was removed laparoscopically through a rent in the gallbladder and cystic duct stump clipped. We report this case to enhance awareness amongst surgeons regarding the unintended location of a biliary stent.

Keywords: Gallbladder stenting; choledocholithiasis; cholecystectomy; ERCP.

1. INTRODUCTION

The prevalence of gall stones is quite high in the general population, being as high as 15% [1]. Of these, 10-18% are found to have common bile duct (CBD) stones [1]. Most CBD stones are secondary having migrated from the gallbladder. A possibility of CBD stone must be excluded prior to laparoscopic cholecystectomy. Since the introduction of endoscopic retrograde cholangiopancreatography (ERCP) over 40 years ago, the ability to tackle CBD stones with minimal complications has improved [2]. Though complications such as pancreatitis, haemorrhage, viscous perforation exists following ERCP, but those of stent migration, wrongful stenting into the cystic duct are quite rare [3].

Here we present a rare case of inadvertent biliary stenting of the cystic duct and gallbladder following ERCP for choledocholithiasis, which was detected during subsequent laparoscopic cholecystectomy.

2. CASE PRESENTATION

A 38 year old lady presented with right upper abdominal pain and jaundice since 10 days with similar complaints in the past. On examination

she was icteric, with normal vital signs. There was tenderness over the right hypochondrium with no features of peritonitis. Abdominal ultrasonography and magnetic resonance cholangiopancreatography (MRCP) revealed cholelithiasis and choledocholithiasis with a distal CBD calculus of size 12 mm as showed in Fig. 1. She underwent ERCP with stone extraction and biliary stenting with no adverse events. Following the procedure patient was relieved of symptoms and discharged to follow up for cholecystectomy.

She was readmitted after 6 weeks with the plan for laparoscopic cholecystectomy. Repeat ultrasonography revealed contracted gallbladder with cholelithiasis and no CBD stones and stent in situ in the CBD. She underwent the standard 4 port laparoscopic cholecystectomy. Intraoperatively, there was minimal adhesions in the region of Calot's triangle as shown in Fig. 2. During Calot's dissection, the cystic duct was found rigid. It was observed that the biliary stent was present within the cystic duct and entering into the gallbladder. It was removed via a rent in the gallbladder as shown in Fig. 3, following which the cholecystectomy was completed. Postoperative period was uneventful and patient was discharged to follow up on post operative day 3.




Fig. 1. MRCP showing CBD calculus



Fig. 2. Intraoperative photograph showing adhesions in the region of Calot's triangle




Fig. 3. Intraoperative photograph showing the removal of the stent through the gallbladder

3. DISCUSSION

ERCP with biliary stenting is now the standard of care as the first line management for certain cases of choledocholithiasis. It involves either a sphincterotomy or balloon dilation of the common bile duct followed by sweeping of the stones [2]. A biliary stent is usually placed to prevent recurrence or stricture. Such patients after ERCP are candidates for laparoscopic cholecystectomy either within the first 48 hours or after 6 weeks, although there are varied guidelines [4]. With many documented complications of ERCP, some are rare and less understood [5]. Of these inadvertent biliary stenting into the cystic duct and gallbladder is one such complication with none to very few cases reported [6]. Stent migration, another known complication of ERCP is not very common. Stent migration can occur either proximally into the hepatic ducts or distally into the intestinal lumen [7, 8].

Our case describes this complication of inadvertent biliary stenting into the cystic duct

and gallbladder, which was later detected during laparoscopic cholecystectomy. Although this is not a fatal complication, it is important for the surgeon to be aware of such occurrences as it can pose trouble while ligating the cystic duct during cholecystectomy. Should a surgeon encounter such a finding intraoperatively, it is best tackled by stent extraction laparoscopically by a rent in the gallbladder, followed by completing the cholecystectomy as depicted in our case report. If identified preoperatively by means of any radiological investigation, a repeat ERCP and stenting may be tried to reposition the stent without any adverse consequences.

4. CONCLUSION

Awareness among surgeons regarding the complication of inadvertent biliary stenting into the cystic duct and gallbladder is necessary to avoid intraoperative surgical surprises during interval laparoscopic cholecystectomy following ERCP for choledocholithiasis.

CONSENT

As per international standard or university standard, patient(s) written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Martin DJ, Vernon DR, Toouli J. Surgical versus endoscopic treatment of bile duct stones. *Cochrane Database Syst Rev*. 2006;CD003327.
2. Al-Haddad M. ERCP for common bile duct stone extraction: Sphincterotomy, balloon dilation, or both?. *Saudi J Gastroenterol*. 2015;21:181-2
3. Freeman ML, Nelson DB, Sherman S, Haber GB, Herman ME, Dorsher PJ, Moore JP, Fennerty MB, Ryan ME, Shaw MJ, Lande JD, Pheley AM. Complications of endoscopic biliary sphincterotomy. *N Engl J Med*. 1996;335(13):909-18.
4. Friis C, Rothman JP, Burcharth J, Rosenberg J. Optimal timing for laparoscopic cholecystectomy after endoscopic retrograde cholangiopancreatography: a systematic review. *Scand J Surg*. 2018;107(2):99-106. DOI:10.1177/1457496917748224. Epub 2017 Dec 26. PMID: 29277136.
5. Talukdar R. Complications of ERCP. *Best Pract Res Clin Gastroenterol*. 2016;30(5):793-805. DOI: 10.1016/j.bpg.2016.10.007. Epub 2016 Oct 27. PMID: 27931637.
6. Yagnik VD, Patel A, Mannari GM, Garg P, Dawka S. Migration of biliary stent into the gallbladder: A surprising intraoperative finding. *J Minim Access Surg*. 2022;18(1):151-153. DOI: 10.4103/jmas.JMAS_47_21. PMID: 35017405; PMCID: PMC8830582.
7. Morimachi M, Ogawa M, Yokota M, Kawanishi A, Kawashima Y, Mine T. Successful endoscopic removal of a biliary stent with stent-stone complex after long-term migration. *Case Rep Gastroenterol*. 2019;13:113-7.
8. Márquez HR, Sanchez JS, Jaimes ES. Proximal migration of biliary stent: Case report. *EC Gastroenterol Dig Syst*. 2019;6:155-62.

© 2023 Biswal et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/95737>