



## Effect of Different Fertility Management Practices on Plant Population and Mortality in Chickpea (*Cicer arietinum* L.)

Pooja Bhatt <sup>a\*</sup>, V. K. Singh <sup>a#</sup>, Rohitashv Singh <sup>a#</sup>, Naresh Malik <sup>a#</sup>  
and Ramesh Chandra <sup>a#</sup>

<sup>a</sup> Department of Agronomy, College of Agriculture, GBPUAT, Pantnagar 263145, Uttarakhand, India.

### Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/IJPSS/2022/v34i2131323

### Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/89985>

Original Research Article

Received 20 May 2022  
Accepted 21 July 2022  
Published 25 July 2022

## ABSTRACT

**Aims:** To assess the effect of Humic acid (HA), Plant growth-promoting bacteria (PGPR), and Recommended dose of fertilizer (RDF) alone or in combination on plant population and mortality percentage in Chickpea (*Cicer arietinum* L.).

**Study of Design:** The experiment was laid out in Randomized block design (RBD) with three replications.

**Place and Duration:** A field experiment was conducted during the *rabi* season in the year 2020-21 and 2021-2022 at D<sub>6</sub> Block, Norman E. Borlaug Crop Research Center at G. B. Pant University of Agriculture and Technology, Pantnagar, Uttarakhand (India).

**Treatments Details:** Total ten treatments were used. The details of the treatments are as follows: T<sub>1</sub>- Absolute control, T<sub>2</sub> - Humic acid @ 2.5 kg/ha (soil application), T<sub>3</sub> -Humic acid @ 20 g/kg of seed (seed treatment), T<sub>4</sub> - PGPR @ 20 g/kg of seed (seed treatment), T<sub>5</sub>- RDF through 150 kg NPK mixture (12:32:16 Grade), T<sub>6</sub> -RDF + HA (soil application), T<sub>7</sub> -RDF +HA (seed treatment), T<sub>8</sub>- RDF +PGPR (Seed treatment), T<sub>9</sub>- HA (soil application) +PGPR (seed treatment), T<sub>10</sub>- RDF + HA (soil application)+PGPR (seed treatment).

**Methodology:** For plant population, the number of plants that emerged in the second row from both sides of the field were counted at 30 DAS and at harvest stage then averaged and calculate

\*Ph.D. Scholar;

#Professor;

\*Corresponding author: E-mail: [bhattpooja1991@gmail.com](mailto:bhattpooja1991@gmail.com);

the number of plants for 1 m<sup>2</sup> area. The mortality percentage was calculated by using the following formula:

$$\text{Plant mortality (\%)} = \frac{\text{No. of plants at 30 DAS} - \text{No. of plants at harvest}}{\text{No. of Plants at 30 DAS}} \times 100$$

**Results:** Findings of the investigation revealed that the effect of different fertility management on plant population was found not significant, but it was found significant in mortality percentage. Among the sole application of treatments T<sub>2</sub> (Humic acid @ 2.5 kg/ha (soil application) and T<sub>4</sub> (PGPR @ 20 g/kg of seed (seed treatment) recorded minimum mortality percentage, while among the combination treatments, T<sub>10</sub> (RDF+HA (soil application) + PGPR as seed treatment) recorded minimum percentage of mortality.

**Keywords:** Humic acid; plant growth-promoting bacteria; chickpea.

## 1. INTRODUCTION

Chickpea (*Cicer arietinum* L.) is one of the first grain legumes domesticated by mankind. It is an important *rabi* pulse in India. Chickpea is the world's most significant legume crop, and its seed contains 16-31 percent protein, 3 percent fiber, 38-73 percent carbohydrate, 0.3 percent Phosphorus, vitamins (B and C), and minerals (Zn, Fe, K, Mg) [1]. In India, chickpea occupied 9.85 Mha with a production of 11.99 Mt and a 1217 kg/ha productivity. Madhya Pradesh, Rajasthan, and Maharashtra are India's top chickpea-producing [2]. Chickpea belongs to a *Leguminosae* family, which can fix atmospheric nitrogen (N<sub>2</sub>) in association with Rhizobium bacteria. For the proper germination of chickpea, the edaphic and climatic conditions should be favorable. Uniform plant population is directly correlated with the yield. A high mortality rate can reduce the economic yield of chickpeas. Some sustainable approach needs to adapt to minimize the mortality percentage and improve the soil fertility status.

Humic acid is a dark black substance resistant to further decomposition made up of plant and animal materials through microbial degradation. Leonhardite (sedimentation layers), an oxidation result of lignite, is a key source of humic acid [3]. Humic acid consists of 51 to 57% Carbon, 4 to 6% Nitrogen, and 0.2 to 1.0% Phosphorus and other terrace nutrients in minute amounts. Humic acid directly enhances plant growth by improving the soil's physical, chemical & biological properties and indirectly by increasing chlorophyll content, membrane permeability, and respiration in plants [4]. Humic acid's structure comprises a wide range of compounds that cannot be defined by a single structural formula; hence its molecular weight ranges from 1000 to 30,000. HA provides more surface area and possesses more negative charge resulting in more water retention and Cation-exchange capacity of the

soil [5]. It is made up of chemically composite non-biochemical organic compounds, which improve soil health and balanced plant nutrition. Thus, it is important for soil and plant productivity [6]. It is also tolerant of abiotic stress [7].

Plant growth-promoting rhizobacteria (PGPR) are a diverse set of bacteria found in the rhizosphere, on root surfaces, and in close proximity to roots that can, directly and indirectly, promote the growth and development of plants [8]. PGPR has been found to release siderophores, which are extracellular metabolites. Microbial Fe-chelating low molecular weight molecules are referred to as siderophores. The siderophores help to facilitate the accessibility of Fe<sup>+3</sup> to plants and protect the plant from soil-borne pathogens, which improves plant development and agricultural productivity [9]. Rhizobium inoculant applied directly to planting furrows aids germination and nitrogen fixation of foreign strains [10].

The rising population of the world has become a severe issue in respect to soil health due to the vast cultivation of crops for fulfilling the requirements of the present and future generation. In this approach, the soil mass has been widely examined without thinking for its health. If we can find some alternative to inorganic fertilizers, we can protect the soil. Two organic sources i.e humic acid and PGPR are being studied here to evaluate the influence on crop performance.

## 2. MATERIALS AND METHODS

The present study was conducted during the *rabi* season in 2020-21 and 2021-2022 at Norman E. Borlaug, a Crop research center at G.B. Pant University of agriculture and technology, district Udham Singh Nagar, Uttarakhand, India. The location is situated at 29° N latitude, 79.5° E longitude and an altitude of 243.83 m above

mean sea level in the foothills of the Himalayas. The soil order of this region is Mollison, and the texture is sandy loam. Soil has a poor or moderately drain condition. The climate is humid and subtropical.

The experiment was laid out in a Randomized block design with three replications. The treatments included the application of humic acid, PGPR, and RDF alone or in combination. The details of the treatments are as follows: T<sub>1</sub>- Absolute control, T<sub>2</sub> - Humic acid @ 2.5 kg/ha (soil application), T<sub>3</sub> -Humic acid @ 20 g/kg of seed (seed treatment), T<sub>4</sub> . PGPR I @ 20 g/kg of seed (seed treatment), T<sub>5</sub>- RDF through 150 kg NPK mixture (12:32:16 Grade), T<sub>6</sub> -RDF + HA (soil application), T<sub>7</sub> -RDF +HA (seed treatment), T<sub>8</sub>- RDF +PGPR (Seed treatment), T<sub>9</sub>- HA (soil application) + PGPR (seed treatment), T<sub>10</sub>- RDF + HA (soil application)+PGPR (seed treatment). There were a total 30 plots, and each plot size was 18 m<sup>2</sup>. The crop was sown at 30x10 cm spacing. PG-186 variety of chickpea was used in the experiment.

### 2.1 Sampling and Calculation

For plant population, the number of plants that emerged in the second row from both sides of the field were counted at 30 DAS and at harvest stage then averaged and calculate the number of plants for 1 m<sup>2</sup> area. The mortality percentage was calculated by using the following formula:

$$\text{Plant mortality (\%)} = \frac{\text{No. of initial plants} - \text{No. of plants at harvest}}{\text{No. of initial plants}} \times 100$$


### 3. RESULTS AND DISCUSSION

The data presented in Table 1 revealed that the plant population at both stages do not significantly affect by fertility management practices during both years. At harvest, the plant population during both years was reduced in all the treatments. The effect of treatments on mortality percentage was found to be significant during both years. In the first year of the experiment, the combination treatments, i.e., T<sub>6</sub>, T<sub>7</sub>, T<sub>8</sub>, T<sub>9</sub>, and T<sub>10</sub>, were found at par and were significantly superior to the sole application of treatments while in the second year, T<sub>6</sub>, T<sub>9</sub>, and T<sub>10</sub> were found at par. The lowest mortality was recorded in T<sub>10</sub> (RDF+HA+PGPR), and the highest mortality was observed in T<sub>1</sub> (Control) Fig. 1. It may be due to the effect of combining humic acid, PGPR, and RDF. Among the sole application of humic acid and PGPR treatments, PGPR using treatment recorded lower mortality in the first year. Humic acid (T<sub>2</sub>) soil application recorded lower mortality in the second year. Among the combination treatments, T<sub>6</sub> obtained a lower percentage of mortality in both the years.

Among the sole application of treatments, T<sub>2</sub>, T<sub>3</sub>, T<sub>4</sub>, and T<sub>5</sub> recorded 14.51%, 9.38%, 22.41%, and 11.12% less mortality in the first year and 22.15%, 10.56%, 18.42%, and 15.26% in the second year, respectively as compared to T<sub>1</sub>(control). Among the combination treatments, humic acid (soil application) and PGPR using treatments recorded the lowest mortality. This might be due to the application of HA which improves the soil structure by increasing the aggregation between the soil particles, which facilitates better aeration and provides strength

**Table 1. Effect of different fertility management practices on plant population and mortality percentage in chickpea**

Treatment	Plant Population/m <sup>2</sup>				Mortality %	
	At 30 DAS		At harvest		2020-21	2021-22
	2020-21	2021-22	2020-21	2021-22		
T <sub>1</sub>	34.0	34.0	27.3	27.3	19.5	19.5
T <sub>2</sub>	34.0	35.0	28.3	29.6	16.6	15.2
T <sub>3</sub>	33.0	34.3	27.3	28.3	17.6	17.5
T <sub>4</sub>	35.0	33.6	29.6	28.3	15.1	15.9
T <sub>5</sub>	34.0	34.3	28.0	28.6	17.3	16.6
T <sub>6</sub>	33.0	35.3	29.3	31.3	11.0	11.4
T <sub>7</sub>	33.0	32.6	28.6	28.0	12.7	14.4
T <sub>8</sub>	34.6	33.0	30.3	28.6	12.4	13.1
T <sub>9</sub>	35.0	35.0	30.6	30.6	12.4	12.2
T <sub>10</sub>	33.0	33.0	29.6	30.0	10.1	9.2
<b>S. Em±</b>	0.9	1.0	1.2	1.3	1.1	1.1
<b>C.D. (P=0.05)</b>	NS	NS	NS	NS	3.5	3.5



**Fig. 1. Effect of different fertility management practices on mortality percentage in chickpea**

to the soil to hold the plant against abnormal weather conditions [11]. On the other hand, PGPR releases some antioxidant which prevents the soil-borne diseases and also provides the favorable environment to the crop by regulating the rhizobacteria [9] similar results were obtained by [12].

#### 4. CONCLUSION

This investigation concluded that the application of humic acid or PGPR along with RDF could reduce the mortality percentage and contribute to more grain yield in chickpeas. As humic acid and PGPR are organic substances that help improve the soil properties and enhance the microbial population in soil on a long-term basis. Application of humic acid and PGPR can reduce the amount of fertilizer that may sustainably protect the soil. This approach can reduce the fertilizer and lead to a step forward in sustainable agriculture.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Özer SS, Karaköy T, Toklu F, Baloch FS, Kilian B, Özkan H. Nutritional and

physicochemical variation in Turkish kabuli chickpea (*Cicer arietinum* L.) landraces. *Euphytica*. 2010;175(2):237–249.

Available: <https://doi.org/10.1007/S10681-010-0174-3>

2. Directorate of Economics and Statistics, Ministry of Agriculture, Government of India; 2021.

Available: <https://eands.dacnet.nic.in/>

3. RW O'donnell - Soil Science, & 1973. (n.d.). The auxin-like effects of humic preparations from leonardite. *Journals.Lww.Com*. Retrieved July 18, 2022.

Available: [https://journals.lww.com/soilsci/Abstract/1973/08000/The\\_Auxin\\_Like\\_Effects\\_of\\_Humic\\_Preparations\\_From.7.aspx](https://journals.lww.com/soilsci/Abstract/1973/08000/The_Auxin_Like_Effects_of_Humic_Preparations_From.7.aspx)

4. Rajpar I, Bhatti M, A ZHPJ. of, & 2011, undefined. (n.d.). Humic acid improves growth, yield and oil content of *Brassica campestris* L. *Researchgate.Net*. Retrieved July 18, 2022.

Available: [https://www.researchgate.net/profile/Dr\\_Zia-ul-hassan\\_Shah/publication/230846587\\_Humic\\_acid\\_improves\\_growth\\_yield\\_and\\_oil\\_content\\_of\\_Brassica\\_compestris\\_L/links/0a85e52d9857a10a54000000/Humic-acid-improves-growth-yield-and-oil-content-of-Brassica-compestris-L.pdf](https://www.researchgate.net/profile/Dr_Zia-ul-hassan_Shah/publication/230846587_Humic_acid_improves_growth_yield_and_oil_content_of_Brassica_compestris_L/links/0a85e52d9857a10a54000000/Humic-acid-improves-growth-yield-and-oil-content-of-Brassica-compestris-L.pdf)

5. Rong Q, Zhong K, Huang H, Li C, Zhang C, Sciences XNA, & 2020, undefined. (n.d.). Humic acid reduces the available cadmium, copper, lead, and zinc in soil and their uptake by tobacco. Mdpi.Com. Retrieved July 18, 2022. Available:<https://www.mdpi.com/634112>
6. Adani F, Spagnol M, Genevini P. Biochemical origin and refractory properties of humic acid extracted from the maize plant. Biogeochemistry. 2006;78(1): 85–96. Available:<https://doi.org/10.1007/S10533-005-2902-7>
7. Shah F, WW. Soil and crop management strategies to ensure higher crop productivity within sustainable environments. Mdpi.Com; 2019. Available:<https://doi.org/10.3390/su11051485>
8. Bhattacharjya S, Research-An RCL, 2013, undefined. Effect of inoculation methods of Mesorhizobium ciceri and PGPR in chickpea (*Cicer arietinum* L.) on symbiotic traits, yields, nutrient uptake and soil properties. Indianjournals.Com; 2013. Available:<https://www.indianjournals.com/ijor.aspx?target=ijor:lr&volume=36&issue=4&article=010>
9. Verma JP, Yadav J, Tiwari KN. Enhancement of Nodulation and Yield of Chickpea by Co-inoculation of Indigenous *Mesorhizobium* spp. and Plant Growth-Promoting Rhizobacteria in Eastern Uttar Pradesh. Communications in Soil Science and Plant Analysis. 2012;43(3):605–621. Available:<https://doi.org/10.1080/00103624.2012.639110>
10. ES Jensen - Plant and, undefined. Inoculation of pea by application of Rhizobium in the planting furrow. Springer. 1987;97(1):63–70. Available:<https://doi.org/10.1007/BF02149824>
11. Chaney K, Swift RS. Studies on aggregate stability. 11. The effect of humic substances on the stability of re- formed soil aggregates. Journal of Soil Science. 1986;37(2):337–343. Available:<https://doi.org/10.1111/J.1365-2389.1986.TB00036.X>
12. Sahni S, Prasad BD. Management of collar rot disease using vermicompost and a PGPR strain *Pseudomonas* sp. and their effect on defense-related enzymes in chickpea. Indian Phytopathology. 2020; 73(2):301–311. Available:<https://doi.org/10.1007/S42360-020-00203-4>

© 2022 Bhatt et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/89985>