



## Determination of the Sexual Maturity of Threadfins *Polydactylus quadrifilis* (Cuvier, 1829), *Galeoides decadactylus* (Bloch, 1795) and *Pentanemus quinquarius* (Linné, 1758) of the Artisanal Marine Fishery of Grand-Lahou (Ivory Coast)

Kouassi Sylvain Konan<sup>1\*</sup>, Ebram Luc Gervais Djadji<sup>1</sup>,  
Yao Nicolas Amon<sup>2</sup> and Konan N'da<sup>3</sup>

<sup>1</sup>Department of Aquaculture, Oceanologic Research Centre P.O.Box V 18, Abidjan, Côte d'Ivoire.

<sup>2</sup>Department of Animal Biology, Peleforo Gon Coulibaly University, P.O.Box 1328, Korhgo, Côte d'Ivoire.

<sup>3</sup>Department of Nature Sciences, Nangui Abrogoua University, 02 P.O.Box 801, Abidjan, Côte d'Ivoire.

### Authors' contributions

This work was done with a frank collaboration among all the authors. Authors KSK and ELGD collected samples and designed the study then, author YNA performed the data processing and the all, drafted the first version of the manuscript. Author KN wrote the protocol and corrected the first versions of the manuscript. All authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/JSRR/2019/v23i130112

#### Editor(s):

(1) Dr. Ana Ribeiro-Barros, Biotrop Center - Environment, Agriculture and Development, Tropical Research Institute (IICT), Portugal.

(2) Dr. Khadiga Ahmed Ismail, Parasitology Department, Faculty of Medicine, Ain Shams University, Cairo, Egypt.

(3) Dr. Rahul Kumar Jaiswal, National Institute of Hydrology, WALMI Campus, Bhopal, India.

#### Reviewers:

(1) Tiogué Tekounegning Claudine, The University of Dschang, Cameroon.

(2) Ali Türker Mugla, Sıtkı Koçman University, Turkey.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/47885>

Received 12 January 2019

Accepted 30 March 2019

Published 10 April 2019

Original Research Article

### ABSTRACT

Reproduction of the threadfins of the costal shelf of Grand-Lahou was studied with a monthly sampling frequencies range from may 2009 to april 2011. 534 specimens of *Polydactylus quadrifilis*, composed of 315 males and 219 females and 648 *Galeoides decadactylus* composed of 420 males and 228 females, then 453 individuals of *Pentanemus quinquarius* composed of 210 males and 243 females were used.

\*Corresponding author: E-mail: [kokouasy@yahoo.fr](mailto:kokouasy@yahoo.fr);

The sexual maturity parameters of threadfins fishes indicate a size of first maturity ( $L_{50} = 67.5$  cm) for females of *Polydactylus quadrifilis* and ( $L_{50} = 55.5$  cm) with their males. Concerning the specimens of *Pentanemus quinquarius*, females present ( $L_{50} = 15$  cm) and their males ( $L_{50} = 14.5$  cm). This parameter has identical values with the females of *Galeoides decadactylus* ( $L_{50} = 15.5$  cm) and their males ( $L_{50} = 15.5$  cm). The individuals of *Polydactylus quadrifilis* and the specimens of *Galeoides decadactylus* and *Pentanemus quinquarius* become mature at 1 year of age. The sex ratio determination show that specimens of *Polydactylus quadrifilis* show a predominance of males during the warm season, from October (73.68%) to March (64.28%) which weakens during the cold season, from May (30%) to July (58%). For *Galeoides decadactylus*, the sex ratio shows a predominance of males, from August (88%) to March (94.73%). Concerning specimens of *Pentanemus quinquarius*, the sex ratio is in favor of the females whose lowest value is during the cold season in July (21.74%). All threafin species reach sexual maturity from their first year of life with different sex ratios variations.

**Keywords:** *Reproduction; sexual maturity; sex ratio; polynemidae; Ivory Coast.*

## 1. INTRODUCTION

Fish are the main source of animal protein consumed worldwide, especially in many developing countries. They also ensure food security for people living near watercourses [1]. To increase national production, States define sectoral policies involving the development of marine and lagoon fisheries, by the rational exploitation of all fishing potential, the modernization of artisanal production means, and the pursuit of efforts in fisheries research [2]. According to FAO [3], promoting sustainable fisheries can encourage better management of the whole ecosystem.

In fact, the exploitation of brackish and marine water fish populations in Africa is intensifying more and more by constantly increasing local populations and especially the alarming acceleration of all the processes of degradation of the natural environment that make consider the major risk of regression and disappearance of species [4]. One of the most significant effects of fishing is in terms of demography, reduction of the average size of the species and the disappearance of large individuals [5].

Therefore, the knowledge of the reproductive biology of the main species among others, threadfins fishes, *Polydactylus quadrifilis*, *Galeoides decadactylus* and *Pentanemus quinquarius* landed on the Ivorian coast is important.

The present study consists in determining the sexual maturity of these threadfins fishes of the ivorian maritime artisanal fishery in order to

contribute to the preservation of the stocks. The sexual maturity have been studied through the size of sexual maturity, the age of first maturation and the sex-ratio of each threadfin specie.

## 2. MATERIALS AND METHODS

Reproduction was studied by conducting monthly samplings within the catches made by the sea fishermen of Grand-lahou. It involved 534 specimens of *Polydactylus quadrifilis* including 315 males and 219 females and 648 of *Galeoides decadactylus* composed of 420 males and 228 females, then 453 of *Pentanemus quinquarius* composed of 210 males and 243 females. Sampling frequencies range from May 2009 to April 2011, for a total of 24 months. At each sampling campaign, a sampling by sex and size class of one cm was carried out. The fish were kept in coolers and brought back to the laboratory. The total lengths ( $L_t$ ) were taken and was used for calculation of the first sexual maturity. Sexes have been identified after dissection of each fish individuals. The proportions of males and females were calculated in relation to the total size within the different size classes.

### 2.1 Size of First Sexual Maturity ( $L_{50}$ )

To calculate the size at which 50% of individuals are mature, individuals of each species were ranked in size classes of one cm. In each class, the percentage of fish whose macroscopic stage of sexual maturity is higher than or equal to stage 3 has been calculated, a stage admitted as corresponding to the stage of ovarian development [6,7].

The logistic function  $P = 1/(1+e^{-(b+aL)})$  linking the proportions of mature individuals and the total length of fish [8] was used.

P: % of mature, a and b: constants,  $L_t$ : total length of fish

## 2.2 Age of First Maturation

According to [9], gonad maturation is not achieved at the same age for all individuals in a cohort. The proportion of maturing gonads increases with age from zero to 100% and all individuals are mature from the age at which 100% spawn or spermate for the first time. The histogram or curve that represents these proportions is called the maturation ogive. It determines the age of first maturation ( $t_{mat}$ ) by calculating the proportions of fish whose macroscopic stage of sexual maturity is greater than or equal to 3 in size classes of one cm.

Thus, specimens of age below to  $t_{mat}$  are considered juveniles and those of equal or higher age are considered as adults [9]. Calculations of the proportions of fish with a macroscopic stage of sexual maturity greater than or equal to 3, by size classes of one cm, made it possible to determine the age of first maturation ( $t_{mat}$ ) of the threadfins.

## 2.3 Sex-Ratio

The sex ratio reflects the masculinity or femininity rate of the population considered. It is defined as the proportion of male or female individuals respectively in relation to the size of females or males and gives an idea of the gender balance (males and females) within the population.

Sex ratio =  $M \times 100 / F$

with; F: number of females and M: number of males

## 3. RESULTS AND DISCUSSION

### 3.1 Size of First Sexual Maturity ( $L_{50}$ )

The logistic model equations determined for the different sexes of the Polynemidae were presented in the table 1. Fig. 1 and 2 respectively showed the graphical representations of the sizes of first sexual maturity of the males and females threadfins. The first maturity sizes obtained with *Polydactylus quadrifilis* were ( $L_{50} = 67.5$  cm) for females and ( $L_{50} = 55.5$  cm) for

males. About the *Pentanemus quinquarius* specimens, females had the size of first sexual maturity ( $L_{50} = 15$  cm) while males had a size of first sexual maturity ( $L_{50} = 14.5$  cm). The size of first sexual maturity were identical with females of *Galeoides decadactylus* ( $L_{50} = 15.5$  cm) and their males ( $L_{50} = 15.5$  cm).

### 3.2 Age of First Maturation

*Polydactylus quadrifilis* individuals became mature at 1 year of age (Fig. 3.a). It was the same for the specimens of *Galeoides decadactylus* (Fig. 3.b) and *Pentanemus quinquarius* (Fig. 3.c).

### 3.3 Sex-Ratio

For the specimens of *Polydactylus quadrifilis* (Fig. 4.a), the sex ratio indicated a predominance of males during the period from October (73.68%) to March (64.28%), with a peak of 83, 33% in November. This sex ratio became low during the cold season, from May (30%) to the end of July (58%). According to *Galeoides decadactylus*, the sex ratio showed a predominance of males, from August (88%) to March (94.73%), except for the month of October where females (88.88%) dominate males (11.11%) (Fig. 4.b). Concerning specimens of *Pentanemus quinquarius*, the sex ratio was in favor of the females whose lowest value were during the cold season, in July (21.74%) (Fig. 4.c).

In order to understand the maturation of threadfins on the costal shelf of Grand-Lahou during their reproductive cycle, this study looked at the size and age corresponding to their first sexual maturity and their sex ratio.

According to Dadebo et al. [10], knowledge of the size of first sexual maturity is important in the management of fisheries resources. It is essential in determining the minimum catch size. In our study, the size of first sexual maturity was reached at 55.5 cm with males and 67.5 cm for females of *Polydactylus quadrifilis*. About of specimens of *Galeoides decadactylus*, males and females simultaneously reached the size of first sexual maturity at 15.5 cm. As well, males and females of *Pentanemus quinquarius* had respectively a size of first sexual maturity ( $L_{50} = 14.5$  cm) and ( $L_{50} = 15$  cm). Compared with the types of allometries observed with threadfin fishes, differential growth did not influence

reaching the size of first sexual maturity. Indeed, specimens of *Polydactylus quadrifilis* and *Pentanemus quinquarius*, those didn't showed differential growth, had different sizes of first sexual maturity between males and females. In contrast, specimens of *Galeoides decadactylus* that showed differential growth between their two sexes, reached simultaneously the size of first sexual maturity [11]. Similarly, [12] report that males frequently dominated catches of juveniles because they was younger but live shorter lives. Thus, sizes of first sexual maturity ( $L_{50} = 15.5$  cm) identical to our results were determined by [13] on the Nigerian coasts with males and

females of *Galeoides decadactylus*. However, results different from ours were obtained by Samba [14] ( $L_{50} = 18.5$  cm) and [15] ( $L_{50} = 13.7$  cm) with *Galeoides decadactylus*, respectively on the Congolese and Senegalese coasts. Similarly, [16] observed lower first maturity sizes ( $L_{50} = 12$  cm) with *Galeoides decadactylus* at the Guinean littoral. In addition, [7] have obtained the fork size of  $L_{50} = 13$  cm with *Galeoides decadactylus* specimens of the Guinean coastal sea. The sizes of first maturity determined with threadfin fishes was corresponded to one year of age. That means these fish were able to contribute to restocking from one year of age.





Fig. 1. First sexual maturity length of males of *Polydactylus quadrifilis* (a), *Galeoides decadactylus* (b) and *Pentanemus quinquarius* (c)



**Fig. 2. First sexual maturity length of females of *Polydactylus quadrifilis* (a), *Galeoides decadactylus* (b) and *Pentanemus quinquarius* (c)**

The size of first sexual maturity would therefore be a function of the physicochemical conditions and the availability in primary productions of their living environments. [17] have noticed the idea that reproduction is closely related to the adaptability of breeders to certain factors in their environment, such as temperature, salinity and trophic resources. This would justify size dimorphism between males and females of *Polydactylus quadrifilis* and *Pentanemus quinquarius*. According to Reichert [18], the best

growth rates were linked to the quality of the diet adopted by the fish and their ability to adapt to the temperature of the environment. Du Buit [19] added that temperature could promote metabolic activities and accelerates growth and aging of animals.

The sex ratio of *Polydactylus quadrifilis* specimens indicated a predominance of males during the warm season and becomes low during the cold season. However, the sex ratio remains

balanced between males and females during the cold season. According to *Galeoides decadactylus* individuals, the sex ratio showed a predominance of males throughout the fishing season. Concerning the specimens of *Pentanemus quinquarius*, the sex ratio were in favor of females. The sex ratio were favorable for males of *Polydactylus quadrifilis* and *Galeoides*

*decadactylus*. However, it was in favor of females with the specimens of *Pentanemus quinquarius*. According to Lopez [15] and Sidibe [7], respectively at the Senegalese and Guinean seacoasts, there were a sex inversion of the young males of *Galeoides decadactylus* during their life cycle. It would be a gradual transformation of males into females until




Fig. 3. First sexual maturity age of threadfins *Polydactylus quadrifilis* (a) *Galeoides decadactylus* (b) and *Pentanemus quinquarius* (c)




Fig. 4. Monthly variation of male (M) and female (F) proportions of thredfins *Polydactylus quadrifilis* (a), *Galeoides decadactylus* (b) and *Pentanemus quinquarius* (c)

Table 1. Size of first sexual maturity (L50) and equations of logistic function P of thredfins *Polydactylus quadrifilis*, *Galeoides decadactylus* and *Pentanemus quinquarius*

Species	Sex	Effectives	L <sub>50</sub> (cm)	P Equations
<i>Polydactylus quadrifilis</i>	Male	315	55,5	$P = \frac{1}{1 + e^{(-7,44 + 2,02Lt)}}$
	Female	219	67,5	$P = \frac{1}{1 + e^{(-1,18 + 0,67Lt)}}$
<i>Galeoides decadactylus</i>	Male	420	15,5	$P = \frac{1}{1 + e^{(-7,48 + 2,41Lt)}}$
	Female	228	15,5	$P = \frac{1}{1 + e^{(-7,93 + 2,49Lt)}}$

Species	Sex	Effectives	L <sub>50</sub> (cm)	P Equations
	Male	210	14,5	$P = \frac{1}{1 + e^{(-3,64 + 1,54Lt)}}$
<i>Pentanemus quinquarius</i>	Female	243	15	$P = \frac{1}{1 + e^{(-6,32 + 2,17Lt)}}$

hermaphroditism were functional. This phenomenon would result in a favorable sex ratio for females. These results was contrary to ours because the sex ratio obtained with *Galeoides decadactylus* was rather favorable to the males. Sidibe [7] argues that changes in sex ratio and size frequency probably had a significant influence on stock availability. The sex ratio in favor of the males results from the fact that the females of *Polydactylus quadrifilis* and *Galeoides decadactylus*, according to Motomura [20], would withdraw from the fishing grounds, to take refuge in marine vegetation 50 meters deep and berries to lay. These females would avoid the fish ermen when those do not fish at shorelines and bays.

#### 4. CONCLUSION

This study shows that specimens of *Polydactylus quadrifilis* and *Pentanemus quinquarius* have different sizes of first sexual maturity between males and females. However, the males and females of *Galeoides decadactylus* simultaneously reach the size of first sexual maturity. Specimens of *Polydactylus quadrifilis* show a predominance of males during the warm season, females during the cold season. Males of *Galeoides decadactylus* are predominant while females predominate in specimens of *Pentanemus quinquarius*. Individuals of *Polydactylus quadrifilis*, *Galeoides decadactylus* and *Pentanemus quinquarius* become mature at one year of age.

#### ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

#### ACKNOWLEDGEMENTS

Authors acknowledge authorities of the Oceanologic Research Centre, the Peleforo Gon Coulibaly University and the Nangui Abrogoua University. They thank laboratory of the aquaculture department of the Oceanologic Research Centre.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. FAO. Accroissement de la contribution des pêches artisanales à la lutte contre la pauvreté et à la sécurité alimentaire. FAO Rome. 2006;108.
2. FPA. Contrat-cadre pour la réalisation d'évaluation, d'étude d'impact et de suivi, concernant les accords de partenariat dans le domaine de la pêche conclus entre la communauté et les pays tiers, étude du secteur de la pêche en Côte d'Ivoire et analyse ex-ante du protocole 2003-2004, rapport final mai. 2004;150.
3. FAO. La Situation Mondiale des Pêches et de l'Aquaculture. FAO Rome. 2012; 241.
4. Laleye P, Chikou A, Philippart JC, Teugels G, Yandewalle P. Étude de la diversité ichtyologique du bassin du fleuve Ouémé au Bénin, Cybium. 2004;329-339.
5. Kraidy LAB, Kone N, Berte S, Konan GN, Yao SS, Kouamelan PE. Pêche et paramètres de reproduction de *Pellonula leonensis*, dans le lac de Taabo (Fleuve Bandama, Côte d'Ivoire): Implications pour une exploitation durable du stock, Int. J. Biol. Chem. Sci. 2014;8(1):75-88.
6. Fehri-Bedoui R, Gharbi H, El Abed A. Période de reproduction et maturité sexuelle de *Liza aurata* des côtes est et sud tunisiennes. Bulletin de l'institut national des sciences et techniques de la mer de Salanumbô. 2002;29:43-56.
7. Sidibe A. Les ressources halieutiques démersales côtières de la guinée: exploitation, biologie et dynamique des principales espèces de la communauté à sciaénidés. Thèse de Doctorat Halieutique ENSA-Rennes. 2003;320.
8. Ghorbel AO, Bradai MN, Bouain A. Période de reproduction et maturité sexuelle de *symphodus (Crenilabrus) tinca* (labridae), des côtes des sfax (tunisie)



- bulletin de l'Institut National des Sciences et Techniques de la Mer de Salanumbô*. 2002;30:76-89.
9. Cadima EL. Manuel d'évaluation des ressources halieutiques. FAO, document technique sur les pêches. Rome. 2002; 291.
  10. Dadebo E, Ahlgrea G, Ahlgreen L. Aspect of reproductive biology of *Labeo horie* (Pisces; Cyprinidae) in Lake Chamo. African Journal of Ecology. 2003;41:31-38.
  11. Konan KS, Diaby M, Agnissan AJP, Kone A, N'da K. Croissance et âge des poissons capitaines: *Polydactylus quadrifilis* (Cuvier, 1829), *Galeoides decadactylus* (Bloch, 1795) et *Pentanemus quinquarius* (Linné, 1758) de la pêche artisanale maritime de Grand-Lahou (Côte d'Ivoire). Int. J. Biol. Chem. Sci. 2012;6(3):1112-1127. [ISSN 1991-8631]
  12. Ozcan G, Balik S. Age and growth of Bassan barbel, *Barbus pectoralis* (Actinopterygii: Cypriniformes: Cyprinidae), under conditions of a dam reservoir. ACTA Ichthyologica et Piscatoria. 2009;39:27-32.
  13. Longhurst AR. Bionomics of the Sciaenidae of tropical west Africa. J. Cons. Inter. Explor. Mer. 1964;1:93-114.
  14. Samba G. Contribution à l'étude de la biologie et de la dynamique d'un Polynemidae ouest africain *Galeoides decadactylus* (Boch., 1795). Thèse de 3e cycle, Université de Bordeaux I, France. 1974;114.
  15. Lopez J. Biologie de la reproduction de *Galeoides decadactylus* au Sénégal. Document Scientifique Prov. Centre de Recherche Océanographique Dakar-Thiaroye. 1979;68:191-204.
  16. Domain F, Bah MO. Description des fonds du plateau continental. In: La pêche côtière en Guinée – Ressources et Exploitation. Domain F, Chavance P, Diallo A. (Eds.), Editions IRD/CNSHB, Paris. 2000;37-49.
  17. Poulet N. Le sandre (*Sander lucioperca* L.): Biologie, comportement et dynamique des populations en Camargue (Bouches du Rhône, France). Thèse de Doctorat de l'Université de Toulouse III. 2004; 185.
  18. Reichert MJM. Diet, consumption, and growth of juvenile flounder (*Etropus crossotus*); A test of the "maximum growth/optimum food hypothesis" in a subtropical nursery area. Journal of sea Research. 2003;50:97-116.
  19. Du Buit MH. Alimentation du merlan *Merlangius merlangus* L. en mer Celtique. Revue Travaux de l'Institut des Pêches maritimes. 1987;49:5-12.
  20. Motomura K, Iwatsuki Y. Threadfins of the World, FAO Species Catalogue for Fishery Purposes. 2002;3:65-71.

© 2019 Konan et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<http://www.sdiarticle3.com/review-history/47885>